

**DYE FREE PAIN RELIEVER CHILDRENS- acetaminophen suspension**  
**Safeway, Inc.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**Drug Facts**

**Active ingredient (in each 5 mL)**

**Acetaminophen 160 mg**

**Purpose**

**Pain reliever/fever reducer**

**Uses**

- temporarily reduces fever
- temporarily relieves minor aches and pains due to:
  - the common cold
  - flu
  - headache
  - sore throat
  - toothache

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or non-prescription). If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if your child has**

liver disease.

**Ask a doctor or pharmacist before use if your child is**

taking the blood thinning drug warfarin.

**When using this product,**

**do not exceed recommended dosage (see overdose warning)**

**Stop use and ask a doctor if**

- new symptoms occur
- redness or swelling is present
- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days

These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

**Directions**

- **this product does not contain directions or complete warnings for adult use**
- **do not take more than directed (see overdose warning)**
- **shake well before using**
- mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device.
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give more than 5 days unless directed by a doctor

<b>Weight (lb)</b>	<b>Age (year)</b>	<b>Dose (mL)*</b>
under 24	under 2	ask a doctor
24-35	2-3	5 mL
36-47	4-5	7.5 mL
48-59	6-8	10 mL

60-71	9-10	12.5 mL
72-95	11	15 mL

\*or as directed by a doctor

### **Other information**

- **each 5 mL contains:**sodium 2 mg
- store between 20-25°C (68-77°F)
- do not refrigerate
- see bottom panel for lot number and expiration date

### **Inactive ingredients**

butylparaben, carboxymethylcellulose sodium, citric acid, flavor, glycerin, high fructose corn syrup, microcrystalline cellulose, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum

### **Questions or comments?**

Call **1-888-723-3939 Monday-Friday 7AM-6PM PST**

### **Principal Display Panel**

FOR AGES 2-11 YEARS

### **Children's Dye-Free Pain Reliever**

Acetaminophen 160 mg per 5 mL

Pain Reliever / Fever Reducer

Oral Suspension

- Alcohol free
- Aspirin free
- Ibuprofen free
- Dosage cup included

GRAPE FLAVOR

FL OZ (mL)

\*\*Compare to Children's Tylenol® Oral Suspension active ingredient

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.**

\*\*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Tylenol® Oral suspension.

DISTRIBUTED BY:

BETTER LIVING BRANDS LLC

P.O. BOX 99, PLEASANTON, CA 94566-0009

www.betterlivingbrandsLLC.com

**Product Label**

**Drug Facts (continued)**

- use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device.
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give more than 5 days unless directed by a doctor
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.

Weight (lb)	Age (year)	Dose (mL)*
under 24	under 2	ask a doctor
24-35	2-3	5 mL
36-47	4-5	7.5 mL
48-59	6-8	10 mL
60-71	9-10	12.5 mL
72-85	11	15 mL

\*or as directed by a doctor

**Other information**

- each 5 mL contains: sodium 2 mg
- store between 20-25°C (68-77°F) do not refrigerate
- see bottom panel for lot number and expiration date

**Inactive ingredients** butylparaben, carboxymethylcellulose sodium, citric acid, flavors, glycerin, high fructose corn syrup, microcrystalline cellulose, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum

**Questions or comments?**  
Call 1-888-723-3929 Monday-Friday 7AM-6PM PST

\*\*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Tylenol® Oral Suspension.

**TAMPER EVIDENT. DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.**

DISTRIBUTED BY  
BETTER LIVING BRANDS LLC  
P.O. BOX 99, PLEASANTON, CA 94566-0009  
1-888-723-3929  
www.betterlivingbrandsLLC.com

**OUR PROMISE**  
QUALITY & SATISFACTION  
**100% GUARANTEED**  
OR YOUR MONEY BACK.

NDC 21130-285-04



Quality Guaranteed

FOR AGES 2-11 YEARS

**Children's Dye-Free Pain Reliever**

Acetaminophen  
160 mg per 5 mL

Oral Suspension  
Pain Reliever/Fever Reducer



GRAPE FLAVOR

\*\*Compare to Children's Tylenol® Oral Suspension active ingredient

- Alcohol Free
- Aspirin Free
- Ibuprofen Free
- Dosing cup included

4 FL OZ (118 mL)

NDC 21130-285-04



Quality Guaranteed

FOR AGES 2-11 YEARS

**Children's Dye-Free Pain Reliever**

Acetaminophen  
160 mg per 5 mL

Oral Suspension  
Pain Reliever/Fever Reducer



GRAPE FLAVOR

\*\*Compare to Children's Tylenol® Oral Suspension active ingredient

- Alcohol Free
- Aspirin Free
- Ibuprofen Free
- Dosing cup included

4 FL OZ (118 mL)

**Drug Facts**

Active ingredient (in each 5 mL)	Purpose
Acetaminophen 160 mg	Pain reliever/fever reducer

**Uses**

- temporarily reduces fever
- temporarily relieves minor aches and pains due to
  - the common cold
  - flu
  - headache
  - sore throat
  - toothache

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if your child has liver disease.**

**Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.**

**When using this product, do not exceed recommended dose (see Overdose warning).**

**Stop use and ask a doctor if**

- new symptoms occur
- redness or swelling is present
- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days

These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

**Directions**

- this product does not contain directions or complete warnings for adult use
- do not take more than directed (see Overdose warning)
- shake well before using
- mL = milliliter



PLD-C234D FC004027

Lot No:  
Exp. Date:

**SIGNATURE CARE Children's Dye-Free Pain Reliever Grape Flavor**

**DYE FREE PAIN RELIEVER CHILDRENS**

acetaminophen suspension

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:21130-285
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

**Inactive Ingredients**

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
BUTYLPARABEN (UNII: 3QPI1U3FV8)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

**Product Characteristics**

Color		Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-285-04	1 in 1 BOX	06/30/2015	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	06/30/2015	

**Labeler** - Safeway, Inc. (009137209)