

CLARITIN REDITABS- loratadine tablet, orally disintegrating
Bayer HealthCare LLC

Claritin®

RediTabs®

Drug Facts

Active ingredient (in each tablet)

Loratadine 5 mg

Purpose

Antihistamine

Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- itchy, watery eyes
- sneezing
- itching of the nose or throat

Warnings

Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.

When using this product do not take more than directed. Taking more than directed may cause drowsiness.

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- place 1 tablet on tongue; tablet disintegrates, with or without water

| | |
|--|--|
| adults and children 6 years and over | 1 tablet every 12 hours; not more than 2 tablets in 24 hours |
| children under 6 years of age | ask a doctor |
| consumers with liver or kidney disease | ask a doctor |

Other information

- safety sealed: do not use if the individual blister unit imprinted with Claritin® RediTabs® is open or torn
- store between 20° to 25°C (68° to 77°F)
- use tablet immediately after opening individual blister
- complies with USP Procedure 2 for Assay and Organic Impurities and Test 2 for Disintegration

Inactive ingredients

anhydrous citric acid, gelatin, mannitol, mint flavor

Questions or comments?

1-800-CLARITIN (1-800-252-7484) or www.claritin.com

Distributed by MSD Consumer Care, Inc.,
PO Box 377, Memphis, TN 38151 USA,
a subsidiary of Merck & Co., Inc., Whitehouse Station, NJ USA.

PRINCIPAL DISPLAY PANEL - 10 Tablet Carton

NDC 11523-7202-3

Non-Drowsy*

Claritin®

RediTabs®

loratadine 5 mg/antihistamine

Indoor & Outdoor

Allergies

****When taken as directed. See Drug Facts Panel.***

Hour

Relief of:

- Sneezing
- Runny Nose
- Itchy, Watery Eyes
- Itchy Throat

or Nose

No Water Needed

Melts in Your Mouth

10

ORALLY

DISINTEGRATING TABLETS

Claritin[®] Reditabs[®] **12 Hour**

Non-Drowsy*

Claritin[®]

Reditabs[®]

loratadine 5 mg/antihistamine

Indoor & Outdoor Allergies

No Water Needed Melts in Mouth

ORALLY DISINTEGRATING TABLETS

BAYER

* When taken as directed. See Drug Facts Panel.



Claritin® Reditabs®
12 Hour

Drug Facts (continued)

Other Information

- safety sealed: do not use if the individual blister unit is open or torn
- with Claritin® Reditabs® is open or torn
- store between 20° to 25° C (68° to 77° F)
- use tablet immediately after opening individual blister
- use tablet immediately after opening individual blister
- complies with USP Procedure 2 for Assay and Organic Impurities and Test 2 for Disintegration

Inactive Ingredients

anhydrous citric acid, gelatin, mannitol, mint flavor

Questions or comments?
1-800-CLARITIN (1-800-252-7484) or www.claritin.com

Purpose Antihistamine

(in each tablet)

eyes these symptoms due to hay fever or allergies:
 ■ itchy, watery eyes
 ■ itching of the nose or throat

or had an allergic reaction to this product or

If you have liver or kidney disease. Your doctor may need a different dose.

do not take more than directed. Taking more may increase the risk of drowsiness.

If an allergic reaction to this product occurs, stop taking Claritin and ask a health professional before use. In case of overdose, get medical help or call a poison control center right away.

1 tablet every 12 hours; with or without water

not more than 2 tablets in 24 hours

ask a doctor

ask a doctor

1. Peel back outer edge.

2. Gently push tablet out.

3. Place the tablet on tongue and close mouth. The tablet will disintegrate.

Follow these directions carefully. Do not attempt to push the tablet through the foil.

• No Water Needed • Melts in your mouth

CLARITIN REDITABS

loratadine tablet, orally disintegrating

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|----------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:11523-7202 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------|----------|
| LORATADINE (UNII: 7AJ03BO7QN) (LORATADINE - UNII:7AJ03BO7QN) | LORATADINE | 5 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) | |
| GELATIN (UNII: 2G86QN327L) | |
| MANNITOL (UNII: 3OWL53L36A) | |

Product Characteristics

| | | | |
|---------------|---------------------------------|---------------------|----------|
| Color | white | Score | no score |
| Shape | ROUND (flat faced beveled edge) | Size | 12mm |
| Flavor | MINT | Imprint Code | C5 |

Contains**Packaging**

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:11523-7202-1 | 3 in 1 CARTON | 12/12/2006 | |
| 1 | | 10 in 1 BLISTER PACK; Type 0: Not a Combination Product | | |
| 2 | NDC:11523-7202-3 | 1 in 1 CARTON | 12/12/2006 | |
| 2 | | 10 in 1 BLISTER PACK; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| NDA | NDA021993 | 12/12/2006 | |

Labeler - Bayer HealthCare LLC (112117283)

Revised: 1/2023

Bayer HealthCare LLC