

ASPIRIN LOW DOSE- aspirin tablet, chewable
ARMY AND AIR FORCE EXCHANGE SERVICE

Exchange Select 44-218

Active ingredient (in each chewable tablet)

Aspirin 81 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

for the temporary relief of minor aches and pains or as recommended by your doctor

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- new symptoms occur
- pain gets worse or lasts more than 10 days
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- drink a full glass of water with each dose
- chew or crush tablets completely before swallowing; do not swallow tablets whole
- adults and children 12 years and over: take 4 to 8 chewable tablets every 4 hours not to exceed 48 chewable tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, dextrates hydrated, FD&C yellow #6 aluminum lake, flavor, maltodextrin, sucralose

Questions or comments?

1-800-426-9391

Principal Display Panel

exchange ✓ **select™**

Compare To The Active Ingredient of
Bayer® Chewable Low Dose Aspirin†

Talk to your Doctor
or other Healthcare
provider before
using this product
for your heart.

LOW DOSE ASPIRIN Chewable Tablets

Pain Reliever (**NSAID**)

•Aspirin Regimen •Orange flavored

81 mg

Actual
Size

Chew or crush tablets completely
before swallowing.
Do not swallow tablets whole.

3 BOTTLES INSIDE
36 + 36 + 36 Tablets

✓quality
value

Value Pack
3 bottles x 36 tablets each =
108 tablets

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

†This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Chewable Low Dose Aspirin.

50844 REV1118A21807

“SATISFACTION GUARANTEED OR YOUR MONEY BACK”

Manufactured For Your Military Exchanges

Distributed by: LNK International, Inc.

Hauppauge, NY 11788

1-800-426-9391

exchange select™

LOW DOSE ASPIRIN Chewable Tablets

Compare To The Active Ingredient of
Bayer® Chewable Low Dose Aspirin†

Talk to your Doctor
or other Healthcare
provider before
using this product
for your heart.

Pain Reliever (NSAID)

- Aspirin Regimen • Orange Flavored

 **3 BOTTLES
INSIDE**
36 + 36 + 36 Tablets

Chew or crush tablets completely
before swallowing.
Do not swallow tablets whole.



Actual
Size

quality
value

exchange select™

LOW DOSE ASPIRIN Chewable Tablets

Compare To The Active Ingredient of
Bayer® Chewable Low Dose Aspirin†

Talk to your Doctor
or other Healthcare
provider before
using this product
for your heart.

Pain Reliever (NSAID)

- Aspirin Regimen • Orange Flavored

 **3 BOTTLES
INSIDE**
36 + 36 + 36 Tablets

Chew or crush tablets completely
before swallowing.
Do not swallow tablets whole.



Actual
Size

quality
value

**Value
Pack**
3 bottles x 36 tablets each =
108 tablets

Value Pack

3 bottles x 36 tablets each =
108 tablets

exchange select™

LOW DOSE ASPIRIN Chewable Tablets

Compare To The Active Ingredient of
Bayer® Chewable Low Dose Aspirin†

Talk to your Doctor
or other Healthcare
provider before
using this product
for your heart.

Pain Reliever (NSAID)

- Aspirin Regimen • Orange Flavored

 **3 BOTTLES
INSIDE**
36 + 36 + 36 Tablets

Chew or crush tablets completely
before swallowing.
Do not swallow tablets whole.



Actual
Size

quality
value

exchange select™

**LOW DOSE
ASPIRIN
Chewable Tablets**



TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

†This product is not manufactured or distributed by Bayer AG, owner of the registered
trademark Bayer® Chewable Low Dose Aspirin.
50844 REV1118A21807
"SATISFACTION GUARANTEED OR YOUR MONEY BACK"
Manufactured For Your Military Exchanges
Distributed by: LNK International, Inc.
Hauppauge, NY 11788
1-800-426-9391

B-0066E-218A-07BR
REV1118A21807

Questions or comments?
1-800-426-9391

as medication
■ you have high blood pressure, heart disease,
liver cirrhosis, or kidney disease
■ you have asthma ■ you are taking a diuretic

Drug Facts	
Active ingredient (in each chewable tablet) Aspirin 81 mg (NSAID) Pain reliever nonsteroidal anti-inflammatory drug	Purpose
Uses for the temporary relief of minor aches and pains or as recommended by your doctor	
Warnings Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] ■ have had stomach ulcers or bleeding problems ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed	
Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor ■ if you have ever had an allergic reaction to this product or any of its ingredients Ask a doctor before use if ■ you have a history of stomach problems, such as stomach bleeding warning applies to you ■ you have a history of ulcers, gastritis, or kidney disease	
Drug Facts	
Active ingredient (in each chewable tablet) Aspirin 81 mg (NSAID) Pain reliever nonsteroidal anti-inflammatory drug	Purpose
Uses for the temporary relief of minor aches and pains or as recommended by your doctor	
Warnings Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] ■ have had stomach ulcers or bleeding problems ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed	
Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor ■ if you have ever had an allergic reaction to this product or any of its ingredients Ask a doctor before use if ■ you have a history of stomach problems, such as stomach bleeding warning applies to you ■ you have a history of ulcers, gastritis, or kidney disease	
Other information ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ see end flap for expiration date and lot number Inactive ingredients corn starch, croscarmellose, hydroxypropyl methylcellulose, polyethylene glycol, polyvinylpyrrolidone, sodium lauryl sulfate, titanium dioxide, yellow #6 aluminum lake.	
Directions ■ drink a full glass of water with each dose ■ chew or crush tablets completely before swallowing; do not swallow tablets whole ■ adults and children 12 years and over: take 4 to 8 chewable tablets every 4 hours not to exceed 48 chewable tablets in 24 hours unless directed by a doctor ■ children under 12 years: do not use unless directed by a doctor	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.	
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.	
Directions ■ drink a full glass of water with each dose ■ chew or crush tablets completely before swallowing; do not swallow tablets whole ■ adults and children 12 years and over: take 4 to 8 chewable tablets every 4 hours not to exceed 48 chewable tablets in 24 hours unless directed by a doctor ■ children under 12 years: do not use unless directed by a doctor	
Other information ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ see end flap for expiration date and lot number Inactive ingredients corn starch, croscarmellose, hydroxypropyl methylcellulose, polyethylene glycol, polyvinylpyrrolidone, sodium lauryl sulfate, titanium dioxide, yellow #6 aluminum lake.	
Drug Facts	
Active ingredient (in each chewable tablet) Aspirin 81 mg (NSAID) Pain reliever nonsteroidal anti-inflammatory drug	Purpose
Uses for the temporary relief of minor aches and pains or as recommended by your doctor	
Warnings Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] ■ have had stomach ulcers or bleeding problems ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed	
Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor ■ if you have ever had an allergic reaction to this product or any of its ingredients Ask a doctor before use if ■ you have a history of stomach problems, such as stomach bleeding warning applies to you ■ you have a history of ulcers, gastritis, or kidney disease	
Other information ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ see end flap for expiration date and lot number Inactive ingredients corn starch, croscarmellose, hydroxypropyl methylcellulose, polyethylene glycol, polyvinylpyrrolidone, sodium lauryl sulfate, titanium dioxide, yellow #6 aluminum lake.	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.	
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.	
Directions ■ drink a full glass of water with each dose ■ chew or crush tablets completely before swallowing; do not swallow tablets whole ■ adults and children 12 years and over: take 4 to 8 chewable tablets every 4 hours not to exceed 48 chewable tablets in 24 hours unless directed by a doctor ■ children under 12 years: do not use unless directed by a doctor	
Other information ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ see end flap for expiration date and lot number Inactive ingredients corn starch, croscarmellose, hydroxypropyl methylcellulose, polyethylene glycol, polyvinylpyrrolidone, sodium lauryl sulfate, titanium dioxide, yellow #6 aluminum lake.	
Drug Facts	
Active ingredient (in each chewable tablet) Aspirin 81 mg (NSAID) Pain reliever nonsteroidal anti-inflammatory drug	Purpose
Uses for the temporary relief of minor aches and pains or as recommended by your doctor	
Warnings Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] ■ have had stomach ulcers or bleeding problems ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed	
Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor ■ if you have ever had an allergic reaction to this product or any of its ingredients Ask a doctor before use if ■ you have a history of stomach problems, such as stomach bleeding warning applies to you ■ you have a history of ulcers, gastritis, or kidney disease	
Other information ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ see end flap for expiration date and lot number Inactive ingredients corn starch, croscarmellose, hydroxypropyl methylcellulose, polyethylene glycol, polyvinylpyrrolidone, sodium lauryl sulfate, titanium dioxide, yellow #6 aluminum lake.	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.	
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.	
Directions ■ drink a full glass of water with each dose	

aspirin tablet, chewable

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients	
Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
DEXTROSE MONOHYDRATE (UNII: LX22YL083G)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	

SUCRALOSE (UNII: 96K6UQ3ZD4)				
Product Characteristics				
Color	orange	Score	no score	
Shape	ROUND	Size	8mm	
Flavor	ORANGE	Imprint Code	44;218	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55301-821-07	3 in 1 CARTON	09/12/1990	
1		36 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013		09/12/1990	

Labeler - ARMY AND AIR FORCE EXCHANGE SERVICE (001695568)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(55301-821)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(55301-821)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(55301-821)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(55301-821)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(55301-821)

