BAYER- as pirin tablet Lil' Drug Store Products, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

BAYER® ASPIRIN

Drug Facts

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)¹

1 nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

temporarily relieves

- headache
- muscle pain
- toothache
- menstrual pain
- pain and fever of colds
- minor pain of arthritis

Warnings

Reve's syndrome

Children and teenagers who have or are recovering form chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- redness or swelling is present
- fever lasts more than 3 days
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours
- children under 12 years: consult a doctor

Other information

- save carton for full directions and warnings
- store at room temperature

Inactive ingredients

2 may contain this ingredient

Questions or comments?

1-800-331-4536 or www.bayeraspirin.com manufactured by Bayer Corporation

PRINCIPAL DISPLAY PANEL - 4 Tablet Carton

GENUINE ASPIRIN BAYER®
ASPIRIN (NSAID) PAIN RELIEVER/FEVER REDUCER

THE WONDER DRUG®

325^{mg}

Safe Pain Relief Plus Lifesaving Benefits

4 COATED TABLETS

Lil' DrugStore®



DO NOT USE IF INNER POUCH OR CARTON IS OPEN OR DAMAGED!

Drug Facts

Active ingredient (in each tablet) Purposes Aspirin 325 mg (NSAID)*......Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug

Uses temporarily relieves

- headache
- · muscle pain
- · toothache
- menstrual pain
- pain and fever of colds
 minor pain of arthritis

Reye's syndrome: Children and teenagers who have or are recovering form chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives facial swelling asthma (wheezing) shock Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
 take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

LIFT PANEL FOR CONTINUED DRUG FACTS



Drug Facts (continued)

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours
- · children under 12 years: consult a doctor

Other information

- save carton for full directions and warnings
- store at room temperature

Inactive ingredients carnauba wax*, corn starch, hypromellose, powdered cellulose, triacetin *may contain this ingredient

Questions or comments?

1-800-331-4536 or www.bayeraspirin.com

This product is repackaged by Lil' Drug Store Products, Inc. 1201 Continental Place NE, Cedar Rapids, IA 52402, 1-877-507-6516 (M-F 8AM-4:30PM CST), www.lildrugstore.com from a product manufactured by Bayer Corporation; Bayer® is a registered trademark of Bayer Corporation, used with permission. 97262C-US-02-12







Drug Facts (continued)

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- · you have asthma

Ask a doctor or pharmacist before use if you are taking a

· arthritis gout

Stop use and ask a doctor if

- · an allergic reaction occurs. Seek medical help right away.
- · you experience any of the following signs of stomach bleeding:
- · feel faint · vomit blood · have bloody or black stools
- · have stomach pain that does not get better pain gets worse or lasts more than 10 days
- redness or swelling is present
 fever lasts more than 3 days
- · new symptoms occur
- ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get

BAYER

aspirin tablet

Product Information

Product TypeHUMAN OTC DRUG LABELItem Code (Source)NDC:66715-9726

Route of Administration ORAL DEA Schedule

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength
ASPIRIN (ASPIRIN) ASPIRIN 325 mg

Inactive Ingredients

Ingredient Name Strength

carnauba wax
starch, corn
hypromelloses
powdered cellulose
triacetin

Product Characteristics

Color WHITE Score no score

Shape ROUND Size 10 mm

Flavor Imprint Code BAYER;BAYER

Contains

Packaging

# Item Code	Package Description	Marketing Start Date	Marketing End Date	
1 NDC:66715-9726-2	2 in 1 CARTON			
1	2 in 1 POUCH			

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part343	07/01/2009	

Labeler - Lil' Drug Store Products, Inc. (093103646)

Revised: 1/2013 Lil' Drug Store Products, Inc.