

READY CARE XL KIT - first aid kit
Dukal Corporation

Front Label

Physicians Care

FIRST AID

50 Person

ReadyCare XL Kit

Contains over 355 Pieces

Back Label

Physicians Care

CLEANSE

20 Antiseptic Wipes

20 Alcohol Pads

TREAT

6 Antibiotic Ointment Packets

6 Burn Ointment Packets

10 3" Cotton Tip Applicators

PROTECT

3 Knuckle Bandages

3 Fingertip Bandages

35 3/8" x 1 1/2" Bandages

16 1" x 3" Bandages

200 3/4" x 3" Bandages

1 1/2" Tape

1 Triangular Bandage

4 3" x 3" Sterile Gauze Pads

1 Abdominal Compress

Medicine

3 Acetaminophen Packets (packet of 2)

3 Aspirin Packets (packets of 2)

3 Antacid Packets (packets of 2)

Other

Gloves (2 Pair)

1 Cold Pack

1 First Aid Guide

10 Finger Splints

5 Disposable Thermometers

1 Tweezer

Acme United Corporation
60 Round Hill Road

Fairfield CT 06824

www.acmeunited.com

Designed in the USA | Made in China

CAUTION: This product may contain natural rubber latex which may cause allergic reactions. This kit contains products that have expiration dates. Please check before use.

BZK Towelette Labeling

Reorder 855

NDC 65517-0004-1

Dukal

BZK TOWELETTE

Contains Benzalkonium Chloride

For External Use Only

Helps Prevent Infection

1 / Pouch

DUKAL CORPORATION

(631) 656-3800

Ronkonkoma, NY 11779 www.dukal.com

Made in China

Drug Facts

Active Ingredients.....Benzalkonium Chloride 0.133% w/v

Purpose..... *First Aid Antiseptic*

USE: Antiseptic Cleansing of face, hands and body without soap and water. Airs dries in seconds

DO NOT USE: in the eyes or apply over large areas of the body.

STOP USE: If irritation, redness or other symptoms develop. Consult a doctor if the conditions persists or gets worse.

CAUTION: Keep out of reach of children. If swallowed get medical help or contact a Poison Control Center right away.

DIRECTIONS: Tear open packet, unfold and use as a washcloth.

INACTIVE INGREDIENTS: Distilled Water

Prep Pad Labeling

Reorder 852

Dukal Corporation

NDC 65517-0001-1

NPN 80003156

ALCOHOL PREP PAD

Saturated with 70% Isopropyl Alcohol

For External Use Only

1 / Pouch

Dukal Corporation

Ronkonkoma, NY 11779

631-656-3800

www.dukal.com

Made in China

Drug Facts**Active Ingredients**

Isopropyl Alcohol 70%

Purpose

Antiseptic Cleanser

Use

For Preparation of Skin prior to an injection

Warnings

- For External Use Only
- Flammable, Keep away from fire or flame

Do Not Use

- with electrocautery procedures
- In the Eyes. If contact occurs, flush eyes with water

Stop Use

If irritation and redness develop. If condition persists, consult your health care practitioner.

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Wipe injection site vigorously and discard.

Other Information

Store at Room Temperature 15 - 30 C (59 - 86 F)

Inactive Ingredient

purified water

Triple Antibiotic Labeling

WaterJel

Triple Antibiotic

First Aid Ointment

To Help Prevent Infection

Each Gram contains

Bacitracin Zinc 400 units

Neomycin Sulphate 5 mg

(equivalent to 3.5 mg Neomycin base)

Polymyxin B Sulfate 5000 units

Water-Jel Technologies

Carlstadt, NJ 07072

Drug Facts

Uses to help prevent infection in
minor cuts, scrapes, burns

Warnings

For external use only

Do not use

in the eyes or apply over large areas of the body

If you are allergic to any of the ingredients

longer than 1 week unless directed by a doctor

Stop use and ask a doctor if

the condition persists or gets worse

a rash or other allergic reaction develops

Keep out of reach of children

if ingested contact Poison Control Center right away

Directions

clean affected area apply a small amount of product

(an amount equal to the surface area of the tip of a finger)

on the area 1 to 3 times daily may be covered with a sterile bandage

First Aid Burn Cream Labeling

WaterJel

First Aid Burn Cream

Antiseptic Pain Relief with Aloe

Active Ingredients:

Benzalkonium Chloride 0.13%

Lidocaine HCL 0.5%
Water-Jel Technologies
Carlstadt, NJ 07072

Drug Facts

Purpose

First Aid Antiseptic, External analgesic

Uses

first aid to help prevent infection and for temporary relief of pain an itching associated with minor cuts, scrapes, burns

Warnings

For external use only

Do not use

in the eyes
in large quantities over raw or blistered areas or on deep puncture wounds, animal bites, or serious burns

Keep out of reach of children

if ingested contact Poison Control Center right away

Directions

clean affected area apply a small amount not more than 3 times daily may be covered with a sterile bandage

Other Information

Store at room temperature

APAP Labeling

Drug Facts

Active Ingredients (In each tablet).....Purposes
Acetaminophen 325 mg.....Pain Reliever/Fever Reducer

Uses:

For the temporary relief of minor aches and pains associated with:

- headache • muscular aches • minor arthritis pain
- common cold • toothache • menstrual cramps

For the reduction of fever.

Warnings:

Alcohol Warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers/fever reducers. Acetaminophen may cause liver damage.

Do not use:

- with any other product containing acetaminophen
- for more than 10 days for pain unless directed by a doctor
- for more than 3 days for fever unless directed by a doctor

Stop Using and Ask a Doctor If:

- symptoms do not improve
- new symptoms occur
- pain or fever persists or gets worse
- redness or swelling is present

Do not exceed recommended dosage. Keep this and all drugs out of the reach of children. In case of accidental overdose, contact a physician or Poison Control Center immediately. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

If pregnant or breast feeding, ask a health professional before use.

Directions:

Do not use more than directed

Adults and children: (12 years and older)

Take 2 tablets every 4 to 6 hours as needed. Do not take more than 12 tablets in 24 hours, or as directed by a doctor.

Children under 12 years:

Do not give to children under 12 years of age.

Other Information:

- Store at room temperature
- Tamper-Evident Sealed Packets
- Do Not Use any Opened or Torn Packets

Inactive Ingredients:

Corn starch*, croscarmellose sodium, crospovidone, hypromellose, microcrystalline cellulose, mineral oil, opadry clear, pharmaceutical glaze, polyethylene glycol, povidone*, pregelatinized starch*, sodium carboxymethylcellulose, sodium starch glycolate, stearic acid*, talc*, titanium dioxide.

*may contain

Questions or comments? 1-800-634-7680

Alcalak Labeling

Drug Facts

Active Ingredients (In each tablet).....Purposes

Calcium Carbonate 420 mgAntacid

Uses:

For the relief of the following symptoms associated with:

- acid indigestion • sour stomach • heartburn • upset stomach

Warnings:

Ask a doctor or health professional before use if you have:

- been taking a prescription drug. Antacids may interact with certain prescription drugs.
- kidney stones
- a calcium-restricted diet

Stop using this product and ask a doctor if:

- symptoms last more than 2 weeks

Do not exceed recommended dosage.

Keep this and all drugs out of the reach of children. If you are pregnant or breast feeding, ask a health professional before use.

Directions:

- Do not use more than directed

Adults and children: (12 years and older)

Take 2 tablets every 2 or 3 hours as symptoms occur or as directed by a physician. Do not take more than 19 tablets in a 24 hour period, or use the maximum dosage of this product for more than 2 weeks, except under the advice and supervision of a physician.

Children under 12 years:

Do not give to children under 12 years of age.

Other Information:

- Phenylketonurics: Contains Phenylalanine 1.5 mg per tablet
- Each tablet contains 168 mg of elemental calcium
- Store at room temperature in a dry place
- Tamper-Evident Sealed Packets
- Do not Use any Opened or Torn Packets

Inactive Ingredients:

aspartame*, carageenan*, croscarmellose sodium*, glycine*, magnesium stearate, mint flavor*, silica*, sorbitol, spearmint flavor*, stearic acid*, stevix*, xylitol*

*may contain

Questions or comments? 1-800-634-7680

Aspirin Labeling**Drug Facts**

Active Ingredients (In each tablet).....Purposes

Aspirin 325mg (NSAID*)Pain Reliever / Fever Reducer

*nonsteroidal anti-inflammatory drug

Uses:

Temporarily relieves minor aches and pains due:

- headache • muscular aches • minor arthritis pain • backache
- common cold • toothache • menstrual cramps

Temporarily reduces fever.

Warnings:

Reye's Syndrome: Children and teenagers should not use this medicine for chicken pox, or flu symptoms before a doctor is consulted about Reye's Syndrome, a rare but serious illness reported to be associated with aspirin.

Allergy Alert: Aspirin may cause a severe allergic reaction which may include:

- hives • skin reddening • facial swelling • rash • asthma (wheezing) • blisters • shock

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach Bleeding Warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding.

The chance is higher if you:

- are age 60 or older • have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcohol drinks every day while using this product
- take more or for a longer time than directed

Do not use:

- if you have ever had an allergic reaction to any other pain reliever/ fever reducer.
- right before or after heart surgery

Ask a Doctor before use if you have:

- problems or serious side effects from taking pain relievers or fever reducers
- stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain • ulcers • bleeding problems • high blood pressure
- heart or kidney disease • taken a diuretic • reached age 60 or older

Ask a Doctor or Pharmacist before use if you are:

- taking any other drug containing and NSAID (prescription or nonprescription)
- taking a blood thinning (anticoagulant) or steroid drug
- under a doctor's care for any serious condition • taking any other drug

When using this product:

- take with food or milk if stomach upset occurs
- long term continuous use may increase the risk of heart attack or stroke

Stop Using and Ask a Doctor If:

- you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- you develop heartburn
- stomach pain or upset gets worse or lasts
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions:

- Do not take more than directed.
- The smallest effective dose should be used.
- Do not take longer than 10 days, unless directed by a doctor (see Warnings)
- Drink a full glass of water with each dose.

Adults and children: (12 years and older)

Take 1 or 2 tablets with water every 4 hours as needed. Do not take more than 12 tablets in 24 hours, or as directed by a doctor.

Children under 12 years:

Do not give to children under 12 years of age.

Other Information:

- Read all product information before using
- Store at room temperature
- Avoid excessive heat and humidity
- Tamper evident sealed packets
- Do not use any opened or torn packets

Inactive Ingredients:

Corn starch, croscarmellose sodium*, hypromellose*, microcrystalline cellulose*, mineral oil*, polyethylene glycol, propylene glycol, silicon dioxide, starch, stearic acid, talc, titanium dioxide*.

*may contain

Questions or comments? 1-800-634-7680

Ready Care XL Kit Label




Ready Care XL Kit Back Label

PhysiciansCARE

Distributed by: Acme United Corporation | Fairfield, CT 06624 | 203-254-6090. PhysiciansCare® is a registered trademark of Acme United Corporation | www.acmeunited.com | Designed in the USA | Manufactured in China

CAUTION: This product may contain natural rubber latex which may cause allergic reactions. This kit contains products that have expiration dates. Please check before use.

CLEANSE		PROTECT		MEDICINE		
20 Antiseptic Wipes  Item refill #51028 Made in China	20 Alcohol pads  Item refill #51019 Made in China	35 3/8" X 1 1/4" Bandages  Made in China		3 Acetaminophen Packets (packet of 2)  Item refill #90016 Made in the USA	3 Aspirin Packets (packet of 2)  Item refill #90014 Made in the USA	3 Antacid Packets (packet of 2)  Item refill #90099 Made in the USA
TREAT		16 1" X 3" Bandages  Item refill #51005 Made in China		OTHER		
6 Antibiotic Ointment Packets  Item refill #12944 Made in the USA	6 Burn Ointment Packets  Item refill #51014 Made in the USA	200 3/4" X 3" Bandages  Item refill #90600 Made in China		Medical Exam Latex Gloves (2 pair)  Item refill# 51016 Made in China	10 Finger Splints  Made in China	
10 3" Cotton Tip Applicators  Item refill #12941 Made in China		1 1/2" x 10yd tape  Item refill #12302 Made in China	4 3" x 3" Gauze  Made in China	1 Cold Pack  Item refill# 51013 Made in China	5 Disposable Thermometers  Made in China	
PROTECT		1 Triangular Bandage  Item refill #51012 Made in China	1 Abdominal Compress  Item refill# 90020 Made in China	1 First Aid Guide  Printed in China	1 Tweezers  Item refill #90024 Made in China	
3 Knuckle Bandages  Item refill #51010 Made in China	3 Fingertip Bandages  Item refill #51009 Made in China					
						<small>000151004 Item 1</small>

Prep Pad Labeling



**REORDER
852**

**NDC 65517-0001-1
NPN 8003156**

ALCOHOL PREP PAD

●●●●●●●●●●●●●●●●

1

**Saturated with 70%
Isopropyl Alcohol**

For External Use Only



DUKAL CORPORATION • (631) 656-3800

Ronkonkoma, NY 11779

Made in China

D11040501 Rev1

Drug Facts

Active Ingredients	Purpose
Isopropyl Alcohol 70%.....	Antiseptic Cleanser
Use For preparation of the skin prior to an injection	
Warnings	
<ul style="list-style-type: none"> • For external use only • Flammable, keep away from fire or flame 	
Do Not Use	
<ul style="list-style-type: none"> • with electrocautery procedures • In the eyes. If contact occurs, flush eyes with water 	
Stop Use If irritation and redness develop. If condition persists consult your health care practitioner.	
Keep out of reach of children. If swallowed get medical help or contact a Poison Control Center right away.	
Directions	
<ul style="list-style-type: none"> • Wipe injection site vigorously and discard 	
Other Information	
Store at room temperature 15° - 30° C (59° - 86° F)	
Inactive Ingredient purified water	

Antiseptic Towelette Label



REORDER
855

NDC 65517-0004-1

BZK TOWELETTE



Contains Benzalkonium Chloride
For External Use Only
Helps Prevent Infection

1
/Pouch

2 DUKAL CORPORATION • (631) 656-3800
Ronkonkoma, NY 11779 • www.dukal.com

Made in China

DRUG FACTS:

Active Ingredient	Purpose
Benzalkonium Chloride, 0.133% w/v...	First Aid Antiseptic

USE: Antiseptic Cleansing of face, hands and body without soap and water. Air dries in seconds.

DO NOT USE: in the eyes or apply over large areas of the body.

STOP USE: if irritation, redness or other symptoms develop. Consult a doctor if the condition persists or gets worse.

CAUTION: Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

DIRECTIONS: Tear open packet, unfold and use as a washcloth.

INACTIVE INGREDIENTS: Distilled Water

DUKAL CORPORATION

D07110604 Rev1

Tear Here

Antacid Label

Medique
2 Tablets Made in USA

Alcalak

Active Ingredient (In Each Tablet)	Purpose
Calcium Carbonate 420 mg.....	Antacid

Uses:
For the relief of the following symptoms associated with:

- acid indigestion • sour stomach
- heartburn • upset stomach

Warnings:
Ask a doctor or health professional before use if you have:

- been taking a prescription drug. Antacids may interact with certain prescription drugs.
- kidney stones • a calcium-restricted diet

Store at room temperature in a dry place
Tamper-Evident Sealed Packets
Do Not Use any Opened or Torn Packets
Mfd. for MEDIQUE PRODUCTS • Fort Myers, FL 33967

Warnings: (continued)

Stop using this product and ask a doctor if:

- symptoms last more than 2 weeks.

Do not exceed recommended dosage. Keep this and all drugs out of the reach of children. As with any drug, if you are pregnant or nursing a baby, seek the advice of a health professional before using this product.

Directions:
Do not use more than directed.

Adults and children: (12 years and older) Take 2 tablets every 2 or 3 hours as symptoms occur or as directed by a physician. Do not take more than 19 tablets in a 24 hour period, or use the maximum dosage of this product for more than 2 weeks, except under the advice and supervision of a physician.

Children under 12 years: Do not give to children under 12 years of age.

Other Information:

- Phenylketonurics: Contains Phenylalanine 1.5 mg per tablet.
- Each tablet contains: 168 mg of elemental calcium.

SEE CARTON FOR COMPLETE PRODUCT INFORMATION

Lot 1747 Exp. 11-2013

Acetaminophen Label

Medique[®] Made in USA

2 Tablets **APAP**

Active ingredient (in each tablet) Purpose
 Acetaminophen 325 mg..... Pain reliever/
 fever reducer

Uses
 For the temporary relief of minor aches and pains associated with

- headache ■ muscular aches ■ minor arthritis pain
- toothache ■ common cold ■ menstrual cramps

For the reduction of fever.

Warnings
 Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 tablets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Store at 59°-86°F (15-30°C) • Tamper-evident sealed packets • Do not use any opened or torn packets
 Mfd. for **MEDIQUE PRODUCTS • Fort Myers, FL 33967**

Warnings (continued)
Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for more than 10 days for pain unless directed by a doctor
- for more than 3 days for fever unless directed by a doctor

Ask a doctor before use if you have

- liver disease

Ask a doctor or pharmacist before use if

- you are taking the blood thinning drug warfarin

Stop using and ask a doctor if

- symptoms do not improve ■ new symptoms occur
- pain or fever persists or gets worse
- redness or swelling is present

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

If pregnant or breast-feeding, ask a health professional before use.

Directions ■ do not use more than directed

Adults and children: (12 years and older)
 Take 2 tablets every 4 to 6 hours as needed. Do not take more than 12 tablets in 24 hours.

Children under 12 years:
 Do not give to children under 12 years of age.

SEE CARTON FOR COMPLETE PRODUCT INFORMATION

Lot 1831 Exp. 01-2013

Aspirin Label

2 Tablets **Medique** Made in USA
Aspirin

Active ingredient (in each tablet) Aspirin (NSAID) 325 mg
Purpose Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses Temporarily relieves minor aches and pains due to
 ■ headache ■ muscular aches ■ minor arthritis pain ■ backache
 ■ common cold ■ toothache ■ menstrual cramps
 Temporarily reduces fever.

Warnings
 Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
 Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ skin reddening ■ facial swelling ■ rash ■ asthma (wheezing) ■ blisters ■ shock
 If an allergic reaction occurs, stop use and seek medical help right away.
 Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
 ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinner (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Store at 59°-86°F (15°-30°C) • Tamper-Evident Sealed Packets
 • Do Not Use any Opened or Torn Packets
 Mfd. for MEDIQUE PRODUCTS • Fort Myers, FL 33967

Warnings (continued)
 Do not use
 ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery ■ if you are taking a prescription drug for gout, diabetes or arthritis

Ask a doctor before use if
 ■ stomach bleeding warning applies to you
 ■ you have a history of stomach problems such as heartburn
 ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are
 ■ under a doctor's care for any serious condition ■ taking any other drug

When using this product
 ■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if
 ■ you experience any of the following signs of stomach bleeding:
 ■ feel faint ■ vomit blood ■ have bloody or black stools
 ■ have stomach pain that does not get better
 ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ you have difficulty swallowing ■ if ringing in the ears or loss of hearing occurs ■ redness or swelling is present in the painful area
 ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions: Do not use more than directed
 Adults and children: (12 years and older) Take 1 or 2 tablets with water every 4 hours as needed. Do not take more than 12 tablets in 24 hours, or as directed by a doctor.
 Children: (under 12 years) Do not give to children under 12 years of age.

SEE CARTON FOR COMPLETE PRODUCT INFORMATION

Lot 1856 Exp. 11-2012

Triple Antibiotic Label

WATERMEL

Triple Antibiotic

First Aid Ointment
 To Help Prevent Infection

EACH GRAM CONTAINS:
 Bacitracin Zinc 400 units
 Neomycin Sulfate 5 mg
 (equivalent to 3.5 mg
 Neomycin base)
 Polymyxin B Sulfate 5000 units

NET WT 0.9g (1/2" OZ)

Water Jet Technology
 Carlisle, NJ 07017

Drug Facts
Uses to help prevent infection in
 ■ minor cuts ■ scrapes ■ burns

Warnings
For external use only
Do not use ■ in the eyes or apply over large areas of the body ■ if you are allergic to any of the ingredients
 ■ longer than 1 week unless directed by a doctor

Stop use and ask a doctor if
 ■ the condition persists or gets worse ■ a rash or other allergic reaction develops

Keep out of reach of children
 If ingested contact Poison Control Center right away.

Directions ■ clean affected area ■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily ■ may be covered with a sterile bandage

LOT 8718 EXP 01/2011

Burn Cream Label



READY CARE XL KIT

first aid kit kit

Product Information

Product Type	MEDICAL DEVICE	Item Code (Source)	NDC:65517-0021
---------------------	----------------	---------------------------	----------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65517-0021-1	1 in 1 PACKAGE, COMBINATION		

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	20 POUCH	8.0 mL
Part 2	6 PACKET	5.4 g
Part 3	6 PACKET	5.4 g
Part 4	20 POUCH	28.0 mL
Part 5	3 PACKET	6
Part 6	3 PACKET	6
Part 7	3 PACKET	6

Part 1 of 7

DUKAL ALCOHOL PREP PAD

isopropyl alcohol swab

Product Information

Item Code (Source) NDC:65517-0001

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ISOPROPYL ALCOHOL (UNII: ND2M416302) (ISOPROPYL ALCOHOL - UNII:ND2M416302)	ISOPROPYL ALCOHOL	0.70 mL in 1.0 mL

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65517-0001-1	0.4 mL in 1 POUCH		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333A	01/01/2000	

Part 2 of 7

WATER-JEL 3-IN1 ANTIBIOTIC

bacitracin zinc, polymyxin b sulfate and neomycin sulfate ointment

Product Information

Item Code (Source) NDC:59898-740

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BACITRACIN ZINC (UNII: 89Y4M234ES) (BACITRACIN - UNII:58H6RWO52I)	BACITRACIN ZINC	400 [iU] in 1 g
NEOMYCIN SULFATE (UNII: 057Y626693) (NEOMYCIN - UNII:H6QD7X297)	NEOMYCIN SULFATE	3.5 mg in 1 g
POLYMYXIN B SULFATE (UNII: 19371312D4) (POLYMYXIN B - UNII:J2VZ07J96K)	POLYMYXIN B SULFATE	5000 [iU] in 1 g

Inactive Ingredients

Ingredient Name	Strength
PETROLATUM (UNII: 4T6H12BN9U)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59898-740-01	0.9 g in 1 PACKET		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part333	01/01/2010	

Part 3 of 7

FIRST AID BURN

lidocaine hydrochloride and benzalkonium chloride cream

Product Information

Item Code (Source)	NDC:59898-902
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y)	BENZALKONIUM CHLORIDE	0.13 g in 100 g
LIDOCAINE HYDROCHLORIDE (UNII: V13007Z41A) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE HYDROCHLORIDE	0.5 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
EDETATE DISODIUM (UNII: 7FLD91C86K)	
LIGHT MINERAL OIL (UNII: N6K5787QVP)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
CETYL ALCOHOL (UNII: 936JST6JCN)	
TROLAMINE (UNII: 9O3K93S3TK)	
GLYCERIN (UNII: PDC6A3C0OX)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
DIAZOLIDINYL UREA (UNII: H5RIZ3MPW4)	
ALOE VERA LEAF (UNII: ZY81Z83H0X)	
WATER (UNII: 059QF0KO0R)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
GLYCERYL MONOSTEARATE (UNII: 230OU9XXE4)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59898-902-01	0.9 g in 1 PACKET		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part333	01/01/2010	

Part 4 of 7

BZK TOWELETTE

benzalkonium chloride swab

Product Information

Item Code (Source)	NDC:65517-0004
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y)	BENZALKONIUM CHLORIDE	0.00186 mL in 1.4 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	1.39814 mL in 1.4 mL

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65517-0004-1	1.4 mL in 1 POUCH		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333A	01/01/2006	

Part 5 of 7

MEDIQUE APAP

acetaminophen tablet, film coated

Product Information

Item Code (Source)	NDC:47682-145
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSE (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	white (white)	Score	no score
Shape	ROUND (ROUND)	Size	10mm
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-145-99	2 in 1 PACKET		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part343	12/30/2008	

Part 6 of 7**MEDIQUE ASPIRIN**

aspirin tablet, film coated

Product Information

Item Code (Source)	NDC:47682-116
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSE (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	white (white)	Score	no score
Shape	ROUND (ROUND)	Size	10 mm
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-116-99	2 in 1 PACKET		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part343	12/30/2008	

Part 7 of 7**MEDIQUE ALCALAK**

calcium carbonate tablet, chewable

Product Information

Item Code (Source)	NDC:47682-088
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Calcium Carbonate (UNII: H0G9379FGK) (Calcium - UNII:SY7Q814VUP)	Calcium Carbonate	420 mg

Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
SUCROSE (UNII: C151H8M554)	

Product Characteristics

Color	white (white)	Score	no score
Shape	ROUND (ROUND)	Size	12mm
Flavor	MINT (MINT)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-088-99	2 in 1 PACKET		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part331	12/30/2008	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part333	05/01/2010	

Labeler - Dukal Corporation (791014871)