

**TUSSIN CF NON DROWSY- dextromethorphan hbr, guaifenesin,
phenylephrine liquid
QUALITY CHOICE (Chain Drug Marketing Association)**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredients (in each 10 mL)

Dextromethorphan HBr 20 mg

Guaifenesin 200 mg

Phenylephrine HCl 10 mg

Purposes

Cough suppressant

Expectorant

Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
 - nasal congestion
 - cough due to minor throat and bronchial irritation

Warnings

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- diabetes
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic

bronchitis or emphysema

Ask a doctor or pharmacist before use if you are

taking any other oral nasal decongestant or stimulant.

When using this product,

do not use more than directed.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age
- adult and children 12 years and over: 10 mL every 4 hours
- children under 12 years: do not use

Other information

- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

anhydrous citric acid, FD&C red #40, flavor, glycerin, lactic acid, menthol, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose

Questions or comments?

Call **1-248-449-9300 Monday-Friday 9AM-5PM EST**

Principal Display Panel

*Compare to the active ingredients in Robitussin® Peak Cold Multi-Symptom Cold CF

Non-Drowsy

Tussin CF

Multi-Symptom Cold

Dextromethorphan HBr Cough Suppressant

Guaifenesin Expectorant

Phenylephrine HCl Nasal Decongestant

Relieves:

Cough

Mucus

Nasal Congestion

For Ages 12 & Over

Dosing Cup Included

Alcohol Free

FL OZ (mL)

*This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributors of Robitussin® Peak Cold Multi-Symptom Cold CF.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER THE CAP IS BROKEN OR MISSING.

Distributed by C.D.M.A., Inc. ©

43157 W. Nine Mile

Novi, MI 48375-0995

www.qualitychoice.com

Questions: 248-449-9300

Package Label

Drug Facts (continued)

Inactive ingredients citric acid, FD&C red #40, flavor, glycerin, lactic acid, menthol, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose

Questions or comments?
Call 1-800-935-2362 Monday-Friday 9AM-5PM EST

*This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Robitussin® Peak Cold Multi-Symptom Cold CF.

TAMPER EVIDENT: DO NOT USE IF CAPTOPIN IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org



Dosing Cup Included



Distributed by C.D.M.A., Inc.®
43157 W 9 Mile Rd
Novi, MI 48375
www.qualitychoice.com
Questions: 800-935-2362



NDC 63868-244-08

*Compare to the active ingredients in Robitussin® Peak Cold Multi-Symptom Cold CF

Non-Drowsy Tussin CF

Multi-Symptom Cold

Dextromethorphan HBr
Cough Suppressant
Guaifenesin
Expectorant
Phenylephrine HCl
Nasal Decongestant

Relieves:
Cough
Mucus
Nasal Congestion

For Ages 12 & Over
Alcohol-Free

8 FL OZ (237 mL)



NDC 63868-244-08

*Compare to the active ingredients in Robitussin® Peak Cold Multi-Symptom Cold CF

Non-Drowsy Tussin CF

Multi-Symptom Cold

Dextromethorphan HBr
Cough Suppressant
Guaifenesin
Expectorant
Phenylephrine HCl
Nasal Decongestant

Relieves:
Cough
Mucus
Nasal Congestion

For Ages 12 & Over
Alcohol-Free

8 FL OZ (237 mL)



PLD-E322C F0007425

Lot No.:
Exp. Date:

Drug Facts

Active ingredients (in each 10 mL)
Dextromethorphan HBr 20 mg.....Cough suppressant
Guaifenesin 200 mg.....Expectorant
Phenylephrine HCl 10 mg.....Nasal decongestant

Uses
■ helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
■ temporarily relieves these symptoms occurring with a cold
■ nasal congestion
■ cough due to minor throat and bronchial irritation

Warnings
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have
■ heart disease ■ high blood pressure
■ diabetes ■ thyroid disease
■ trouble urinating due to an enlarged prostate gland
■ cough that occurs with too much phlegm (mucus)
■ cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are taking any other oral nasal decongestant or stimulant.

When using this product, do not use more than directed.

Stop use and ask a doctor if
■ nervousness, dizziness, or sleeplessness occur
■ symptoms do not get better within 7 days or are accompanied by fever
■ cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions
■ do not take more than 6 doses in any 24-hour period
■ measure only with dosing cup provided. Do not use any other dosing device.
■ keep dosing cup with product
■ mL = milliliter
■ this adult product is not intended for use in children under 12 years of age
■ adults and children 12 years and over: 10 mL every 4 hours
■ children under 12 years: do not use

Other information
■ store between 20-25°C (68-77°F). Do not refrigerate.

QUALITY CHOICE Non-Drowsy Tussin CF Multi-Symptom Cold

TUSSIN CF NON DROWSY

dextromethorphan hbr, guaifenesin, phenylephrine liquid

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:63868-244 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|----------------------------------|-------------------|
| DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 20 mg in 10 mL |

| | | |
|---|--------------------------------|--------------------|
| GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ) | GUAIFENESIN | 200 mg in 10 mL |
| PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV) | PHENYLEPHRINE HYDROCHLORIDE | 10 mg in 10 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| WATER (UNII: 059QF0KO0R) | |
| SODIUM BENZOATE (UNII: OJ245FE5EU) | |
| FD&C RED NO. 40 (UNII: WZB9127XOA) | |
| LACTIC ACID (UNII: 33X04XA5AT) | |
| MENTHOL (UNII: L7T10EIP3A) | |
| SORBITOL (UNII: 506T60A25R) | |
| SUCRALOSE (UNII: 96K6UQ3ZD4) | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:63868-244-04 | 1 in 1 BOX | 06/30/2014 | 12/31/2024 |
| 1 | | 118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | | |
| 2 | NDC:63868-244-08 | 1 in 1 BOX | 06/30/2014 | 12/31/2024 |
| 2 | | 237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC monograph final | part341 | 06/30/2014 | 12/31/2024 |

Labeler - QUALITY CHOICE (Chain Drug Marketing Association) (011920774)

Revised: 5/2023

QUALITY CHOICE (Chain Drug Marketing Association)