

LOPERAMIDE HYDROCHLORIDE- loperamide hydrochloride liquid H E B

HEB Loperamide Hydrochloride Oral Suspension Drug Facts

Active ingredient (in each 7.5 mL)

Loperamide HCl 1 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Do not use

if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease

Ask a doctor or pharmacist before use if you are

taking antibiotics

When using this product

- tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging.

These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- shake well before using
- only use attached measuring cup to dose product

adults and children 12 years and over	30 mL (6 tsp) after the first loose stool; 15 mL (3 tsp) after each subsequent loose stool; but no more than 60 mL (12 tsp) in 24 hours
children 9-11 years (60-95 lbs)	15 mL (3 tsp) after the first loose stool; 7.5 mL (1 1/2 tsp) after each subsequent loose stool; but no more than 45 mL (9 tsp) in 24 hours
children 6-8 years (48-59 lbs)	15 mL (3 tsp) after the first loose stool; 7.5 mL (1 1/2 tsp) after each subsequent loose stool; but no more than 30 mL (6 tsp) in 24 hours
children under 6 years (up to 47 lbs)	ask a doctor

Other information

- each 30 mL (6 tsp) contains: sodium 13 mg
- store between 20-25°C (68-77°F)
- do not use if printed plastic neckband is broken or missing
- see side panel for lot number and expiration date

Inactive ingredients

anhydrous citric acid, carboxymethylcellulose sodium, D&C yellow no. 10, FD&C blue no. 1, glycerin, microcrystalline cellulose, natural and artificial mint flavor, propylene glycol, purified water, simethicone, sodium benzoate, sucralose, titanium dioxide, xanthan gum

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Compare to Imodium® A-D active ingredient

LOPERAMIDE HYDROCHLORIDE ORAL SUSPENSION

Anti-Diarrheal

Controls the Symptoms of Diarrhea

Mint Flavor

GLUTEN FREE

1 mg Loperamide Hydrochloride per 7.5 mL

Drug Facts

Active ingredient Purpose (in each 7.5 mL)

Loperamide HCl
1 mg.....Anti-diarrheal

Use controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Do not use if you have bloody or black stool

PEEL BACK HERE

Compare to Imodium® A-D active ingredient*

NDC 37808-645-26



LOPERAMIDE HYDROCHLORIDE ORAL SUSPENSION

Anti-Diarrheal

Controls the Symptoms of Diarrhea

■ Mint Flavor

4 FL OZ
(120 mL)

1 mg Loperamide
Hydrochloride per 7.5 mL

GLUTEN FREE

MADE WITH
PRIDE & CARE FOR H-E-B
SAN ANTONIO, TX 78204

GUARANTEE

We believe the high quality of this H-E-B product makes it an outstanding value. We hope you'll agree. If not, we'll cheerfully refund your money. Thanks for shopping with us.

*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Imodium® A-D.

1108



: 64526 1J FL

Drug Facts (continued)

Ask a doctor before use if you have ■ fever
■ mucus in the stool ■ a history of liver disease

Ask a doctor or pharmacist before use if you are taking antibiotics

When using this product ■ tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

■ symptoms get worse
■ diarrhea lasts for more than 2 days
■ you get abdominal swelling or bulging.
These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison

Drug Facts (continued)

Control Center right away. (1-800-222-1222)

Directions ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

■ find right dose on chart. If possible, use weight to dose; otherwise use age. ■ shake well before using ■ only use attached measuring cup to dose product

adults and children 12 years and over	30 mL (6 tsp) after the first loose stool; 15 mL (3 tsp) after each subsequent loose stool; but no more than 60 mL (12 tsp) in 24 hours
--	---

children 9-11 years (60-95 lbs)	15 mL (3 tsp) after the first loose stool; 7.5 mL (1 1/2 tsp) after each subsequent loose stool; but no more than 45 mL (9 tsp) in 24 hours
------------------------------------	--

Drug Facts (continued) children 6-8 years (48-59 lbs) 15 mL (3 tsp) after the first loose stool; 7.5 mL (1 1/2 tsp) after each subsequent loose stool; but no more than 30 mL (6 tsp) in 24 hours children under 6 years (up to 47 lbs) ask a doctor		Drug Facts (continued) do not use if printed plastic neckband is broken or missing see side panel for lot number and expiration date Inactive ingredients anhydrous citric acid, carboxymethylcellulose sodium, D&C yellow no. 10, FD&C blue no. 1, glycerin, microcrystalline cellulose, natural and artificial mint flavor, propylene glycol, purified water, simethicone, sodium benzoate, sucralose, titanium dioxide, xanthan gum
Other information each 30 mL (6 tsp) contains: sodium 13 mg store between 20-25°C (68-77°F)		Questions or comments? 1-800-719-9260

LOPERAMIDE HYDROCHLORIDE

loperamide hcl liquid

Product Information

Product Type	HUMAN OTC DRUG LABEL	Item Code (Source)	NDC:37808-645
Route of Administration	ORAL	DEA Schedule	

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (LOPERAMIDE)	LOPERAMIDE HYDROCHLORIDE	1 mg in 7.5 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID	
CARBOXYMETHYLCELLULOSE SODIUM	
D&C YELLOW NO. 10	
FD&C BLUE NO. 1	
GLYCERIN	
CELLULOSE, MICROCRYSTALLINE	
PROPYLENE GLYCOL	
WATER	
SODIUM BENZOATE	
SUCRALOSE	
TITANIUM DIOXIDE	
XANTHAN GUM	

Product Characteristics

Color	GREEN (opaque, viscous)	Score	
-------	-------------------------	-------	--

Shape		Size		
Flavor	MINT	Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37808-645-26	120 mL in 1 BOTTLE		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANDA	ANDA091292		03/05/2012	

Labeler - HEB (007924756)