

## **UP AND UP ACETAMINOPHEN- acetaminophen tablet**

### **Target Corporation**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

### **Target Corporation Acetaminophen Drug Facts**

#### **Active ingredient (in each tablet)**

Acetaminophen 325 mg

#### **Purpose**

Pain reliever/fever reducer

#### **Uses**

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- backache
- minor pain of arthritis
- the common cold
- toothache
- premenstrual and menstrual cramps
- temporarily reduces fever

#### **Warnings**

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

#### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if the user has ever had an allergic reaction to this product or any of its ingredients

#### **Ask a doctor before use if the user**

- has liver disease
- is a child with pain of arthritis

#### **Ask a doctor or pharmacist before use if the user is**

taking the blood thinning drug warfarin

#### **Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days (for adults) or 5 days (for children)
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

#### **If pregnant or breast-feeding,**

ask a health professional before use.

#### **Keep out of reach of children.**

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### **Directions**

- do not take more than directed (see overdose warning)

adults and children 12 years and over	<ul style="list-style-type: none"> <li>• take 2 tablets every 4 to 6 hours while symptoms last</li> <li>• do not take more than 10 tablets in 24 hours</li> <li>• do not use for more than 10 days unless directed by a doctor</li> </ul>
children 6-11 years	<ul style="list-style-type: none"> <li>• take 1 tablet every 4 to 6 hours while symptoms last</li> <li>• do not take more than 5 tablets in 24 hours</li> <li>• do not use for more than 5 days unless directed by a doctor</li> </ul>
children under 6 years	ask a doctor

#### **Inactive ingredients**

croscarmellose sodium\*, povidone, pregelatinized starch, stearic acid \*may contain this ingredient

### **Questions?**

Call 1-888-547-7400

### **Principal Display Panel**

Compare to active ingredient in Regular Strength Tylenol® Tablets

regular strength

acetaminophen tablets, 325 mg

pain reliever/fever reducer

100 TABLETS

ACTUAL SIZE

100 TABLETS

regular strength  
**acetaminophen tablets,**  
**325 mg**

pain reliever/fever reducer



Compare to active ingredient  
in Regular Strength Tylenol® Tablets\*\*

NDC 11673-444-78

regular strength  
**acetaminophen tablets,**  
**325 mg**

pain reliever/fever reducer



100 TABLETS



ACTUAL SIZE

**100**

TABLETS

EXP. DATE

LOT NO.



DO NOT USE IF PRINTED SEAL UNDER CAP IS  
BROKEN OR MISSING

**Drug Facts**

**Active ingredient**  
(in each tablet)

**Purpose**

Acetaminophen 325 mg...Pain reliever/fever reducer

**Uses** ■ temporarily relieves minor aches and pains due to:

- the common cold
- headache
- minor pain of arthritis
- backache
- muscular aches
- toothache
- premenstrual and menstrual cramps

■ temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if:

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if the user has ever had an allergic reaction to this product or any of its ingredients

**Drug Facts (continued)**

Ask a doctor before use if the user:

- has liver disease
- is a child with pain of ear, throat, or tonsils

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin.

Stop use and ask a doctor if:

- new symptoms occur
- pain gets worse or lasts more than 10 days (for adults) or 5 days (for children)
- fever gets worse or lasts more than 3 days
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. **Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you don't notice any signs or symptoms.

**Directions**

■ do not take more than directed (see overdose warning)

adults and children 12 years and over

- take 2 tablets every 4 to 6 hours while symptoms last
- do not take more than 10 tablets in 24 hours
- do not use for more than 10 days unless directed by a doctor


children 6-11 years

- take 1 tablet every 4 to 6 hours while symptoms last
- do not take more than 5 tablets in 24 hours
- do not use for more than 5 days unless directed by a doctor

children under 6 years

ask a doctor

Dr. Inc. 100 Tablets  
NDC 11673-444-78  
Upsher-Khan  
Reg. U.S. Pat. & Tm. Off.  
© 2014



## UP AND UP ACETAMINOPHEN

acetaminophen tablet

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-444
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

### Inactive Ingredients

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
POVIDONE (UNII: FZ989GH94E)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

### Product Characteristics

Color	WHITE	Score	no score
Shape	ROUND (beveled edge)	Size	10mm
Flavor		Imprint Code	325MG;L403
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-444-78	1 in 1 CARTON	08/20/2014	
1		100 in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	08/20/2014	

