

**TUSSIN CF - dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride liquid**

**Chain Drug Consortium, LLC**

-----  
**Active ingredients**

**DRUG FACTS**

**Active ingredients**

**(in each 5 mL tsp)**

Dextromethorphan HBr, USP 10 mg

Guaifenesin, USP 100 mg

Phenylephrine HCL, USP 5 mg

**Purposes**

**Purposes**

Cough Suppressant

Expectorant

Nasal Decongestant

**Keep out of reach of children**

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Uses**

**Uses**

- help loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold
- nasal congestion
- cough due to minor throat and bronchial irritation

**Warnings**

**Warnings**

**Do not use**

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug

contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use Ask a doctor or pharmacist before use  
Ask a doctor before use if you have**

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

**Ask a doctor or pharmacist before use if you are**  
taking any other oral nasal decongestant or stimulant.

**When using this product do not use more than directed.**

**Stop use and ask a doctor if**

**Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- symptoms do not get better within 7 days, or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache.

These could be signs of a serious condition.

**If pregnant or breast-feeding**

**If pregnant or breast feeding,**

ask a health professional before use.

**Directions**

**Directions**

- Do not take more than 6 doses in any 24 hour period

- This adult strength product is not intended for use in children under 12 years of age

Age	Dose
adults and children 12 years and over	2 teaspoons every 4 hours
children under 12	do not use

### **Other information**

#### **Other information**

- **Keep carton for full Direction for use.**
- store at 20-25 ° C (68-77 ° F)
- do not refrigerate
- dosage cup provided
- sodium 3 mg per teaspoonful

### **Inactive ingredients**

#### **Inactive ingredients**

anhydrous citric acid, FD and C red no. 40, glycerin, menthol, natural and artificial flavor, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

### **Questions?**

#### **Questions?**

Call weekdays from 9:30 AM to 4:30 PM EST at

**1-877-798-5944**

### **Product Label**

NDC 68016-126-04

**\*COMPARE TO THE ACTIVE INGREDIENTS IN ROBITUSSIN® PEAK COLD MULTI-SYMP TOM COLD**

**Premier Value ®  
Tussin CF**

**Dextromethorphan HBr/ Guaifenesin / Phenylephrine HCL**

**COUGH SUPPRESSANT  
EXPECTORANT/  
NASAL DECONGESTANT  
NON-DROWSY**

Multi-Symptom Formula

For ages 12 and over

8 FL OZ (237 mL)

**INDEPENDENTLY TESTED SATISFACTION GUARANTEED PV**

If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.

\*This product is not manufactured or distributed by Pfizer, owner of the trademark Robitussin®

DISTRIBUTED BY: CHAIN DRUG CONSORTIUM 3301 NW BOCA RATON BLVD SUITE 101,  
BOCA RATON, FL 33431  
BX-018

**DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING**

<b>Drug Facts</b>	
<b>Active ingredients (in each 5 mL tsp)</b>	<b>Purposes</b>
Dextromethorphan HBr, USP 10mg...Cough Suppressant Guaifenesin, USP 100 mg..... Expectorant Phenylephrine HCl, USP 5 mg.... Nasal Decongestant	
<b>Uses</b>	
<ul style="list-style-type: none"> <li>help loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes</li> <li>temporarily relieves these symptoms occurring with a cold:               <ul style="list-style-type: none"> <li>nasal congestion</li> <li>cough due to minor throat and bronchial irritation</li> </ul> </li> </ul>	
<b>Warnings</b>	
<p><b>Do not use</b> if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p><b>Ask a doctor before use if you have</b></p> <ul style="list-style-type: none"> <li>heart disease</li> <li>high blood pressure</li> <li>thyroid disease</li> <li>diabetes</li> <li>trouble urinating due to an enlarged prostate gland</li> <li>cough that occurs with too much phlegm (mucus)</li> <li>cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema</li> </ul> <p><b>Ask a doctor or pharmacist before use if you are</b> taking any other oral nasal decongestant or stimulant.</p> <p><b>When using this product do not use more than directed.</b></p> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>you get nervous, dizzy or sleepless</li> <li>symptoms do not get better within 7 days or are accompanied by fever</li> <li>cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.</li> </ul> <p><b>If pregnant or breast-feeding,</b> ask a health professional before use.</p> <p><b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.</p>	

<b>Drug Facts</b> (continued)	
<b>Directions</b>	
<ul style="list-style-type: none"> <li>Do not take more than 6 doses in any 24 hour period</li> <li>This adult strength product is not intended for use in children under 12 years of age</li> </ul>	
Age	Dose
adults and children 12 years and over	2 teaspoons every 4 hours
children under 12	do not use
<b>Other information</b>	
<ul style="list-style-type: none"> <li>Keep carton for full Direction for use.</li> <li>store at 20-25°C (68-77°F)</li> <li>do not refrigerate</li> <li>dosage cup provided</li> <li>sodium 3 mg per teaspoonful</li> </ul>	
<b>Inactive ingredients</b>	
anhydrous citric acid, FD&C red no. 40, glycerin, menthol, natural & artificial flavor, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose	
<b>Questions?</b> Call weekdays from 9:30 AM to 4:30 PM EST at <b>1-877-798-5944</b>	

\*This product is not manufactured or distributed by Pfizer, owner of the registered trademark Robitussin®

**DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING**



BX-004

If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.

DISTRIBUTED BY:  
CHAIN DRUG CONSORTIUM  
3301 NW BOCA RATON BLVD  
SUITE 101, BOCA RATON, FL 33431

NDC 68016-126-04

\*COMPARE TO THE ACTIVE INGREDIENTS IN ROBITUSSIN® PEAK COLD MULTI-SYMPTOM COLD



**Tussin CF**

Dextromethorphan HBr /  
Guaifenesin / Phenylephrine HCl

**COUGH SUPPRESSANT  
EXPECTORANT/  
NASAL DECONGESTANT**

NON-DROWSY

Multi-Symptom Formula  
For ages 12 & over

4 FL OZ (118 mL)



## Drug Facts

### Active ingredients (in each 5 mL tsp) Purposes

Dextromethorphan HBr, USP 10mg...Cough Suppressant  
Guaifenesin, USP 100 mg..... Expectorant  
Phenylephrine HCl, USP 5 mg.... Nasal Decongestant

### Uses

- help loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
  - nasal congestion
  - cough due to minor throat and bronchial irritation

### Warnings

**Do not use** if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

**Ask a doctor or pharmacist before use if you are** taking any other oral nasal decongestant or stimulant.

**When using this product do not use more than directed.**

### Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

## Drug Facts (continued)

### Directions

- Do not take more than 6 doses in any 24 hour period
- This adult strength product is not intended for use in children under 12 years of age

Age	Dose
adults and children 12 years and over	2 teaspoons every 4 hours
children under 12	do not use

### Other information

- Keep carton for full Direction for use.**
- store at 20-25°C (68-77°F)
- do not refrigerate
- dosage cup provided
- sodium 3 mg per teaspoonful

### Inactive ingredients

anhydrous citric acid, FD&C red no. 40, glycerin menthol, natural & artificial flavor, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

**Questions?** Call weekdays from 9:30 AM to 4:30 PM EST at **1-877-798-5944**

\*This product is not manufactured or distributed by Pfizer, owner of the registered trademark Robitussin®

**DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING**



BX-004

If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.

DISTRIBUTED BY:  
CHAIN DRUG CONSORTIUM  
3301 NW BOCA RATON BLVD  
SUITE 101, BOCA RATON, FL 33431

NDC 68016-126-04

\*COMPARE TO THE ACTIVE INGREDIENTS IN ROBITUSSIN® PEAK COLD MULTI-SYMP TOM COLD



# Tussin CF

Dextromethorphan HBr /  
Guaifenesin / Phenylephrine HCl

**COUGH SUPPRESSANT  
EXPECTORANT/  
NASAL DECONGESTANT**

NON-DROWSY

Multi-Symptom Formula  
For ages 12 & over

4 FL OZ (118 mL)



res

## TUSSIN CF

dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:68016-126
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 5 mL
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg in 5 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 5 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>MENTHOL</b> (UNII: L7T10EIP3A)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KOOR)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	

## Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CHERRY (cherry)	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68016-126-04	118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/15/2012	
2	NDC:68016-126-08	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/15/2012	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/15/2012	

**Labeler** - Chain Drug Consortium, LLC (101668460)

**Registrant** - AptaPharma Inc. (790523323)

## Establishment

Name	Address	ID/FEI	Business Operations
AptaPharma Inc.		790523323	manufacture(68016-126)