

COLD AND ALLERGY D- pseudoephedrine hcl, triprolidine hcl tablet, film coated
Walgreen Company

Walgreens 44-287

Active ingredients (in each tablet)

Pseudoephedrine HCl 60 mg
Triprolidine HCl 2.5 mg

Purpose

Nasal decongestant
Antihistamine

Uses

- temporarily relieves these symptoms due to hay fever (allergic rhinitis) or other upper respiratory allergies:
 - runny nose
 - itchy, watery eyes
 - nasal congestion
 - sneezing
 - itching of the nose or throat
- temporarily relieves these symptoms due to the common cold:
 - runny nose
 - sneezing
 - nasal congestion

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- high blood pressure
- heart disease
- thyroid disease
- diabetes
- glaucoma
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers.

When using this product

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- use caution when driving a motor vehicle or operating machinery
- drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or occur with fever

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours. Do not take more than 4 tablets in 24 hours.
- children under 12 years: do not use

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

hypromellose, magnesium stearate, polyethylene glycol, silicified microcrystalline cellulose, titanium dioxide

Questions or comments?

1-800-426-9391

Principal display panel

NDC 0363-0287-22

Walgreens

• WALGREENS •
PHARMACIST RECOMMENDED†

Cold & Allergy D

PSEUDOEPHEDRINE HCl / NASAL DECONGESTANT
TRIPROLIDINE HCl / ANTIHISTAMINE

Maximum Strength

• Relieves runny nose, nasal congestion,
sneezing & itchy, watery eyes

48

TABLETS

ONE TABLET PER DOSE

ACTUAL SIZE

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

†Our pharmacists recommend
the Walgreens brand.

50844 ORG112328722

DISTRIBUTED BY:

**WALGREEN CO.,
DEERFIELD, IL 60015**

walgreens.com ©2022 Walgreen Co.

W30RG1022-F
REV-0124



DISTRIBUTED BY:
WALGREENS CO.,
DEERFIELD, IL 60015
walgreens.com ©2022 Walgreen Co.

Questions or comments? 1-800-426-9391

Inactive ingredients
hypromellose, magnesium stearate, polyethylene glycol, stifted microcrystalline cellulose, titanium dioxide

Other information
TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED
OR BLISTER IS TORN OR BROKEN
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ see end flap for expiration date and lot number

Directions
■ do not take more than directed
adults and children 12 years and over: take 1 tablet every 4 to 6 hours. Do not take more than 4 tablets in 24 hours.
■ children under 12 years: do not use

Stop use and ask a doctor if
■ nervousness, dizziness, or sleeplessness occur
■ symptoms do not improve within 7 days or occur with fever
■ If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)
drowsiness may occur ■ avoid alcoholic beverages
■ excitability may occur, especially in children
■ use caution when driving a motor vehicle or operating machinery
■ alcohol, sedatives, and tranquilizers may increase drowsiness
Stop use and ask a doctor if

When using this product
■ do not exceed recommended dosage
or tranquilizers.

Ask a doctor or pharmacist before use if you are taking sedatives
■ difficulty in urination due to enlargement of the prostate gland
■ a breathing problem such as emphysema or chronic bronchitis

Ask a doctor before use if you have
■ high blood pressure
■ heart disease ■ thyroid disease ■ diabetes ■ glaucoma
■ a breathing problem such as emphysema or chronic bronchitis
MAOI, ask a doctor or pharmacist before taking this product.

Warnings
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Uses
temporarily relieves these symptoms due to hay fever
temporarily relieves these symptoms due to the common cold:
■ runny nose ■ sneezing ■ nasal congestion
■ itchy, watery eyes ■ nasal congestion
■ sneezing ■ itching of the nose or throat

Purpose
Pseudoephedrine HCl 60 mg... Nasal decongestant
Triprolidine HCl 2.5 mg... Antihistamine

Drug Facts
COMPLETE PRODUCT INFORMATION
KEEP OUTER PACKAGE FOR

Our pharmacists recommend the Walgreens brand.

50844 ORG112328722

No print area
Lot no/Exp date



TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

B-2201-287A-22-HR
ORG112328722

NDC 0363-0287-22

Walgreens



Cold & Allergy D

PSEUDOEPHEDRINE HCl / NASAL DECONGESTANT
TRIPROLIDINE HCl / ANTIHISTAMINE

Maximum Strength

- Relieves runny nose, nasal congestion, sneezing & itchy, watery eyes

48

TABLETS

ONE TABLET PER DOSE

ACTUAL SIZE



Walgreens 44-287A

COLD AND ALLERGY D

pseudoephedrine hcl, triprolidine hcl tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0287
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PSEUDOEPHEDRINE HYDROCHLORIDE (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	60 mg
TRIPROLIDINE HYDROCHLORIDE (UNII: YAN7R5L890) (TRIPROLIDINE - UNII:2L8T9S52QM)	TRIPROLIDINE HYDROCHLORIDE	2.5 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	287
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0287-22	2 in 1 CARTON	11/29/2022	
1		24 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	11/29/2022	

Labeler - Walgreen Company (008965063)**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(0363-0287)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(0363-0287) , pack(0363-0287)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(0363-0287)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(0363-0287)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(0363-0287)

Revised: 2/2024

Walgreen Company