UP AND UP CHILDRENS IBUPROFEN- ibuprofen suspension Target Corporation

Target Corporation Children's Ibuprofen Oral Suspension Drug Facts

Active ingredient (in each 5 mL = 1 teaspoonful)

Ibuprofen 100 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- give with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding
- feels faint
- vomits blood
- has bloody or black stools
- has stomach pain that does not get better
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed measuring cup
- if needed, repeat dose every 6-8 hours

- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance
- wash dosage cup after each use

| Dosing Chart | | | |
|---------------|------------------------------------|--------------------|--|
| Weight (lbs) | Age (yrs) Dose (teaspoonful or mL) | | |
| under 2 years | | ask a doctor | |
| 24-35 lbs | 2-3 years | 1 tsp or 5 mL | |
| 36-47 lbs | 4-5 years | 1 ½ tsp or 7.5 mL | |
| 48-59 lbs | 6-8 years | 2 tsp or 10 mL | |
| 60-71 lbs | 9-10 years | 2 ½ tsp or 12.5 mL | |
| 72-95 lbs | 11 years | 3 tsp or 15 mL | |

Other information

- each teaspoon contains: sodium 2 mg
- do not use if printed neckband is broken or missing
- store at 20-25°C (68-77°F)
- · do not freeze
- see bottom panel for lot number and expiration date

Inactive ingredients

anhydrous citric acid, D&C red #33, FD&C blue #1, FD&C red #40, glycerin, high fructose corn syrup, hypromellose, natural and artificial grape flavor, polysorbate 80, purified water, sodium benzoate, sorbitol solution, xanthan gum

Questions?

1-800-910-6874

Principal Display Panel

children's

ibuprofen oral suspension

100 mg per 5 mL (teaspoonful)

pain reliever/fever reducer (NSAID)


Compare to active ingredient in Children's Motrin® Grape Flavor

lasts up to 8 hours

alcohol free

GRAPE FLAVOR

AGE 2 to 11 YEARS



child has problems or serious side effects

child has problems or senous side effects from taking pain relievers or fever reducers child has not been drinking fluids
 child has lost a lot of fluid due to vomiting or diarrhea
 child has high blood pressure, heart disease, liver cirrhosis, or kidney disease child has extens.

Ask a doctor or pharmacist before use if the child is

under a doctor's care for any serious condition

When using this product
■ give with food or milk if stomach upset occurs
■ the risk of heart attack or stroke may increase if you use more than directed or for

Stop use and ask a doctor if a child experiences any of the following signs of stomach bleeding

feels faint

womits blood

has bloody or black stools

■ has stomach pain that does not get better the child does not get any relief within first day (24 hours) of treatment

fever or pain gets worse or lasts more than

days
 redness or swelling is present in the painful area
 any new symptoms appear

Directions

resistance

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

This product does not contain directions or complete warnings for adult use

do not give more than directed
shake well before using
find right dose on chart. If possible, use weight to dose; otherwise use age.

use only enclosed measuring cup
if needed, repeat dose every 6-8 hours
do not use more than 4 times a day
replace original bottle cap to maintain child resistance

vash dosage cup after each use

child has asthma

■ child is taking a diuretic

taking any other drug

longer than directed Stop use and ask a doctor if



Dosing Chart

Weight (lbs) Age (yrs) Dose (teaspoonful or mL)

each teaspoon contains: sodium 2 mg
 do not use if printed neckband is broken

Inactive ingredients anhydrous citric acid, D&C red #33, FD&C blue #1, FD&C red #40,

glycerin, high fructose corn syrup, hypromellose, natural and artificial grape flavor, polysorbate 80, purified water, sodium benzoate, sorbitol solution, xanthan gum

Questions? Call 1-800-910-6874

*This product is not manufactured or distributed

by McNeil Consumer Healthcare, distributor of Children's Motrin® Grape Flavor.

to make good products at good prices for good people. Enjoy!

Target gives 5% of its income to the community to support education, the

arts, social services and voluntee

■ do not use in printed neckband is bi or missing ■ store at 20-25°C (68-77°F) ■ do not freeze ■ see bottom panel for lot number and expiration date

24-35 lbs | 2-3 years

36-47 lbs 4-5 years

48-59 lbs 6-8 years

60-71 lbs 9-10 years

72-95 lbs 11 years

Other information

GLUTEN FREE

094 01 0245 ID201416

Distributed by Target Corporation

All Rights Reserved Shop Target.com

Minneapolis, MN 55403 ® & © 2012 Target Brands, Inc.

%.

ask a doctor

1 tsp or 5 mL

1 1/2 tsp or 7.5 mL

2 tsp or 10 mL

2 1/2 tsp or 12.5 mL

3 tsp or 15 ml

children's ibuprofen

oral suspension

100 mg per 5 mL (teaspoonful) pain reliever/fever reducer (NSAID)



children's

100 mg per 5 mL (teaspoonful) pain reliever/fever reducer (NSAID)



Important: Read all product information before using. Keep this box for important information. This product is intended for use in children. **Drug Facts**

Active ingredient Purposes

(in each 5 mL = 1 teaspoonful) Ibuprofen 100 mg (NSAID)*.....Pain relie .Pain reliever fever reducer nonsteroidal anti-inflammatory drug

- Uses temporarily:
 relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings
Allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

— hives

— hives

- hives facial swelling
 asthma (wheezing) shock
 skin reddening rash ■ blisters

■ blisters
If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems ■ takes a blood thinning (anticoagulant) or
- takes a blood thinning (anticoagulant) or steroid drug
 takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 takes more or for a longer time than directed Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult dector promptly De not use serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

■ if the child has ever had an allergic reaction to any other pain reliever/fever reducer right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn

NDC 11673-660-26

ibuprofen

oral suspension

Compare to active ingredient in Children's Motrin® Grape Flavor**

lasts up to 8 hours alcohol free



4 FL OZ (118 mL)



LOT NO.

EXP.

: PPOSP OM CA



UP AND UP CHILDRENS IBUPROFEN

ibuprofen suspension

| Product Information | | | |
|-------------------------|----------------------|--------------------|---------------|
| Product Type | HUMAN OTC DRUG LABEL | Item Code (Source) | NDC:11673-660 |
| Route of Administration | ORAL | DEA Sche dule | |
| | | | |

| Active Ingredient/Active Moiety | | | |
|---------------------------------|-------------------|----------------|--|
| Ingredient Name | Basis of Strength | Strength | |
| IBUPROFEN (IBUPROFEN) | IB UPRO FEN | 100 mg in 5 mL | |

| Inactive Ingredients | | | |
|----------------------|--|--|--|
| Strength | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |

| Product Characteristics | | | |
|-------------------------|-----------------|--------------|--|
| Color | PURPLE (opaque) | Score | |
| Shape | | Size | |
| Flavor | GRAPE | Imprint Code | |
| Contains | | | |

| Packaging | | | |
|--------------------|---------------------|----------------------|--------------------|
| # Item Code | Package Description | Marketing Start Date | Marketing End Date |
| 1 NDC:11673-660-26 | 1 in 1 CARTON | | |
| 1 | 118 mL in 1 BOTTLE | | |

| Marketing Information | | | |
|-----------------------|--|----------------------|--------------------|
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
| ANDA | ANDA074937 | 07/08/2009 | |
| | | | |

Labeler - Target Corporation (006961700)

Revised: 1/2013 Target Corporation