

**HISTEX-DM- dextromethorphan hydrobromide, phenylephrine hydrochloride,
and triprolidine hydrochloride syrup**
Allegis Pharmaceuticals, LLC

HISTEX™ -DM Syrup

Drug Facts

<i>Active ingredients (in each 5 mL teaspoonful)</i>	<i>Purpose</i>
Dextromethorphan HBr 20 mg	Cough Suppressant
Phenylephrine HCl 10 mg	Decongestant
Triprolidine HCl 2.5 mg	Antihistamine

Uses

temporarily relieves these symptoms due to common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- cough due to minor throat or bronchial irritation
- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes
- nasal congestion
- reduces swelling of nasal passages

Warnings

Do not exceed recommended dosage.

Do not use this product

- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- a persistent or chronic cough such occurs with smoking, asthma, chronic bronchitis, or emphysema
- a cough that occurs with too much phlegm (mucus)

- trouble urinating due to an enlarged prostate gland

Ask a doctor before use if you are taking sedatives or tranquilizers

When using this product

- excitability may occur, especially in children
- may cause drowsiness
- alcohol, sedatives and tranquilizers may increase the drowsiness effect
- avoid alcoholic beverages
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- cough or nasal congestion persists for more than 1 week, tends to recur or is accompanied by a fever, rash or persistent headache. These could be signs of a serious condition.
- new symptoms occur

If pregnant or breast-feeding, ask a health professional before use.

Keep out of the reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Do not exceed recommended dosage.

AGE	DOSE
Adults and Children 12 years of age and older:	1 teaspoonful (5 mL) every 4 hours, not to exceed 4 teaspoonfuls (20 mL) in 24 hours, or as directed by a doctor
Children 6 to under 12 years of age	½ teaspoonful (2.5 mL) every 4 hours, not to exceed 2 teaspoonfuls (10 mL) in 24 hours, or as directed by a doctor.
Children under 6 years of age	Consult a doctor

Other Information

Store at 15°-30° C (59°-86° F).

Tamper evident by foil seal under cap. Do not use if foil seal is missing or broken.

Dispense in a tight, light-resistant container with a child-resistant cap.

Inactive ingredients

Citric Acid, Glycerin, Propylene Glycol, Purified Water, Sodium Citrate, Sodium Saccharin, Sorbitol, Grape Flavor.

Questions? Comments?

Call 1-866-633-9033.

PRINCIPAL DISPLAY PANEL - 473 mL Bottle Label

NDC 28595-804-16

Antihistamine • Decongestant

Cough Suppressant

HISTEX™ -DM

Syrup

Each teaspoonful (5 mL)

contains:

Dextromethorphan HBr 20 mg

Phenylephrine HCl 10 mg

Tripolidine HCl 2.5 mg

Sugar-Free • Dye Free

Alcohol Free

Grape Flavor

16 fl oz (473 mL)

NDC 28595-804-16

Antihistamine • Nasal Decongestant
Cough Suppressant

HISTEX-DMTM Syrup

Each teaspoonful (5 mL)
contains:

Dextromethorphan HBr ... 20 mg
Phenylephrine HCl 10 mg
Triprolidine HCl 2.5 mg

Sugar-Free • Dye Free
Alcohol Free

Grape Flavor

Tamper evident by foil seal under cap.
Do not use if foil seal is missing or broken.



16 fl oz (473 mL)

Drug Facts

Active ingredients
(in each 5 mL teaspoonful)

Dextromethorphan HBr 20 mg	Cough Suppressant
Phenylephrine HCl 10 mg	Nasal Decongestant
Triprolidine HCl 2.5 mg	Antihistamine

Purpose

Uses temporarily relieves these symptoms due to common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- cough due to minor throat and bronchial irritation
- runny nose ■ sneezing ■ itching of the nose or throat
- itchy, watery eyes ■ nasal congestion
- reduces swelling of nasal passages

Warnings

Do not exceed recommended dosage.

Do not use this product

- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- heart disease ■ high blood pressure
- thyroid disease ■ diabetes ■ glaucoma
- a persistent or chronic cough such occurs with smoking, asthma, chronic bronchitis, or emphysema
- a cough that occurs with too much phlegm (mucus)
- trouble urinating due to an enlarged prostate gland

Ask a doctor before use if you are taking sedatives or tranquilizers

When using this product

- excitability may occur, especially in children
- may cause marked drowsiness
- alcohol, sedatives and tranquilizers may increase the drowsiness effect
- avoid alcoholic beverages

Drug Facts (continued)

- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- cough or nasal congestion persists for more than 1 week, tends to recur or is accompanied by a fever, rash or persistent headache. These could be signs of a serious condition.
- new symptoms occur

If pregnant or breast-feeding, ask a health professional before use.

Keep out of the reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions Do not exceed recommended dosage.

AGE	DOSE
Adults and Children 12 years of age and older	1 teaspoonful (5 mL) every 4 hours, not to exceed 4 teaspoonfuls (20 mL) in 24 hours, or as directed by a doctor
Children 6 to under 12 years of age	½ teaspoonful (2.5 mL) every 4 hours, not to exceed 2 teaspoonfuls (10 mL) in 24 hours, or as directed by a doctor.
Children under 6 years of age	Consult a doctor

Other Information

Store at 15°-30° C (59°-86° F).

Tamper evident by foil seal under cap. Do not use if foil seal is missing or broken.

Inactive ingredients Citric acid, glycerin, grape flavor, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium saccharin, sorbitol solution

Questions? Comments? Call 1-866-633-9033.

Manufactured for: Allegris Pharmaceuticals, LLC
www.allegrispharma.com Canton, MS 39046

Rev. 03/22

HISTEX-DM

dextromethorphan hydrobromide, phenylephrine hydrochloride, and triprolidine hydrochloride syrup

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:28595-804
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 5 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 5 mL
TRIPROLIDINE HYDROCHLORIDE (UNII: YAN7R5L890) (TRIPROLIDINE - UNII:2L8T9S52QM)	TRIPROLIDINE HYDROCHLORIDE	2.5 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	

SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SORBITOL (UNII: 506T60A25R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:28595-804-16	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/19/2014	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/19/2014	

Labeler - Allegis Pharmaceuticals, LLC (792272861)

Revised: 1/2024

Allegis Pharmaceuticals, LLC