

LAXATIVE ENTERIC COATED- bisacodyl tablet, coated
AmerisourceBergen Drug Corporation (Good Neighbor Pharmacy) 24385

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredient (in each tablet)

Bisacodyl USP 5 mg

Purpose

Stimulant laxative

Use

- for temporary relief of occasional constipation and irregularity
- this product generally produces a bowel movement in 6 to 12 hours

Warnings

Do not use

if you cannot swallow without chewing.

Ask a doctor before use if you have

- stomach pain, nausea or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

When using this product

- do not chew or crush tablet(s)
- do not use within 1 hour after taking an antacid or milk
- it may cause stomach discomfort, faintness and cramps

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take with a glass of water

adults and children 12 years of age and over	take 1 to 3 tablets in a single daily dose
children 6 to under 12 years of age	take 1 tablet in a single daily dose
children under 6 years of age	ask a doctor

Other information

- store between 20°-25°C (68°-77°F)

Inactive ingredients

acacia, anhydrous calcium sulfate, anhydrous lactose, carnauba wax, colloidal silicon dioxide, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, gelatin, iron oxide, iron oxide black, iron oxide yellow, magnesium stearate, microcrystalline cellulose, polyethylene glycol (PEG) 400, polyvinyl acetate phthalate, povidone, shellac, sodium starch glycolate, stearic acid, sugar, talc, titanium dioxide

Questions or Comments?

Call **1-877-753-3935 Monday-Friday 9AM-5PM EST**

Principal Display Panel

*Compare to Dulcolax® active ingredient

Laxative

Bisacodyl, 5 mg

Gentle, Predictable Overnight Relief

Enteric coated tablets

*This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals, Inc., distributor of Dulcolax® Laxative.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Distributed By AmerisourceBergen

1300 Morris Drive, Chesterbrook, PA 19087

Visit us at www.mygnp.com

Package Label

***Compare to Dulcolax® Laxative active Ingredient**



**GOOD
NEIGHBOR
PHARMACY**

NDC 24385-903-63

Laxative

bisacodyl, 5 mg

Gentle, Predictable Overnight Relief

25 enteric coated tablets



TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Lot / Exp.:

Product of:

*This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals, Inc., distributor of Dulcolax® Laxative.

Distributed By AmerisourceBergen
1300 Morris Drive, Chesterbrook, PA 19087
Questions or Concerns?
Visit us at www.mygnp.com



PLD-D23K
FC003324

ABC#10098892



Questions or comments? Call
1-877-753-3935 Monday-Friday 9AM-5PM EST

Inactive ingredients acacia, anhydrous calcium sulfate, anhydrous lactose, carnauba wax, colloidal silicon dioxide, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, gelatin, iron oxide black, iron oxide yellow, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl acetate phthalate, povidone, shellac, sodium starch glycolate, stearic acid, sugar, talc, titanium dioxide

Other information
■ store between 20-25°C (68-77°F)

- When using this product**
- a sudden change in bowel habits that lasts more than 2 weeks
 - do not chew or crush tablet(s)
 - do not use within 1 hour after taking an antacid or milk
 - it may cause stomach discomfort, faintness and cramps
- Stop use and ask a doctor if**
- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
 - you need to use a laxative for more than 1 week

■ stomach pain, nausea or vomiting

Ask a doctor before use if you have

Do not use if you cannot swallow without chewing.

Warnings

■ this product generally produces a bowel irregularity

■ for temporary relief of occasional constipation and movement in 6 to 12 hours

Use

Bisacodyl USP 5 mg.....Stimulant laxative

(in each tablet)

Active ingredient

Purpose

Drug Facts

(continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

■ take with a glass of water

adults and children 12 years of age and over

take 1 to 3 tablets in a single daily dose

children 6 to under 12 years of age

take 1 tablet in a single daily dose

children under 6 years of age

ask a doctor

Drug Facts

(continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

■ take with a glass of water

adults and children 12 years of age and over

take 1 to 3 tablets in a single daily dose

children 6 to under 12 years of age

take 1 tablet in a single daily dose

children under 6 years of age

ask a doctor

Good Neighbor Pharmacy Laxative

LAXATIVE ENTERIC COATED

bisacodyl tablet, coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:24385-903
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
BISACODYL (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y)		BISACODYL	5 mg
Inactive Ingredients			
Ingredient Name			Strength
ACACIA (UNII: 5C5403N26O)			
CALCIUM SULFATE ANHYDROUS (UNII: E934B3V59H)			
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)			
CARNAUBA WAX (UNII: R12CBM0EIZ)			
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
STARCH, CORN (UNII: O8232NY3SJ)			
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)			
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)			
ALUMINUM OXIDE (UNII: LMI26O6933)			
GELATIN (UNII: 2G86QN327L)			
FERRIC OXIDE RED (UNII: 1K09F3G675)			
FERROSOFERRIC OXIDE (UNII: XM0M87F357)			
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)			

POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
Polyvinyl Acetate Phthalate (UNII: 58QVG85GW3)	
POVIDONE (UNII: FZ989GH94E)	
SHELLAC (UNII: 46N107B71O)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
SUCROSE (UNII: C151H8M554)	

Product Characteristics

Color	orange, yellow	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	TCL003
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:24385-903-63	1 in 1 CARTON	10/24/2011	12/27/2024
1		25 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	10/24/2011	12/27/2024

Labeler - AmerisourceBergen Drug Corporation (Good Neighbor Pharmacy) 24385 (007914906)

Revised: 9/2022

AmerisourceBergen Drug Corporation (Good Neighbor Pharmacy) 24385