

UP AND UP ACETAMINOPHEN- acetaminophen tablet, film coated, extended release
Target Corporation

Target Corporation Acetaminophen Drug Facts

Active ingredient (in each caplet)

Acetaminophen 650 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- premenstrual and menstrual cramps
- the common cold
- headache
- toothache
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

liver disease

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)

adults	<ul style="list-style-type: none">• take 2 caplets every 8 hours with water• swallow whole; do not crush, chew, split or dissolve• do not take more than 6 caplets in 24 hours• do not use for more than 10 days unless directed by a doctor
under 18 years of age	<ul style="list-style-type: none">• ask a doctor

Other information

- store at 20-25°C (68-77°F). Avoid excessive heat 40°C (104°F).

Inactive ingredients

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, hypromellose, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, pregelatinized starch, stearic acid, titanium dioxide

Questions? Call

1-888-547-7400

Principal Display Panel

see new warning

Compare to active ingredient in Tylenol® 8HR Arthritis Pain
arthritis pain

acetaminophen extended-release tablets, 650 mg

pain reliever / fever reducer

for the temporary relief of minor arthritis pain

DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

100 CAPLETS*

100 CAPLETS* 650 mg EACH

*(CAPSULE-SHAPED TABLETS)

see new warning NDC 11673-966-78

Compare to active ingredient in
Tylenol® 8HR Arthritis Pain**

arthritis pain
acetaminophen
extended-release tablets, 650 mg
pain reliever/fever reducer
for the temporary relief of minor arthritis pain

up&up

DO NOT USE WITH OTHER
MEDICINES CONTAINING
ACETAMINOPHEN

THIS PACKAGE FOR
HOUSEHOLDS WITHOUT
YOUNG CHILDREN

100
CAPLETS*

100 CAPLETS* 650 mg EACH
*(CAPSULE-SHAPED TABLETS)

Drug Facts (continued)
Questions? Call
1-888-547-7400

**This product is not manufactured or
distributed by McNeil Consumer Healthcare,
distributor of Tylenol® 8HR Arthritis Pain.
U.S. Patent 7,997,172

GLUTEN FREE

094 01 0496 R01
Distributed by Target Corporation
Minneapolis, MN 55403
© 2015 Target Brands, Inc.
All Rights Reserved Shop Target.com

Contains No Aspirin
**DO NOT USE IF PRINTED FOIL
UNDER CAP IS BROKEN OR MISSING**

: 96678 UW F2

Drug Facts

Active ingredient (in each caplet)	Purpose
---	----------------

Acetaminophen 650 mg.....	Pain reliever/ fever reducer
------------------------------	---------------------------------

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches ■ backache
 - premenstrual and menstrual cramps
 - the common cold ■ headache ■ toothache
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur

■ redness or



swelling is
present
These could
be signs of
a serious
condition. ▶

3 **PEEL BACK HERE ▶**

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not take more than directed (see overdose warning)

adults	■ take 2 caplets every 8 hours with water ■ swallow whole; do not crush, chew, split or dissolve ■ do not take more than 6 caplets in 24 hours ■ do not use for more than 10 days unless directed by a doctor
under 18 years of age	■ ask a doctor

Other information ■ store at 20-25°C (68-77°F). Avoid excessive heat 40°C (104°F).

Inactive ingredients carnauba wax, colloidal silicon dioxide, croscarmellose sodium, hypromellose, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, pregelatinized starch, stearic acid, titanium dioxide ▼

UP AND UP ACETAMINOPHEN

acetaminophen tablet, film coated, extended release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-966
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTO DEXTRIN (UNII: 7CVR7L4A2D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POVIDONE (UNII: FZ989GH94E)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	WHITE	Score	no score
Shape	CAPSULE	Size	19mm
Flavor		Imprint Code	L544
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-966-78	100 in 1 BOTTLE; Type 0: Not a Combination Product	03/15/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075077	03/18/2015	

