ADAPALENE- adapalene gel Walgreens

Adapalene Gel USP 0 1 percent 1 6 oz

Active ingredient

Adapalene USP 0.1% (retinoid)*

*read consumer information leaflet

Purpose

Acne treatment

Use

• For the treatment of acne

Warnings

For external use only

Do not use

- on damaged skin (cuts, abrasions eczema, sunburn)
- if you are allergic to adapalene or any of the ingredients in this product

Pregnancy/Breastfeeding,

ask a doctor before use.

When using this product

- limit sun exposure, including light from tanning beds, and use sunscreen when going outdoors.
- do not wax to remove hair in areas where the product has been applied
- during the early weeks of use, your acne may appear to worsen before it improves (this is normal); continue using as directed, unless you get irritation that become severe
- irritation (redness, itching, dryness, burning) is more likely to occur:
 - in the first few weeks of use
 - if using more than one topical acne medication at a time
 - but irritation usually lessens with continued use of this product
- it may take up to 3 months of once daily use to see result
- avoid product contact eyes, lips, and month. If contact occurs immediately flush the area with water.
- wash hands after use

Stop use and ask a doctor if

- you become pregnant, or planning to become pregnant while using this product
- you have symptoms of an allergic reaction (such as itching, rash, hives, swelling of the lips, eyelids, and shortness of breath)
- irritation become severe
- you see no improvement after 3 months of once daily use

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

Adults and children 12 years of age and older:

- use **once** daily
- clean the skin gently and pat dry before applying the product
- cover the entire affected area with a thin layer. For example, if your acne in on the face, apply the product to the entire face.
- do not use more than one time a day. Applying more than directed will not provide faster or better results, but may worsen skin irritation.

Children under 12 years of age: ask a doctor

Other information

- store at rooms temperature 68° to $77^{\circ}F$
- protect from freezing

Inactive ingredients

carbomer homopolymer typeC, edetate disodium, methylparabem, poloxamer 182, propylene glycol, purified water. May contain hydrochloric acid and/or sodium hydroxide to adjust sodium hydroxide pH.

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to the active ingredient in Differin® Gelt

Adapalene Gel

USP 0.1%

ACNE TREATMENT

Oil Free

Fragrance Free

- Previously available only by prescription
- Dermatologist developed & tested
- Once daily topical retinoid*

NET WT OZ (g)

***READ CONSUMER INFORMATION LEAFLET BEFORE USE.**

*Read carton and enclosed consumer information leaflet before using this product. Keep this carton and consumer information leaflet. They contain important information.

TAMPER EVIDENT: DO NOT USE IF SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

<code>++This product</code> is not manufactured or distributed by Galderma Laboratories, L.P. distributor of Differin® Gel.

DISTRIBUTED BY WALGREEN CO.

DEERFIELD, IL 60015

Product Label

Drug Facts

Active ingredient Purpose Adapalene, USP 0.1% (retinoid)*. Acne treatment *read consumer information leaflet

Use For the treatment of acne

Warnings

For external use only

Do not use

 on damaged skin (cuts, abrasions, eczema, sunburn) If you are allergic to adapalene or any of the ingredients in this product

If pregnant or breast-feeding, ask a doctor before use.

When using this product

- Imit sun exposure, including light from tanning beds, and use sunscreen when going outdoors.
- do not wax to remove hair in areas where the product has been applied
- during the early weeks of use, your acne may appear to worsen before it improves (this is normal); continue using
- as directed, unless you get irritation that becomes severe irritation (redness, itching, dryness, burning) is more likely to occur:
- in the first few weeks of use
- if using more than one topical acne medication at a time but irritation usually lessens with continued use of this product
- it may take up to 3 months of once daily use to see results avoid product contact with eyes, lips, and mouth. If contact occurs, immediately flush the area with water. wash hands after use

Stop use and ask a doctor if

- vou become pregnant, or are planning to become pregnant
- while using the product you have symptoms of an allergic reaction (such as itching, rash, hives, swelling of the lips, eyelids, and shortness of breath)
- Irritation becomes severe

vou see no improvement after 3 months of once daily use

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

Adults and children 12 years of age and older: use once daily

- clean the skin gently and pat dry before applying the product cover the entire affected area with a thin layer. For example, if
- your acne is on the face, apply the product to the entire face. do not use more than one time a day. Applying more than
- directed will not provide faster or better results, but may worsen skin irritation.

Children under 12 years of age: ask a doctor

Other information

store at room temperature 68° to 77°F protect from freezing

Inactive ingredients carbomer homopolymer type C, edetate disodium, methylparaben, poloxamer 182, propylene glycol, purified water. May contain hydrochloric acid and/or sodium hydroxide to adjust pH.

Questions or comments? Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

Our pharmacists recommend the Walgreens brand. We This product is not manufactured or distributed by Galderma Laboratories, L.P., distributor of Differin® Gel. Read carton and enclosed consumer information leaflet before using this product. Keep this carton and consumer information leaflet. They contain important information.

w

Walgreens Compare to the active ingredient in Differin® Gel"

Adapalene Gel USP 0.1%

ACNE TREATMENT

Oil Free

Fragrance Free

- Previously available only by prescription
- Dermatologist developed & tested
- Once daily topical retinoid*



DISTRIBUTED BY: WALGREEN CO. DEERFIELD, IL 60015 100% SATISFACTION GUARANTEED walgreens.com ©2021 Walgreen Co.

W3ORG1221-F **REV-0923**

ACTUAL SIZE **ONE TUBE INSIDE**



NET WT 1.6 OZ (45 g)



WALGREENS Acne Treatment

ADAPALENE

adapalene gel

	oduct Infor						
Pr	oduct Type		HUMAN OTC DRUG	Item Co	ode (Source)	NDC:0	0363-1401
Ro	oute of Admini	stration	TOPICAL				
Ac	tive Ingredi	ent/Active	Moietv				
Ingredient Name					Basis of Strength Strengt		Strength
ADAPALENE (UNII: 1L4806J2QF) (A					ADAPALENE		
In	active Ingre	dients					
			Ingredient Na	me			Strengt
	RBOMER HOMO 93RCW27E)	POLYMER TYP	E C (ALLYL PENTAERY)	HRITOL C	ROSSLINKED) (UNII:		
ED	ETATE DISODIU	M (UNII: 7FLD91	.С86К)				
	THYLPARABEN		•				
	LOXAMER 182	UNII: IX0HIX6OA	G)				
		-					
PR	OPYLENE GLYC	OL (UNII: 6DC90					
PR W/		OL (UNII: 6DC90 F0KO0R))167V3)				
PR W/ SO	OPYLENE GLYC Ater (Unii: 059Q	OL (UNII: 6DC90 F0KO0R))167V3)				
PR W/ SO	OPYLENE GLYC Ater (UNII: 059Q Dium Hydroxi	OL (UNII: 6DC90 F0KO0R) DE (UNII: 55X04)167V3)		Marketing Start Date	Mar	keting End Date
PR W/ SO Pa #	OPYLENE GLYC ATER (UNII: 059Q DIUM HYDROXI	OL (UNII: 6DC90 F0KO0R) DE (UNII: 55X04	2167V3) QC32I)		-	Mar	_
PR W/ SO Pa #	OPYLENE GLYC ATER (UNII: 059Q DIUM HYDROXI ICKaging Item Code NDC:0363-1401-	OL (UNII: 6DC90 FOKOOR) DE (UNII: 55X04 Pac 1 in 1 CARTON	2167V3) QC32I)	10,	Date	Mar	_
PR W/ SO Pa #	OPYLENE GLYC ATER (UNII: 059Q DIUM HYDROXI ICKaging Item Code NDC:0363-1401-	OL (UNII: 6DC90 FOKOOR) DE (UNII: 55X04 Pac 1 in 1 CARTON 45 g in 1 TUBE	QC32I) kage Description ; Type 0: Not a Combinat	ion	Date	Mar	_
PR W/ SO Pa # 1 1	OPYLENE GLYC ATER (UNII: 059Q DIUM HYDROXI Item Code NDC:0363-1401- 16	OL (UNII: 6DC90 FOKOOR) DE (UNII: 55X04 Pac 1 in 1 CARTON 45 g in 1 TUBE Product 1 in 1 CARTON	QC32I) kage Description ; Type 0: Not a Combinat	ion 10,	Date /25/2023	Mar	_
PR W/ SO Pa # 1 1	OPYLENE GLYC ATER (UNII: 059Q DIUM HYDROXI Item Code NDC:0363-1401- 16	OL (UNII: 6DC90 FOKOOR) DE (UNII: 55X04 Pac 1 in 1 CARTON 45 g in 1 TUBE Product 1 in 1 CARTON 15 g in 1 TUBE	QC32I) kage Description	ion 10,	Date /25/2023	Mar	_
PR W/ SO Pa # 1 1 2 2	OPYLENE GLYC ATER (UNII: 059Q DIUM HYDROXI Item Code NDC:0363-1401- 16	OL (UNII: 6DC90 FOKOOR) DE (UNII: 55X04 Pac 1 in 1 CARTON 45 g in 1 TUBE Product 1 in 1 CARTON 15 g in 1 TUBE Product	QC32I) kage Description ; Type 0: Not a Combinat	ion 10,	Date /25/2023	Mar	_
PR W/ SO Pa # 1 1 2 2	OPYLENE GLYC ATER (UNII: 059Q DIUM HYDROXI Item Code NDC:0363-1401- 16 NDC:0363-1401- 05	OL (UNII: 6DC90 FOKOOR) DE (UNII: 55X04 Pac 1 in 1 CARTON 45 g in 1 TUBE Product 1 in 1 CARTON 15 g in 1 TUBE Product	QC32I) kage Description ; Type 0: Not a Combinat	ion 10, ion 10, ion	Date /25/2023		_

Labeler - Walgreens (008965063)

Revised: 10/2023

Walgreens