

**LEADER LOPERAMIDE HYDROCHLORIDE- loperamide hydrochloride liquid**  
**Cardinal Health**

-----  
**Cardinal Health Loperamide Hydrochloride Oral Suspension Drug Facts**

**Active ingredient (in each 7.5 mL)**

Loperamide HCl 1 mg

**Purpose**

Anti-diarrheal

**Use**

controls symptoms of diarrhea, including Travelers' Diarrhea

**Warnings**

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

**Do not use**

if you have bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool
- a history of liver disease

**Ask a doctor or pharmacist before use if you are**

taking antibiotics

**When using this product**

- tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging.

These may be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## Directions

- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- shake well before using
- only use attached measuring cup to dose product

adults and children 12 years and over	30 mL (6 tsp) after the first loose stool; 15 mL (3 tsp) after each subsequent loose stool; but no more than 60 mL (12 tsp) in 24 hours
children 9-11 years (60-95 lbs)	15 mL (3 tsp) after the first loose stool; 7.5 mL (1 1/2 tsp) after each subsequent loose stool; but no more than 45 mL (9 tsp) in 24 hours
children 6-8 years (48-59 lbs)	15 mL (3 tsp) after the first loose stool; 7.5 mL (1 1/2 tsp) after each subsequent loose stool; but no more than 30 mL (6 tsp) in 24 hours
children under 6 years (up to 47 lbs)	ask a doctor

## Other information

- each 30 mL (6 tsp) contains: sodium 13 mg
- store between 20-25°C (68-77°F)
- do not use if printed plastic neckband is broken or missing
- see side panel for lot number and expiration date

## Inactive ingredients

anhydrous citric acid, carboxymethylcellulose sodium, D&C yellow no. 10, FD&C blue no. 1, glycerin, microcrystalline cellulose, natural and artificial mint flavor, propylene glycol, purified water, simethicone, sodium benzoate, sucralose, titanium dioxide, xanthan gum

## Questions or comments?

1-800-719-9260

## Package/Label Principal Display Panel

Compare to Imodium® A-D active ingredient

Loperamide Hydrochloride Oral Suspension

Anti-Diarrheal

Controls the Symptoms of Diarrhea

Mint Flavor

1 mg Loperamide Hydrochloride per 7.5 mL

**Drug Facts**

**Active ingredient Purpose**  
**(in each 7.5 mL)**

Loperamide HCl  
1 mg.....Anti-diarrheal

**Use** controls symptoms of diarrhea, including Travelers' Diarrhea

**Warnings**

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

**Do not use** if you have bloody or black stool

**PEEL BACK HERE**

NDC 37205-584-26



Compare to **Imodium® A-D** active ingredient\*

**Loperamide Hydrochloride Oral Suspension**

**Anti-Diarrheal**

Controls the Symptoms of Diarrhea

Mint Flavor



1 mg Loperamide Hydrochloride per 7.5 mL

**4 FL OZ (120 mL)**

DISTRIBUTED BY  
CARDINAL HEALTH  
DUBLIN, OHIO 43017  
CIN 4505830  
www.myleader.com  
1-800-200-6313

**Gluten Free**

All **Leader®** Brand products are 100% satisfaction guaranteed or return to place of purchase for a full refund

\*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Imodium® A-D.



0 96295 12100 1

: 64526 E9 F1

**Drug Facts (continued)**

**Ask a doctor before use if you have** ■ fever ■ mucus in the stool ■ a history of liver disease

**Ask a doctor or pharmacist before use if you are taking antibiotics**

**When using this product** ■ tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison

**Drug Facts (continued)**

Control Center right away. (1-800-222-1222)

**Directions** ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

- find right dose on chart. If possible, use weight to dose; otherwise use age. ■ shake well before using ■ only use attached measuring cup to dose product

adults and children 12 years and over 30 mL (6 tsp) after the first loose stool; 15 mL (3 tsp) after each subsequent loose stool; but no more than 60 mL (12 tsp) in 24 hours

children 9-11 years (60-95 lbs) 15 mL (3 tsp) after the first loose stool; 7.5 mL (1 1/2 tsp) after each subsequent loose stool; but no more than 45 mL (9 tsp) in 24 hours

**Drug Facts (continued)**

children 6-8 years (48-59 lbs) 15 mL (3 tsp) after the first loose stool; 7.5 mL (1 1/2 tsp) after each subsequent loose stool; but no more than 30 mL (6 tsp) in 24 hours

children under 6 years (up to 47 lbs) ask a doctor

**Other information**

- each 30 mL (6 tsp) contains: sodium 13 mg
- store between 20-25°C (68-77°F)

**Drug Facts (continued)**

■ **do not use if printed plastic neckband is broken or missing** ■ see side panel for lot number and expiration date

**Inactive ingredients** anhydrous citric acid, carboxymethylcellulose sodium, D&C yellow no. 10, FD&C blue no. 1, glycerin, microcrystalline cellulose, natural and artificial mint flavor, propylene glycol, purified water, simethicone, sodium benzoate, sucralose, titanium dioxide, xanthan gum

**Questions or comments?**  
1-800-719-9260

# LEADER LOPERAMIDE HYDROCHLORIDE

loperamide hcl liquid

## Product Information

Product Type	HUMAN OTC DRUG LABEL	Item Code (Source)	NDC:37205-584
Route of Administration	ORAL	DEA Schedule	

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (LOPERAMIDE)	LOPERAMIDE HYDROCHLORIDE	1 mg in 7.5 mL

## Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID	
CARBOXYMETHYLCELLULOSE SODIUM	
D&C YELLOW NO. 10	
FD&C BLUE NO. 1	
GLYCERIN	
CELLULOSE, MICROCRYSTALLINE	
PROPYLENE GLYCOL	
WATER	
SODIUM BENZOATE	
SUCRALOSE	
TITANIUM DIOXIDE	
XANTHAN GUM	

## Product Characteristics

Color	GREEN (opaque, viscous)	Score	
Shape		Size	
Flavor	MINT	Imprint Code	
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37205-584-26	120 mL in 1 BOTTLE		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091292	03/28/2012	

