

**TOPCARE PAIN AND FEVER CHILDRENS- acetaminophen suspension**  
**Topco Associates LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Topco Associates LLC. Children's Pain & Fever Drug Facts**

**Active ingredient (in each 5 mL)**

Acetaminophen 160 mg

**Purpose**

Pain reliever/fever reducer

**Uses**

temporarily:

- reduces fever
- relieves minor aches and pains due to:
- the common cold
- flu
- headache
- sore throat
- toothache

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not

sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if your child has ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if your child has**

liver disease

**Ask a doctor or pharmacist before use if your child is**

taking the blood thinning drug warfarin

**When using this product**

**do not exceed recommended dose (see overdose warning)**

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

**Directions**

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed (see overdose warning)**
- **shake well before using**
- mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

<b>Weight (lb)</b>	<b>Age (yr)</b>	<b>Dose (mL)*</b>
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL
72-95	11 years	15 mL

\*or as directed by a doctor

**Attention:** use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

### **Other information**

- **each 5 mL contains:** sodium 3 mg
- store at 20-25°C (68-77°F)
- **do not use if printed neckband is broken or missing**

### **Inactive ingredients**

anhydrous citric acid, calcium sulfate, carrageenan, flavor, glycerin, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, propylparaben, purified water, sodium benzoate, sorbitol solution, sucralose, tribasic sodium phosphate

### **Questions or comments?**

**1-888-423-0139**

### **Package/Label Principal Display Panel**

COMPARE TO CHILDREN'S TYLENOL® ORAL SUSPENSION ACTIVE INGREDIENT

children's Pain & Fever

ORAL SUSPENSION

ACETAMINOPHEN, 160 mg per 5 mL

PAIN RELIEVER - FEVER REDUCER

OUR PHARMACISTS RECOMMEND

DYE-FREE

Aspirin Free

Ibuprofen Free

Alcohol Free

Ages 2 to 11 Years

DYE-FREE

GRAPE FLAVOR

4 FL OZ (118 mL)



## TOPCARE PAIN AND FEVER CHILDRENS

acetaminophen suspension

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:36800-397
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength		
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL		
<b>Inactive Ingredients</b>				
Ingredient Name	Strength			
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)				
CALCIUM SULFATE, UNSPECIFIED FORM (UNII: WAT0DDB505)				
CARRAGEENAN (UNII: 5C69YCD2YJ)				
GLYCERIN (UNII: PDC6A3C0OX)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K679OBS311)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
PROPYLPARABEN (UNII: Z8IX2SC1OH)				
WATER (UNII: 059QF0K00R)				
SODIUM BENZOATE (UNII: OJ245FE5EU)				
SORBITOL (UNII: 506T60A25R)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
SODIUM PHOSPHATE, TRIBASIC (UNII: A752Q30A6X)				
<b>Product Characteristics</b>				
Color		Score		
Shape		Size		
Flavor	GRAPE	Imprint Code		
Contains				
<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-397-26	1 in 1 CARTON	02/28/2012	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph not final	part343	02/28/2012		

**Labeler** - Topco Associates LLC (006935977)

Revised: 12/2019

Topco Associates LLC