

VALRUBICIN INTRAVESICAL SOLUTION- valrubicin intravesical solution solution, concentrate
Hikma Pharmaceuticals USA Inc.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use VALRUBICIN INTRAVESICAL SOLUTION safely and effectively. See full prescribing information for VALRUBICIN INTRAVESICAL SOLUTION.

VALRUBICIN intravesical solution, for intravesical use
Initial U.S. Approval: 1981

INDICATIONS AND USAGE

- Valrubicin intravesical solution is an anthracycline topoisomerase inhibitor indicated for intravesical therapy of BCG-refractory carcinoma *in situ* (CIS) of the urinary bladder in patients for whom immediate cystectomy would be associated with unacceptable morbidity or mortality. (1)

DOSAGE AND ADMINISTRATION

- **For Intravesical Use Only.** (2.1)
- Valrubicin intravesical solution is recommended at a dose of 800 mg administered intravesically once a week for six weeks. (2.1)
- Delay administration at least two weeks after transurethral resection and/or fulguration. (2.1)
- Warm valrubicin intravesical solution slowly to room temperature, but do not heat. (2.1)
- Use caution when handling and preparing the solution of valrubicin intravesical solution. (2.2)

DOSAGE FORMS AND STRENGTHS

- Injection: 200 mg/5 mL in single-use vials (3)

CONTRAINDICATIONS

- Perforated bladder or compromised bladder mucosa (4, 5.2)
- Hypersensitivity to anthracyclines or polyoxyl castor oil. (4)
- Concurrent urinary tract infections. (4)
- Patients with a small bladder capacity unable to tolerate a 75 mL instillation. (4)

WARNINGS AND PRECAUTIONS

- Delaying cystectomy can lead to development of metastatic bladder cancer, which is lethal. (5.1)
- Do not administer valrubicin intravesical solution to patients with a perforated bladder or to those in whom the integrity of the bladder mucosa has been compromised. (5.2, 12.3)
- Evaluate the status of the bladder before the intravesical instillation of valrubicin intravesical solution. (5.3)
- Use with caution in patients with severe irritable bladder symptoms. (5.4)
- Embryo-Fetal Toxicity: Valrubicin intravesical solution can cause fetal harm. Advise females of reproductive potential of the potential risk to a fetus and to use effective contraception. (5.5, 8.1, 8.3)

ADVERSE REACTIONS

The most frequently reported adverse reactions ($\geq 5\%$) after administration of valrubicin intravesical solution are urinary frequency, dysuria, urinary urgency, bladder spasm, hematuria, bladder pain, urinary incontinence, cystitis, urinary tract infection, nocturia, local burning symptoms, abdominal pain, and nausea. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Hikma at us.hikma@primevigilance.com or call 1-877-845-0689 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

USE IN SPECIFIC POPULATIONS

- Lactation: Advise not to breastfeed. (8.2)

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Recommended Dosing
- 2.2 Preparation, Handling, and Administration

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Risk of Metastatic Bladder Cancer with Delayed Cystectomy
- 5.2 Risk in Patients with Perforated Bladder
- 5.3 Risk in Patients Undergoing Transurethral Resection of the Bladder (TURB)
- 5.4 Risk in Patients with Irritable Bladder Symptoms
- 5.5 Embryo-Fetal Toxicity

6 ADVERSE REACTIONS

- 6.1 Clinical Trial Experience

7 DRUG INTERACTIONS

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.3 Females and Males of Reproductive Potential
- 8.4 Pediatric Use
- 8.5 Geriatric Use

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

15 REFERENCES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Valrubicin intravesical solution is an anthracycline topoisomerase inhibitor indicated for intravesical therapy of BCG-refractory carcinoma *in situ* (CIS) of the urinary bladder in patients for whom immediate cystectomy would be associated with unacceptable morbidity or mortality.

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dosing

For Intravesical Use Only. Do NOT administer by intravenous or intramuscular routes.

Valrubicin intravesical solution is recommended at a dose of 800 mg administered intravesically once a week for six weeks. Delay administration at least two weeks after transurethral resection and/or fulguration [see *Warnings and Precautions (5.2, 5.3)*].

2.2 Preparation, Handling, and Administration

Handle and dispose of valrubicin intravesical solution in a manner consistent with other cytotoxic drugs.¹ The use of goggles, gloves, and protective gowns is recommended during preparation and administration of the drug.

Valrubicin intravesical solution contains polyoxyl castor oil, which has been known to cause leaching of di(2-ethylhexyl) phthalate (DEHP) a hepatotoxic plasticizer, from polyvinyl chloride (PVC) bags and intravenous tubing. Valrubicin intravesical solution should be prepared and stored in glass, polypropylene, or polyolefin containers and tubing. It is recommended that non-DEHP containing administration sets, such as those that are polyethylene-lined, be used.

Valrubicin intravesical solution is a clear red solution. Visually inspect for particulate matter and discoloration prior to administration. At temperatures below 4°C (39°F), polyoxyl castor oil may begin to form a waxy precipitate. If this happens, the vial should be warmed in the hand until the solution is clear. If particulate matter is still seen, do not administer valrubicin intravesical solution.

For each instillation, slowly allow four 5 mL vials (200 mg valrubicin/5 mL vial) to warm to room temperature, but do not heat. Withdraw 20 mL of valrubicin intravesical solution from the four vials and dilute with 55 mL of 0.9% Sodium Chloride Injection, USP to provide 75 mL of a diluted valrubicin intravesical solution. Valrubicin intravesical solution diluted in 0.9% Sodium Chloride Injection, USP for administration is stable for 12 hours at temperatures up to 25°C (77°F). Since compatibility data are not available, do not mix valrubicin intravesical solution with other drugs.

Insert a urethral catheter into the patient's bladder under aseptic conditions, drain the bladder, and instill the diluted 75 mL valrubicin intravesical solution slowly via gravity flow over a period of several minutes. Withdraw the catheter and retain valrubicin intravesical solution in the bladder for two hours before voiding. At the end of two hours, all patients should void. Some patients may be unable to retain the drug for the full two hours. Instruct patients to maintain adequate hydration following valrubicin intravesical solution treatment [see *Patient Counseling Information (17)*].

3 DOSAGE FORMS AND STRENGTHS

Injection: 200 mg/5 mL sterile, clear red, solution in single-use vials for intravesical instillation upon dilution.

4 CONTRAINDICATIONS

Valrubicin intravesical solution is contraindicated in patients with:

- Perforated bladder [*see Warnings and Precautions (5.2)*]
- Known hypersensitivity to anthracyclines or polyoxyl castor oil
- Active urinary tract infection
- Small bladder capacity and unable to tolerate a 75 mL instillation

5 WARNINGS AND PRECAUTIONS

5.1 Risk of Metastatic Bladder Cancer with Delayed Cystectomy

Inform patients that valrubicin intravesical solution has been shown to induce complete response in only about 1 in 5 patients with BCG-refractory CIS, and that delaying cystectomy could lead to development of metastatic bladder cancer, which is lethal. The exact risk of developing metastatic bladder cancer from such a delay may be difficult to assess [*see Clinical Studies (14)*] but increases the longer cystectomy is delayed in the presence of persisting CIS. If there is not a complete response of CIS to treatment after 3 months or if CIS recurs, reconsider cystectomy.

5.2 Risk in Patients with Perforated Bladder

Evaluate the bladder before the intravesical instillation of drug and do not administer valrubicin intravesical solution to patients with a perforated bladder or to those in whom the integrity of the bladder mucosa has been compromised [*see Contraindications (4)*]. In case of bladder perforation, delay the administration of valrubicin intravesical solution until bladder integrity has been restored. One patient with a perforated bladder who received 800 mg of valrubicin intravesical solution intravesically developed severe leukopenia and neutropenia approximately two weeks after drug administration [*see Clinical Pharmacology (12.3)*].

5.3 Risk in Patients Undergoing Transurethral Resection of the Bladder (TURB)

To avoid systemic exposure to valrubicin intravesical solution for the patients undergoing TURB, evaluate the status of the bladder before the intravesical instillation of drug. Delay administration at least two weeks after transurethral resection and/or fulguration.

5.4 Risk in Patients with Irritable Bladder Symptoms

Use valrubicin intravesical solution with caution in patients with severe irritable bladder symptoms. Bladder spasm and spontaneous discharge of the intravesical instillate may occur; clamping of the urinary catheter is not advised.

5.5 Embryo-Fetal Toxicity

Based on findings in animal studies and its mechanism of action, valrubicin intravesical solution can cause fetal harm when administered to a pregnant woman [see *Clinical Pharmacology (12.1 and 12.3)*]. In animal reproduction studies, intravenous administration of valrubicin to pregnant rats during the period of organogenesis at a dose about 0.2 times the recommended human intravesical dose caused embryo-fetal malformations and increased resorptions. Advise females who might become pregnant of the potential risk to a fetus.

Advise females of reproductive potential to use effective contraception during treatment with valrubicin intravesical solution and for 6 months following the final dose [see *Use in Specific Populations (8.1 and 8.3)*].

6 ADVERSE REACTIONS

6.1 Clinical Trial Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The safety of valrubicin intravesical solution was assessed in 230 patients with transitional cell carcinoma of the bladder, including 205 patients who received multiple weekly doses. One hundred seventy-nine of the 205 patients received the approved dose and schedule of 800 mg weekly for multiple weeks.

Approximately 84% of patients who received intravesical valrubicin intravesical solution in clinical studies experienced local adverse reactions. The local adverse reactions associated with valrubicin intravesical solution usually occur during or shortly after instillation and resolve within 1 to 7 days after the instillate is removed from the bladder. Seven out of 143 patients (5%) who were scheduled to receive six doses of valrubicin intravesical solution failed to receive all of the planned doses because of the occurrence of local bladder symptoms.

TABLE 1 displays the frequency of the local adverse reactions at baseline and during treatment among 179 patients who received 800 mg doses of valrubicin intravesical solution in a multiple-cycle treatment regimen.

Adverse Reaction	Before Treatment	During 6-week Course of Treatment
ANY LOCAL BLADDER SYMPTOM	45%	88%
Urinary Frequency	30%	61%
Dysuria	11%	56%
Urinary Urgency	27%	57%
Bladder Spasm	3%	31%
Hematuria	11%	29%
Bladder Pain	6%	28%
Urinary Incontinence	7%	22%

Cystitis	4%	15%
Nocturia	2%	7%
Local Burning Symptoms - Procedure Related	0%	5%
Urethral Pain	0%	3%
Pelvic Pain	1%	1%
Hematuria (Gross)	0%	1%

TABLE 2 displays the adverse reactions other than local bladder symptoms that occurred in 1% or more of the 230 patients who received at least one dose of valrubicin intravesical solution in a clinical trial.

TABLE 2	
Systemic Adverse Reactions ($\geq 1\%$) Following Intravesical Administration of Valrubicin Intravesical Solution (N=230)	
Body System	
Preferred Term	
Body as a Whole	
Abdominal Pain	5%
Asthenia	4%
Headache	4%
Malaise	4%
Back Pain	3%
Chest Pain	3%
Fever	2%
Cardiovascular	
Vasodilation	2%
Digestive	
Nausea	5%
Diarrhea	3%
Vomiting	2%
Flatulence	1%
Hemic and Lymphatic	
Anemia	2%
Metabolic and Nutritional	
Hyperglycemia	1%
Peripheral Edema	1%
Musculoskeletal	
Myalgia	1%
Nervous	
Dizziness	3%
Respiratory	
Pneumonia	1%
Skin and Appendages	
Rash	3%

Urogenital	
Urinary Tract Infection	15%
Urinary Retention	4%
Hematuria (miscroscopic)	3%

Adverse reactions other than local reactions that occurred in less than 1% of the patients who received valrubicin intravesical solution intravesically in clinical trials are listed below.

Digestive System: Tenesmus

Metabolic and Nutritional: Nonprotein nitrogen increased

Skin and Appendages: Pruritus

Special Senses: Taste loss

Urogenital System: Local skin irritation, poor urine flow, and urethritis

7 DRUG INTERACTIONS

No drug interaction studies were conducted.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Based on findings in animal studies and its mechanism of action, valrubicin intravesical solution can cause fetal harm when administered to a pregnant females [see *Clinical Pharmacology (12.1 and 12.3)*]. There are no available data in pregnant females to inform the drug-associated risk. In animal reproduction studies, intravenous administration of valrubicin to pregnant rats during the period of organogenesis at a dose about 0.2 times the recommended human intravesical dose caused embryo-fetal malformations and increased resorptions [see *Data*]. Advise females who are or might become pregnant of the potential risk to a fetus.

In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically-recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

Data

Animal Data

Daily intravenous administration of valrubicin to pregnant rats during the period of organogenesis at doses ≥ 12 mg/kg (about 0.2 times the recommended human intravesical dose on a mg/m² basis) was embryo-fetal toxic and teratogenic. Administration of 12 mg/kg resulted in fetal malformations. A dose of 24 mg/kg (about 0.3 times the recommended human intravesical dose on a mg/m² basis) caused numerous, severe alterations in the skull and skeleton of the developing fetuses. This dose also caused an increase in fetal resorptions and a decrease in viable fetuses.

8.2 Lactation

Risk Summary

There are no data on the presence of valrubicin or its metabolites in human milk, the effects of valrubicin on the breast-fed infant, or its effects on milk production. Because of the potential for serious adverse reactions in breast-fed infants from valrubicin, advise a lactating female not to breastfeed during treatment with valrubicin intravesical solution and for 2 weeks after the final dose.

8.3 Females and Males of Reproductive Potential

Contraception

Females

Valrubicin intravesical solution can cause fetal harm when administered to a pregnant female. Advise females of reproductive potential to use effective contraception during treatment with valrubicin intravesical solution and for 6 months after the final dose [see *Use in Specific Populations (8.1)*].

Males

Based on genotoxicity findings, advise men with female partners of reproductive potential to use effective contraception during treatment with valrubicin intravesical solution and for 3 months following the final dose [see *Non Clinical Toxicology (13.1)*].

Infertility

Males

Studies of the effects of valrubicin intravesical solution on human male or female fertility have not been done. Based on findings in animal studies, valrubicin intravesical solution may impair fertility in males of reproductive potential [see *Nonclinical Toxicology (13.1)*].

8.4 Pediatric Use

Safety and effectiveness in pediatric patients have not been established.

8.5 Geriatric Use

Because carcinoma *in situ* of the bladder generally occurs in older individuals, 85% of the patients enrolled in the clinical studies of valrubicin intravesical solution were more than 60 years of age (49% of the patients were more than 70 years of age). In the primary efficacy studies, the mean age of the population was 69.5 years. There are no specific precautions regarding use of valrubicin intravesical solution in geriatric patients who are otherwise in good health.

10 OVERDOSAGE

There is no known antidote for overdoses of valrubicin intravesical solution. The primary anticipated complications of overdose associated with intravesical administration would be consistent with irritable bladder symptoms.

Myelosuppression is possible if valrubicin intravesical solution is inadvertently administered systemically or if significant systemic exposure occurs following intravesical administration (e.g., in patients with bladder rupture/perforation). Under

such inadvertent exposures in the peritoneal cavity, the expected toxicities include leukopenia and neutropenia, beginning within 1 week of dose administration, with nadirs by the second week, and recovery generally by the third week. If valrubicin intravesical solution is administered when bladder rupture or perforation is suspected, weekly monitoring of complete blood counts should be performed for 3 weeks.

11 DESCRIPTION

Valrubicin Intravesical Solution, USP contains valrubicin (N-trifluoroacetyl Adriamycin-14-valerate), which is a semisynthetic analog of the anthracycline doxorubicin as a cytotoxic agent. The chemical name of valrubicin is (2*S*-*cis*)-2-[1,2,3,4,6,11-hexahydro-2,5,12-trihydroxy-7-methoxy-6,11-dioxo-4-[[2,3,6-trideoxy-3-[(trifluoroacetyl)amino]- α -L-lyxo-hexopyranosyl]oxyl]-2-naphthacenyl]-2-oxoethylpentanoate. Valrubicin is an orange or orange-red powder that is highly lipophilic, soluble in methylene chloride, ethanol, methanol and acetone, and relatively insoluble in water. Its chemical formula is $C_{34}H_{36}F_3NO_{13}$ and its molecular weight is 723.65. The chemical structure is shown in FIGURE 1.




FIGURE 1. Chemical Structure of Valrubicin

Valrubicin Intravesical Solution, USP is intended for intravesical administration in the urinary bladder. It is supplied as a nonaqueous solution that should be diluted before intravesical administration. Each vial of Valrubicin Intravesical Solution, USP contains 200 mg valrubicin at a concentration of 40 mg/mL in 5 mL of 50% polyoxyl castor oil/50% v/v dehydrated alcohol, USP without preservatives or other additives. The solution is sterile and nonpyrogenic.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Valrubicin is an anthracycline that affects a variety of interrelated biological functions, most of which involve nucleic acid metabolism. In cells, it inhibits the incorporation of nucleosides into nucleic acids, causes chromosomal damage, and arrests the cell cycle in G₂. Although valrubicin does not bind strongly to DNA, valrubicin metabolites interfere

with the normal DNA breaking-resealing action of DNA topoisomerase II.

12.3 Pharmacokinetics

When 800 mg valrubicin intravesical solution was administered intravesically to patients with carcinoma *in situ*, valrubicin intravesical solution penetrated into the bladder wall. The mean total anthracycline concentration measured in bladder tissue exceeded the levels causing 90% cytotoxicity to human bladder cells cultured *in vitro*. During the two-hour dose-retention period, the metabolism of valrubicin intravesical solution to its major metabolites N-trifluoroacetyl Adriamycin and N-trifluoroacetyl Adriamycinol in bladder tissue was negligible. After retention, the drug was almost completely excreted by voiding the instillate. Mean percent recovery of valrubicin intravesical solution, N-trifluoroacetyl Adriamycin, and total anthracyclines in 14 urine samples from six patients was 98.6%, 0.4%, and 99.0% of the total administered drug, respectively. During the two-hour dose-retention period, only nanogram quantities of valrubicin intravesical solution were absorbed into the plasma. N-trifluoroacetyl Adriamycin and N-trifluoroacetyl Adriamycinol were measured in blood.

Total systemic exposure to anthracyclines during and after intravesical administration of valrubicin intravesical solution is dependent upon the condition of the bladder wall. The mean $AUC_{0-6 \text{ hours}}$ (total anthracyclines exposure) for an intravesical dose of 900 mg of valrubicin intravesical solution administered 2 weeks after transurethral resection of bladder tumors (n=6) was 78 nmol/L•hr. In patients receiving 800 mg of valrubicin intravesical solution 5 to 51 minutes after typical (n=8) and extensive (n=5) TURB tumors, the mean $AUC_{0-6 \text{ hours}}$ values for total anthracyclines were 409 and 788 nmol/L•hr, respectively. The $AUC_{0-6 \text{ hours}}$ total exposure to anthracyclines was 18,382 nmol/L•hr in one patient who experienced a perforated bladder following a transurethral resection that occurred 5 minutes before administration of an intravesical dose of 800 mg of valrubicin intravesical solution. Administration of a comparable intravenous dose of valrubicin intravesical solution (600 mg/m²; n=2) as a 24-hour infusion resulted in an $AUC_{0-6 \text{ hours}}$ for total anthracyclines of 11,975 nmol/L•hr. These results are shown in FIGURE 2.




FIGURE 2. Comparison of Mean AUC₀₋₆ hours in Valrubicin Intravesical Solution Clinical Studies (N=number of patients)

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

The carcinogenic potential of valrubicin has not been evaluated.

In vitro, valrubicin was mutagenic in the bacterial reverse mutation (Ames) assay and clastogenic in the chromosomal aberration assay in Chinese Hamster Ovary (CHO) cells.

Studies in animals evaluating the effects of valrubicin on male or female fertility have not been conducted. Based on effects on male reproductive organs in general toxicology studies in dogs with intravesical instillation, valrubicin may impair fertility in male patients. When instilled into the bladder of male dogs weekly for 6 weeks, valrubicin caused mild to moderate atrophy of the prostate with inflammation, diffuse decrease in acinar size, epithelial changes. It also caused testicular degeneration, marked germ cell depletion, spermatid giant cells and karyomegaly.

14 CLINICAL STUDIES

Valrubicin intravesical solution was administered intravesically to a total of 230 patients with transitional cell carcinoma of the bladder, including 205 patients who received multiple weekly doses ranging from 200 to 900 mg. One hundred seventy-nine of the 205 patients received the approved dose and schedule of 800 mg weekly for multiple weeks. Patients receiving valrubicin intravesical solution for refractory carcinoma *in situ* were monitored for disease recurrence or progression with cystoscopy, biopsy, and urine cytology every 3 months.

In the 90 study patients with BCG-refractory carcinoma *in situ* (CIS), 70% had received at least 2 courses of BCG and 30% had received one course of BCG and at least one additional course of treatment with another agent(s) - e.g., mitomycin, thiotepa, or interferon. Valrubicin intravesical solution was administered beginning at least two weeks after transurethral resection and/or fulguration.

After intravesical administration of valrubicin intravesical solution, 16 patients (18%) had a complete response documented by bladder biopsies and cytology at 6 months following initiation of therapy. Median duration of response from start of treatment varied according to the method of analysis (13.5 months if measured to last bladder biopsy without tumor and 21 months if measured until time of documented recurrence). A retrospective analysis in the 16 patients with complete response to valrubicin intravesical solution demonstrated that time to recurrence of their disease after treatment with valrubicin intravesical solution was longer than time to recurrence after previous courses of intravesical therapy.

Of the 90 patients with BCG-refractory CIS, 11% (10 patients) developed metastatic or deeply-invasive bladder cancer during follow-up; four of these patients, none who underwent cystectomy, died with metastatic bladder cancer and six were found to have developed stage progression to deeply-invasive disease (T3), with lymph node involvement in one patient, at the time of cystectomy. It is uncertain to what extent the development of advanced bladder cancer in these patients was due to the delay in cystectomy required to receive treatment with valrubicin intravesical solution (3 months was the time of follow-up to determine response), as cystectomy was often delayed or was never performed despite failure of treatment with valrubicin intravesical solution. In the 10 patients documented to have invasive bladder cancer or metastatic disease, the delay between the time of treatment failure (when cystectomy should have been performed) and cystectomy or documentation of advanced bladder cancer was a median of 17.5 months.

15 REFERENCES

1. OSHA Hazardous Drugs. OSHA.
<http://www.osha.gov/SLTC/hazardousdrugs/index.html>.

16 HOW SUPPLIED/STORAGE AND HANDLING

Valrubicin Intravesical Solution, USP is a sterile, clear red solution in polyoxyl castor oil/dehydrated alcohol, USP, containing 40 mg valrubicin per mL. Valrubicin Intravesical

Solution, USP is available in single-use, clear glass vials, individually packaged in the following sizes:

NDC 24201-101-04 Carton of four 200 mg/5 mL single-use vials

Store vials under refrigeration at 2°-8°C (36°-46°F) in the carton. DO NOT FREEZE.

The container closure is not made with natural rubber latex.

17 PATIENT COUNSELING INFORMATION

Risk of Metastatic Bladder Cancer with Delayed Cystectomy

- Inform patients that valrubicin intravesical solution has been shown to induce complete responses in only about 1 in 5 patients, and that delaying cystectomy could lead to development of metastatic bladder cancer, which is lethal. Discuss the relative risk of cystectomy versus the risk of metastatic bladder cancer [*see Clinical Trials (14)*] and that the risk increases the longer cystectomy is delayed in the presence of persisting CIS.

Local Adverse Reactions Before and During Treatment

- Inform patients that the major acute toxicities from valrubicin intravesical solution are related to irritable bladder symptoms that may occur during instillation and retention of valrubicin intravesical solution and for a limited period following voiding [*see Adverse Reactions (6.1)*].
- Inform patients that for the first 24 hours following administration, red-tinged urine is typical.
- Advise patients to report prolonged irritable bladder symptoms or prolonged passage of red-colored urine immediately to their physician.
- Instruct patients to maintain adequate hydration following valrubicin intravesical solution treatment.

Embryo-Fetal Toxicity

- Advise females of reproductive potential of the potential risk to a fetus and to use effective contraception during treatment with valrubicin intravesical solution and for 6 months after the last dose. Advise females to inform their healthcare provider of a known or suspected pregnancy [*see Warnings and Precautions (5.5), Use in Specific Populations (8.1 and 8.3)*].
- Advise male patients with female partners of reproductive potential to use effective contraception during treatment with valrubicin intravesical solution and for 3 months after the last dose [*see Use in Specific Populations (8.3)*].

Lactation

- Advise females not to breastfeed during treatment with valrubicin intravesical solution and for 2 weeks after the last dose [*see Use in Specific Populations (8.2)*].

Healthcare professionals can email Hikma at us.hikma@primevigilance.com or call 1-877-845-0689 for information on this product.

Manufactured by:
The University of Iowa Pharmaceuticals
Iowa City, Iowa 52242

Manufactured for:
Hikma
Berkeley Heights, NJ 07922

Rev. June 2022

PACKAGE/LABEL PRINCIPAL DISPLAY PANEL

Caution: Cytotoxic Agent

NDC 24201-101-01 Rx only

**Valrubicin
Intravesical
Solution, USP**
200 mg/5 mL
(40 mg/mL)

For Intravesical Use Only
NOT for Intramuscular or Intravenous Use
5 mL Single Dose Vial.
Discard Unused Portion.

WARNING: Do not administer without diluting. See enclosed package insert for proper handling.
Store refrigerated 2° and 8°C (36° and 46°F).
DO NOT FREEZE
Preservative-free.
The container closure is not made with natural rubber latex.

Mfd for R06/2022
Hikma
Berkeley Heights,
NJ 07922 668115

(01)00324201101015

Package/Label Display Panel

**Valrubicin
Intravesical Solution, USP**
200 mg/5 mL (40 mg/mL)

For Intravesical Use Only. NOT for Intramuscular or Intravenous Use.
WARNING: Do not administer without diluting. See enclosed package insert for proper handling.

Rx only

4 x 5 mL Single Dose Vials.
Discard Unused Portion

Caution: Cytotoxic Agent

NDC 24201-101-04

3 24201 10104 6

**Valrubicin
Intravesical Solution, USP**
200 mg/5 mL (40 mg/mL)

For Intravesical Use Only, NOT for Intramuscular or Intravenous Use.
WARNING: Do not administer without diluting. See enclosed package insert for proper handling.
Usual Dosage: See enclosed package insert.
Description: Each mL contains 40 mg Valrubicin, 50% polyoxyl castor oil, and 50% w/v dehydrated alcohol, USP.
Store in refrigerator, 2°-8°C (36°-46°F).
DO NOT FREEZE
Sterile, Non Pyrogenic, Preservative-free.
The container closure is not made with natural rubber latex.

582219-01

4 x 5 mL Single Dose Vial, Discard Unused Portion
Intravesical Solution, USP
200 mg/5 mL (40 mg/mL)
Valrubicin
NDC 24201-101-04
Caution: Cytotoxic Agent

R06/2022
Manufactured for:
Hikma
Berkeley Heights,
NJ 07922

hikma.

VALRUBICIN INTRAVESICAL SOLUTION

valrubicin intravesical solution solution, concentrate

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:24201-101
Route of Administration	INTRAVESICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
VALRUBICIN (UNII: 2C6NUM6878) (VALRUBICIN - UNII:2C6NUM6878)	VALRUBICIN	40 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	0.5 mL in 1 mL
POLYOXYL 35 CASTOR OIL (UNII: 6D4M1DAL6O)	0.5 mL in 1 mL
NITROGEN (UNII: N762921K75)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:24201-101-04	4 in 1 CARTON	04/23/2019	
1		5 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206430	04/23/2019	

Labeler - Hikma Pharmaceuticals USA Inc. (118707839)

Registrant - Hikma Pharmaceuticals USA Inc. (118707839)

Revised: 6/2022

Hikma Pharmaceuticals USA Inc.