

**TUSSIN CF MULTI SYMPTOM COLD ADULT- dextromethorphan hbr,  
guaifenesin, phenylephrine liquid  
P & L Development, LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**Drug Facts**

**Active ingredients (in each 10 mL)**

Dextromethorphan HBr 20 mg

Guaifenesin 200 mg

Phenylephrine HCl 10 mg

**Purposes**

Cough suppressant

Expectorant

Nasal decongestant

**Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
  - nasal congestion
  - cough due to minor throat and bronchial irritation

**Warnings**

**Do not use**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- diabetes
- high blood pressure
- thyroid disease
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

- trouble urinating due to enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**

taking any other oral nasal decongestant or stimulant.

**When using this product,**

**do not use more than directed.**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL=milliliter
- this adult product is not intended for use in children under 12 years of age
- adults and children 12 years and over: 10 mL every 4 hours
- children under 12 years: do not use

**Other information**

- store between 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients**

anhydrous citric acid, FD&C red #40, flavor, glycerin, lactic acid, menthol, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose

**Questions or comments?**

Call **1-877-753-3935 Monday-Friday 9AM-5PM EST**

**Principal Display Panel**

Compare to active ingredients in Robitussin® Peak Cold Multi-Symptom Cold CF\*

**Adult**

**Tussin CF**

Multi-Symptom Cold

Dextromethorphan HBr

Guaifenesin

Phenylephrine HCl

Relieves:

- Cough
- Mucus
- Nasal Congestion

Non-drowsy

for ages 12 years and over

Alcohol free

Dosing Cup Included

FL OZ (mL)

\*This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributors of Robitussin® Peak Cold Multi-Symptom Cold CF.

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

Manufactured by:

PL Developments

11865 S. Alameda St

Lynwood, CA 90262

**Package Label**

**Drug Facts (continued)**

**Directions**

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age
- adults and children 12 years and over: 10 mL every 4 hours
- children under 12 years: do not use

**Other information**

- store between 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients** citric acid, FD&C red #40, flavor, glycerin, lactic acid, menthol, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose

**Questions or comments?**  
Call 1-877-753-3835 Monday-Friday 9AM-5PM EST

\*This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Robitussin® Peak Cold Multi-Symptom Cold CF.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

**PARENTS:**  
Learn about your medicine online  
[www.StopMedicineInCase.org](http://www.StopMedicineInCase.org)



Dosing Cup Included



Manufactured by:  
**PL Developments**  
11865 S. Alameda St  
Lynwood, CA 90262



Compare to active ingredients in Robitussin® Peak Cold Multi-Symptom Cold CF\*  
NDC 49580-0382-4



relieves:

- cough
- mucus
- nasal congestion

for ages 12 years and over  
non-drowsy  
alcohol-free

4 fl oz (118 mL)



Compare to active ingredients in Robitussin® Peak Cold Multi-Symptom Cold CF\*  
NDC 49580-0382-4



relieves:

- cough
- mucus
- nasal congestion

for ages 12 years and over  
non-drowsy  
alcohol-free

4 fl oz (118 mL)

**Drug Facts**

Active ingredients (in each 10 mL)	Purposes
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant

**Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold
- nasal congestion
- cough due to minor throat and bronchial irritation

**Warnings**

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

**Ask a doctor or pharmacist before use if you are taking any other oral nasal decongestant or stimulant.**

**When using this product, do not use more than directed.**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.



## ReadyinCase Adult Tussin CF Multi-Symptom Cold

### TUSSIN CF MULTI SYMPTOM COLD ADULT

dextromethorphan hbr, guaifenesin, phenylephrine liquid

#### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:49580-0382
<b>Route of Administration</b>	ORAL		

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 10 mL
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 10 mL

<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)		PHENYLEPHRINE HYDROCHLORIDE	10 mg in 10 mL	
<b>Inactive Ingredients</b>				
<b>Ingredient Name</b>			<b>Strength</b>	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)				
GLYCERIN (UNII: PDC6A3C0OX)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
WATER (UNII: 059QF0KO0R)				
SODIUM BENZOATE (UNII: OJ245FE5EU)				
FD&C RED NO. 40 (UNII: WZB9127XOA)				
LACTIC ACID (UNII: 33X04XA5AT)				
MENTHOL (UNII: L7T10EIP3A)				
SORBITOL (UNII: 506T60A25R)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:49580-0382-4	1 in 1 BOX	04/30/2015	04/30/2025
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>	
OTC monograph final	part341	04/30/2015	04/30/2025	

**Labeler** - P & L Development, LLC (101896231)

Revised: 6/2023

P & L Development, LLC