

**COLD AND FLU SEVERE DAYTIME NIGHTTIME- acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride MEIJER**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**MEI-1186-2019-0521**

**Cold + Flu Severe Day**

**Drug Facts**

<b>Active ingredients (in each caplet)</b>	<b>Purpose</b>
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

**Uses**

- for the temporary relief of the following cold/flu symptoms:
  - minor aches and pain
  - headache
  - sore throat
  - nasal congestion
  - cough
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily reduces fever

**Warnings**

**Liver warning**

This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

**When using this product****do not exceed recommended dosage****Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than directed (see overdose warning)

adults and children  
12 years and over

- take 2 caplets every 4 hours
- swallow whole – do not crush, chew, or dissolve
- do not take more than 10 caplets in 24

	hours
children under 12 years	<ul style="list-style-type: none"> <li>■ ask a doctor</li> </ul>

## Other information

- store between 20-25°C (68-77°F) in a dry place
- retain carton for complete product information and warnings

## Inactive ingredients

acesulfame potassium, colloidal silicon dioxide, croscarmellose sodium, crospovidone, D&C yellow #10 aluminum lake, FD&C blue #2 aluminum lake, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, stearic acid, talc, titanium dioxide

## Cold + Flu Severe Night

### Drug Facts

<b>Active ingredients (in each caplet)</b>	<b>Purpose</b>
Acetaminophen 325 mg	Pain reliever/fever reducer
Chlorpheniramine Maleate 2 mg	Antihistamine
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

## Uses

- for the temporary relief of the following cold/flu symptoms:
  - minor aches and pains
  - headache
  - sore throat
  - nasal congestion
  - cough
  - sinus congestion and pressure
  - sneezing and runny nose
- helps clear nasal passages
- relieves cough to help you sleep
- temporarily reduces fever

## Warnings

### Liver warning

This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

### Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

### **Sore throat warning**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma

### **Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

### **When using this product**

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

### **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.**

### **Overdose warning**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- do not use more than directed (see overdose warning)

adults and children 12 years and over	<ul style="list-style-type: none"><li>■ take 2 caplets every 4 hours</li><li>■ swallow whole – do not crush, chew, or dissolve</li><li>■ do not take more than 10 caplets in 24 hours</li></ul>
children under 12 years	<ul style="list-style-type: none"><li>■ ask a doctor</li></ul>

### **Other information**

- store between 20-25°C (68-77°F) in a dry place
- retain carton for complete product information and warnings

### **Inactive ingredients**

acesulfame potassium, colloidal silicon dioxide, croscarmellose sodium, FD&C blue #1 aluminum lake, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, stearic acid, talc, titanium dioxide

### **PRINCIPAL DISPLAY PANEL**

#### **COMBO PACK**

NDC 41250-753-01

Compare to Tylenol® Cold & Flu Severe Day + Night active ingredients†

meijer®

daytime

cold & flu severe

Acetaminophen |Dextromethorphan HBr

Phenylephrine HCl | Guaifenesin

Pain Reliever / Fever Reducer; Cough Suppressant;

Nasal Decongestant; Expectorant

Day Caplet Relieves:

Head & Body Aches

Fever & Sore Throat

Cough, Nasal Congestion

Mucus & Chest Congestion

16 CAPLETS

COOL TASTE

instant cooling sensation

nighttime

cold & flu severe

Acetaminophen | Chlorpheniramine Maleate

Dextromethorphan HBr | Phenylephrine HCl

Pain Reliever/Fever Reducer; Antihistamine;

Cough Suppressant; Nasal Decongestant

Night Caplet Relieves:

Head & Body Aches

Fever & sore Throat


Cough, Nasal Congestion

Runny Nose

8 CAPLETS

COOL TASTE

instant cooling sensation



## COLD AND FLU SEVERE DAYTIME NIGHTTIME

acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide, guaifenesin, and phenylephrine

hydrochloride kit

## Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:41250-753

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-753-01	1 in 1 CARTON	05/21/2019	

## Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	2 BLISTER PACK	16
Part 2	1 BLISTER PACK	8

## Part 1 of 2

### ACETAMINOPHEN, DEXTROMETHORPHAN HYDROBROMIDE, GUAIFENESIN, AND PHENYLEPHRINE HYDROCHLORIDE

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, coated

## Product Information

Route of Administration

ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINO PHEN (UNII: 362O9ITL9D) (ACETAMINO PHEN - UNII:362O9ITL9D)	ACETAMINO PHEN	325 mg
DEXTRO METHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTRO METHORPHAN - UNII:7355X3ROTS)	DEXTRO METHORPHAN HYDROBROMIDE	10 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDRO CHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

## Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
CROSPOVIDONE (UNII: 2S7830E561)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	

MAGNESIUM STEARATE (UNII: 70097M6I30)

MALTO DEXTRIN (UNII: 7CVR7L4A2D)

CELLULOSE, MICRO CRYSTALLINE (UNII: OP1R32D61U)

POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)

POLYVINYL ALCOHOL (UNII: 532B59J990)

POVIDONE (UNII: FZ989GH94E)

STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)

STEARIC ACID (UNII: 4ELV7Z65AP)

TALC (UNII: 7SEV7J4R1U)

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

### Product Characteristics

Color	yellow	Score	no score
Shape	OVAL	Size	19mm
Flavor	MINT	Imprint Code	AAA;1136
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		8 in 1 BLISTER PACK; Type 1: Convenience Kit of Co-Package		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341		

### Part 2 of 2

## ACETAMINOPHEN, CHLORPHENIRAMINE MALEATE, DEXTROMETHORPHAN HYDROBROMIDE, AND PHENYLEPHRINE HYDROCHLORIDE

acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide, and phenylephrine hydrochloride tablet, coated

### Product Information

Route of Administration	ORAL
-------------------------	------

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE -	CHLORPHENIRAMINE MALEATE	2 mg

UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3R0TS)	DEXTROMETHORPHAN HYDROBROMIDE	10
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>ACESULFAME POTASSIUM</b> (UNII: 23OV73Q5G9)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>ALUMINUM OXIDE</b> (UNII: LMI26O6933)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>STARCH, PREGELATINIZED CORN</b> (UNII: O8232NY3SJ)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	blue	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	17mm
<b>Flavor</b>	MINT	<b>Imprint Code</b>	AAA;1139
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		8 in 1 BLISTER PACK; Type 1: Convenience Kit of Co-Package		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	05/21/2019	

Revised: 5/2019

MEIJER