

**TOXIBAN WITH SORBITOL- activated charcoal suspension**  
**LLOYD, Inc. of Iowa**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.*

-----  
**ToxiBan® With Sorbitol**

**INDICATION:**

For use as a decontamination agent via adsorption of organic chemical toxicants in domestic animals.

In all cases of suspected poisonings, consult your veterinarian for complete diagnosis and specific treatment.

**DOSAGE AND ADMINISTRATION:**

Give orally 10 to 20 ml per kg body weight (5 to 10 ml per lb) by causing the animal to consume the calculated dose. The product may be given as is or mixed with a small amount of cold water. Consult a veterinarian should delivery by stomach tube be needed.

**CAUTION:** Do not use in animals with ileus or intestinal obstruction. ToxiBan Suspension with Sorbitol should not be given in conjunction with oral medications as the charcoal may adsorb the therapeutic agent. Maintain patient hydration. Monitor for 4 hours after use for signs of hypernatremia (ataxia, tremors, seizures).

**NOTE:** Osmotic cathartics may be associated with electrolyte disturbances (hypernatremia) due to loss of excess free body water in the gut. Dehydration and hypotension have been reported with excess catharsis. Cathartics should only be used with the initial charcoal dose. If charcoal is still required, use ToxiBan Suspension or Granules at the lower dosage every 6-8 hours for 24 - 48 hours. **Tests for ethylene glycol must be conducted before ToxiBan Suspension is given to prevent false positive reactions.**

**Adsorptive Power:..... Per ml**

Each ml of ToxiBan Suspension with Sorbitol can adsorb at least 55 mg strychnine sulfate.

**SHAKE WELL BEFORE USE**

**KEEP OUT OF REACH OF CHILDREN**

Store at room temperature. Avoid excess humidity.

**Discard within 48 hours after opening.**

**LLOYD, Inc.**  
Shenandoah, IA 51601 USA

List No. 1155

Rev. 0623

# PRINCIPAL DISPLAY PANEL - 240 mL Bottle Label

NDC 11789-115-50

ToxiBan®

Charcoal-Kaolin  
Suspension  
With Sorbitol


## INDICATION:

For use as a decontamination agent via adsorption of organic chemical toxicants in domestic animals.

In all cases of suspected poisonings, consult your veterinarian for complete diagnosis and specific treatment.

LLOYD®

Net Volume: 240 mL (8.1 fl oz)

<p><b>DOSAGE AND ADMINISTRATION:</b> Give orally 10 to 20 mL per kg body weight (5 to 10 mL per lb) by causing the animal to consume the calculated dose. The product may be given as is or mixed with a small amount of cold water. Consult a veterinarian should delivery by stomach tube be needed.</p> <p><b>CAUTION:</b> Do not use in animals with ileus or intestinal obstruction. ToxiBan Suspension with Sorbitol should not be given in conjunction with oral medications as the charcoal may adsorb the therapeutic agent. Maintain patient hydration. Monitor for 4 hours after use for signs of hypernatremia (ataxia, tremors, seizures).</p> <p><b>NOTE:</b> Osmotic cathartics may be associated with electrolyte disturbances (hypernatremia) due to loss of excess free body water in the gut. Dehydration and hypotension have been reported with excess catharsis. Cathartics should only be used with the initial charcoal dose. If charcoal is still required, use ToxiBan Suspension or Granules at the lower dosage every 6-8 hours for 24 - 48 hours. Tests for ethylene glycol must be conducted before ToxiBan Suspension is given to prevent false positive reactions.</p>	<p>NDC 11789-115-50</p> <p><b>ToxiBan®</b> <b>Charcoal-Kaolin</b> <b>Suspension</b> <b>With Sorbitol</b></p> <p><b>INDICATION:</b> For use as a decontamination agent via adsorption of organic chemical toxicants in domestic animals.</p> <p><b>In all cases of suspected poisonings, consult your veterinarian for complete diagnosis and specific treatment.</b></p> <p><b>LLOYD</b></p> <p>Net Volume: 240 mL (8.1 fl oz)</p>	<p>Adsorptive Power: _____ Per mL Each mL of ToxiBan Suspension with Sorbitol can adsorb at least 55 mg strychnine sulfate.</p> <p><b>SHAKE WELL BEFORE USE</b> <b>KEEP OUT OF REACH OF CHILDREN</b></p> <p>Store at room temperature. Avoid excess humidity. <b>Discard within 48 hours after opening.</b></p> <p><b>LLOYD, Inc.</b> Shenandoah, IA 51801 USA</p> <p>List No. 1155</p> <p>Rev. 0623</p> 
--	--	--

## TOXIBAN WITH SORBITOL

activated charcoal suspension

### Product Information

<b>Product Type</b>	OTC ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:11789-115
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

<b>ACTIVATED CHARCOAL</b> (UNII: 2P3VWU3H10) (ACTIVATED CHARCOAL - UNII:2P3VWU3H10)	ACTIVATED CHARCOAL	55 [arb'U] in 1 mL
---	--------------------	--------------------

Inactive Ingredients	
Ingredient Name	Strength
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>KAOLIN</b> (UNII: 24H4NWX5CO)	
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)	

Product Characteristics			
<b>Color</b>	BLACK (BLACK LIQUID)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11789-115-50	240 mL in 1 BOTTLE		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
UNAPPROVED DRUG OTHER		07/07/2023	

**Labeler -** LLOYD, Inc. of Iowa (962286535)

**Registrant -** LLOYD, Inc. of Iowa (007281942)

Establishment			
Name	Address	ID/FEI	Business Operations
LLOYD, Inc. of Iowa		962286535	LABEL, PACK, API MANUFACTURE, MANUFACTURE

Establishment			
Name	Address	ID/FEI	Business Operations
LLOYD, Inc. of Iowa		007281942	ANALYSIS