

**ZOSYN IN GALAXY CONTAINERS- tazobactam sodium and piperacillin sodium injection, solution**  
**Wyeth Pharmaceuticals LLC, a subsidiary of Pfizer Inc.**

**HIGHLIGHTS OF PRESCRIBING INFORMATION**

These highlights do not include all the information needed to use ZOSYN safely and effectively. See full prescribing information for ZOSYN.

**ZOSYN® (piperacillin and tazobactam) injection, for intravenous use**  
**Initial U.S. Approval: 1993**

**RECENT MAJOR CHANGES**

Warnings and Precautions, Hemophagocytic Lymphohistiocytosis (5.3)

4/2022

**INDICATIONS AND USAGE**

ZOSYN is a combination of piperacillin, a penicillin-class antibacterial and tazobactam, a beta-lactamase inhibitor, indicated for the treatment of:

- Intra-abdominal infections in adult and pediatric patients 2 months of age and older (1.1)
- Nosocomial pneumonia in adult and pediatric patients 2 months of age and older (1.2)
- Skin and skin structure infections in adults (1.3)
- Female pelvic infections in adults (1.4)
- Community-acquired pneumonia in adults (1.5)

To reduce the development of drug-resistant bacteria and maintain the effectiveness of ZOSYN and other antibacterial drugs, ZOSYN should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria. (1.6)

**DOSAGE AND ADMINISTRATION**

- If a dose of ZOSYN is required that does not equal 2.25 g, 3.375 g, or 4.5 g, ZOSYN injection in GALAXY Containers is not recommended for use and an alternative formulation of ZOSYN should be considered. (2.1)
- *Adult Patients With Indications Other Than Nosocomial Pneumonia*; The usual daily dosage of ZOSYN for adults is 3.375 g every six hours totaling 13.5 g (12.0 g piperacillin and 1.5 g tazobactam). (2.2)
- *Adult Patients with Nosocomial Pneumonia*: Initial presumptive treatment of patients with nosocomial pneumonia should start with ZOSYN at a dosage of 4.5 g every six hours plus an aminoglycoside, totaling 18.0 g (16.0 g piperacillin and 2.0 g tazobactam). (2.3)
- *Adult Patients with Renal Impairment*: Dosage in patients with renal impairment (creatinine clearance  $\leq$ 40 mL/min) and dialysis patients should be reduced, based on the degree of renal impairment. (2.4)
- *Pediatric Patients by Indication and Age*: See Table below (2.5)

**Recommended Dosage of ZOSYN for Pediatric Patients 2 months of Age and Older, Weighing up to 40 Kg and With Normal Renal Function**

<b>Age</b>	<b>Appendicitis and /or Peritonitis</b>	<b>Nosocomial Pneumonia</b>
<b>2 months to 9 months</b>	90 mg/kg (80 mg piperacillin and 10 mg tazobactam) <u>every 8 (eight) hours</u>	90 mg/kg (80 mg piperacillin and 10 mg tazobactam) <u>every 6 (six) hours</u>
<b>Older than 9 months</b>	112.5 mg/kg (100 mg piperacillin and 12.5 mg tazobactam) <u>every 8 (eight) hours</u>	112.5 mg/kg (100 mg piperacillin and 12.5 mg tazobactam) <u>every 6 (six) hours</u>

- Administer ZOSYN by intravenous infusion over 30 minutes to both adult and pediatric patients (2.2, 2.3, 2.4, 2.5).
- ZOSYN and aminoglycosides should be reconstituted, diluted, and administered separately. Co-administration via Y-site can be done under certain conditions. (2.7)

See the full prescribing information for the preparation and administration instructions for ZOSYN Injection in GALAXY Containers.

### DOSAGE FORMS AND STRENGTHS

ZOSYN® Injection: 2.25 g in 50 mL, 3.375 g in 50 mL, and 4.5 g in 100 mL frozen solution in single-dose GALAXY Containers. (3, 16)

### CONTRAINDICATIONS

Patients with a history of allergic reactions to any of the penicillins, cephalosporins, or beta-lactamase inhibitors. (4)

### WARNINGS AND PRECAUTIONS

- Serious hypersensitivity reactions (anaphylactic/anaphylactoid) reactions have been reported in patients receiving ZOSYN. Discontinue ZOSYN if a reaction occurs. (5.1)
- ZOSYN may cause severe cutaneous adverse reactions, such as Stevens-Johnson syndrome, toxic epidermal necrolysis, drug reaction with eosinophilia and systemic symptoms, and acute generalized exanthematous pustulosis. Discontinue ZOSYN for progressive rashes. (5.2)
- Hemophagocytic lymphohistiocytosis (HLH) has been reported with the use of ZOSYN. If HLH is suspected, discontinue ZOSYN immediately. (5.3)
- Hematological effects (including bleeding, leukopenia and neutropenia) have occurred. Monitor hematologic tests during prolonged therapy. (5.4)
- As with other penicillins, ZOSYN may cause neuromuscular excitability or seizures. Patients receiving higher doses, especially in the presence of renal impairment may be at greater risk. Closely monitor patients with renal impairment or seizure disorders for signs and symptoms of neuromuscular excitability or seizures. (5.5)
- Nephrotoxicity in critically ill patients has been observed; the use of ZOSYN was found to be an independent risk factor for renal failure and was associated with delayed recovery of renal function as compared to other beta-lactam antibacterial drugs in a randomized, multicenter, controlled trial in critically ill patients. Based on this study, alternative treatment options should be considered in the critically ill population. If alternative treatment options are inadequate or unavailable, monitor renal function during treatment with ZOSYN. (5.6)
- *Clostridioides difficile*-associated diarrhea: evaluate patients if diarrhea occurs. (5.8)

### ADVERSE REACTIONS

The most common adverse reactions (incidence >5%) are diarrhea, constipation, nausea, headache, and insomnia. (6.1)

**To report SUSPECTED ADVERSE REACTIONS, contact Pfizer Inc. at 1-800-438-1985 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

### DRUG INTERACTIONS

- ZOSYN administration can significantly reduce tobramycin concentrations in hemodialysis patients. Monitor tobramycin concentrations in these patients. (7.1)
- Probenecid prolongs the half-lives of piperacillin and tazobactam and should not be co-administered with ZOSYN unless the benefit outweighs the risk. (7.2)
- Co-administration of ZOSYN with vancomycin may increase the incidence of acute kidney injury. Monitor kidney function in patients receiving ZOSYN and vancomycin. (7.3)
- Monitor coagulation parameters in patients receiving ZOSYN and heparin or oral anticoagulants. (7.4)
- ZOSYN may prolong the neuromuscular blockade of vecuronium and other non-depolarizing neuromuscular blockers. Monitor for adverse reactions related to neuromuscular blockade. (7.5)

### USE IN SPECIFIC POPULATIONS

Dosage in patients with renal impairment (creatinine clearance  $\leq$ 40 mL/min) should be reduced based on the degree of renal impairment. (2.4, 8.6)

**See 17 for PATIENT COUNSELING INFORMATION.**

**Revised: 12/2022**

- 1.1 Intra-abdominal Infections
- 1.2 Nosocomial Pneumonia
- 1.3 Skin and Skin Structure Infections
- 1.4 Female Pelvic Infections
- 1.5 Community-acquired Pneumonia
- 1.6 Usage

## **2 DOSAGE AND ADMINISTRATION**

- 2.1 Important Administration Instructions
- 2.2 Dosage in Adult Patients With Indications Other Than Nosocomial Pneumonia
- 2.3 Dosage in Adult Patients With Nosocomial Pneumonia
- 2.4 Dosage in Adult Patients With Renal Impairment
- 2.5 Dosage in Pediatric Patients With Appendicitis/Peritonitis or Nosocomial Pneumonia
- 2.6 Directions for Use of ZOSYN Injection
- 2.7 Compatibility With Aminoglycosides

## **3 DOSAGE FORMS AND STRENGTHS**

## **4 CONTRAINDICATIONS**

## **5 WARNINGS AND PRECAUTIONS**

- 5.1 Hypersensitivity Adverse Reactions
- 5.2 Severe Cutaneous Adverse Reactions
- 5.3 Hemophagocytic Lymphohistiocytosis
- 5.4 Hematologic Adverse Reactions
- 5.5 Central Nervous System Adverse Reactions
- 5.6 Nephrotoxicity in Critically Ill Patients
- 5.7 Electrolyte Effects
- 5.8 *Clostridioides difficile*-Associated Diarrhea
- 5.9 Development of Drug-Resistant Bacteria

## **6 ADVERSE REACTIONS**

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience
- 6.3 Additional Experience with Piperacillin

## **7 DRUG INTERACTIONS**

- 7.1 Aminoglycosides
- 7.2 Probenecid
- 7.3 Vancomycin
- 7.4 Anticoagulants
- 7.5 Vecuronium
- 7.6 Methotrexate
- 7.7 Effects on Laboratory Tests

## **8 USE IN SPECIFIC POPULATIONS**

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment
- 8.7 Hepatic Impairment
- 8.8 Patients with Cystic Fibrosis

## **10 OVERDOSAGE**

## **11 DESCRIPTION**

## **12 CLINICAL PHARMACOLOGY**

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

12.4 Microbiology

### **13 NONCLINICAL TOXICOLOGY**

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

### **15 REFERENCES**

### **16 HOW SUPPLIED/STORAGE AND HANDLING**

### **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

#### **1.1 Intra-abdominal Infections**

ZOSYN is indicated in adults and pediatric patients (2 months of age and older) for the treatment of appendicitis (complicated by rupture or abscess) and peritonitis caused by beta-lactamase producing isolates of *Escherichia coli* or the following members of the *Bacteroides fragilis* group: *B. fragilis*, *B. ovatus*, *B. thetaiotaomicron*, or *B. vulgatus*.

#### **1.2 Nosocomial Pneumonia**

ZOSYN is indicated in adults and pediatric patients (2 months of age and older) for the treatment of nosocomial pneumonia (moderate to severe) caused by beta-lactamase producing isolates of *Staphylococcus aureus* and by piperacillin and tazobactam-susceptible *Acinetobacter baumannii*, *Haemophilus influenzae*, *Klebsiella pneumoniae*, and *Pseudomonas aeruginosa* (Nosocomial pneumonia caused by *P. aeruginosa* should be treated in combination with an aminoglycoside) [see *Dosage and Administration (2)*].

#### **1.3 Skin and Skin Structure Infections**

ZOSYN is indicated in adults for the treatment of uncomplicated and complicated skin and skin structure infections, including cellulitis, cutaneous abscesses and ischemic/diabetic foot infections caused by beta-lactamase producing isolates of *Staphylococcus aureus*.

#### **1.4 Female Pelvic Infections**

ZOSYN is indicated in adults for the treatment of postpartum endometritis or pelvic inflammatory disease caused by beta-lactamase producing isolates of *Escherichia coli*.

#### **1.5 Community-acquired Pneumonia**

ZOSYN is indicated in adults for the treatment of community-acquired pneumonia (moderate severity only) caused by beta-lactamase producing isolates of *Haemophilus*

*influenzae*.

## 1.6 Usage

To reduce the development of drug-resistant bacteria and maintain the effectiveness of ZOSYN and other antibacterial drugs, ZOSYN should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

## 2 DOSAGE AND ADMINISTRATION

### 2.1 Important Administration Instructions

If a dose of ZOSYN is required that does not equal 2.25 g, 3.375 g, or 4.5 g, ZOSYN injection in GALAXY Containers is not recommended for use and an alternative formulation of ZOSYN should be considered.

### 2.2 Dosage in Adult Patients With Indications Other Than Nosocomial Pneumonia

The usual total daily dosage of ZOSYN for adult patients with indications other than nosocomial pneumonia is 3.375 g every six hours [totaling 13.5 g (12.0 g piperacillin and 1.5 g tazobactam)], to be administered by intravenous infusion over 30 minutes. The usual duration of ZOSYN treatment is from 7 to 10 days.

### 2.3 Dosage in Adult Patients With Nosocomial Pneumonia

Initial presumptive treatment of adult patients with nosocomial pneumonia should start with ZOSYN at a dosage of 4.5 g every six hours plus an aminoglycoside, [totaling 18.0 g (16.0 g piperacillin and 2.0 g tazobactam)], administered by intravenous infusion over 30 minutes. The recommended duration of ZOSYN treatment for nosocomial pneumonia is 7 to 14 days. Treatment with the aminoglycoside should be continued in patients from whom *P. aeruginosa* is isolated.

### 2.4 Dosage in Adult Patients With Renal Impairment

In adult patients with renal impairment (creatinine clearance  $\leq$  40 mL/min) and dialysis patients (hemodialysis and CAPD), the intravenous dose of ZOSYN should be reduced based on the degree of renal impairment. The recommended daily dosage of ZOSYN for patients with renal impairment administered by intravenous infusion over 30 minutes is described in Table 1.

**Table 1: Recommended Dosage of ZOSYN in Patients with Normal Renal Function and Renal Impairment (As total grams piperacillin and tazobactam)\***

<b>Creatinine clearance,</b>	<b>All Indications (except nosocomial</b>	<b>Nosocomial</b>
------------------------------	---	-------------------

<b>mL/min</b>	<b>pneumonia)</b>	<b>Pneumonia</b>
Greater than 40 mL/min	3.375 every 6 hours	4.5 every 6 hours
20 to 40 mL/min <sup>†</sup>	2.25 every 6 hours	3.375 every 6 hours
Less than 20 mL/min <sup>†</sup>	2.25 every 8 hours	2.25 every 6 hours
Hemodialysis <sup>‡</sup>	2.25 every 12 hours	2.25 every 8 hours
CAPD	2.25 every 12 hours	2.25 every 8 hours

\* Administer **ZOSYN** by intravenous infusion over 30 minutes.

† Creatinine clearance for patients not receiving hemodialysis

‡ 0.75 g (0.67 g piperacillin and 0.08 g tazobactam) should be administered following each hemodialysis session on hemodialysis days

For patients on hemodialysis, the maximum dose is 2.25 g every twelve hours for all indications other than nosocomial pneumonia and 2.25 g every eight hours for nosocomial pneumonia. Since hemodialysis removes 30% to 40% of the administered dose, an additional dose of 0.75 g ZOSYN (0.67 g piperacillin and 0.08 g tazobactam) should be administered following each dialysis period on hemodialysis days. No additional dosage of ZOSYN is necessary for CAPD patients.

## **2.5 Dosage in Pediatric Patients With Appendicitis/Peritonitis or Nosocomial Pneumonia**

The recommended dosage for pediatric patients with appendicitis and/or peritonitis or nosocomial pneumonia aged 2 months of age and older, weighing up to 40 kg, and with normal renal function, is described in Table 2 [see *Use in Specific Populations (8.4)* and *Clinical Pharmacology (12.3)*].

**Table 2: Recommended Dosage of ZOSYN in Pediatric Patients 2 Months of Age and Older, Weighing Up to 40 kg, and With Normal Renal Function\*†**

<b>Age</b>	<b>Appendicitis and/or Peritonitis</b>	<b>Nosocomial Pneumonia</b>
2 months to 9 months	90 mg/kg (80 mg piperacillin and 10 mg tazobactam) <u>every 8 (eight) hours</u>	90 mg/kg (80 mg piperacillin and 10 mg tazobactam) <u>every 6 (six) hours</u>
Older than 9 months of age	112.5 mg/kg (100 mg piperacillin and 12.5 mg tazobactam) <u>every 8 (eight) hours</u>	112.5 mg/kg (100 mg piperacillin and 12.5 mg tazobactam) <u>every 6 (six) hours</u>

\* Administer ZOSYN by intravenous infusion over 30 minutes

† If a dose of ZOSYN is required that does not equal 2.25 g, 3.375 g, or 4.5 g, ZOSYN injection in GALAXY Containers is not recommended for use and an alternative formulation of ZOSYN should be considered [see *Use in Specific Populations (8.4)*].

Pediatric patients weighing over 40 kg and with normal renal function should receive the

adult dose [see *Dosage and Administration (2.2, 2.3)*].

Dosage of ZOSYN in pediatric patients with renal impairment has not been determined.

## **2.6 Directions for Use of ZOSYN Injection**

### Important Administration Instructions for ZOSYN Injection in GALAXY Containers

Administer ZOSYN Injection in GALAXY Containers using sterile equipment, after thawing to room temperature.

ZOSYN containing EDTA is compatible for co-administration via a Y-site intravenous tube with Lactated Ringer's injection, USP.

Do **NOT** add supplementary medication.

Unused portions of ZOSYN Injection should be discarded.

Do **NOT** use plastic containers in series connections. Such use could result in air embolism due to residual air being drawn from the primary container before administration of the fluid from the secondary container is complete.

Handle frozen product containers with care. Product containers may be fragile in the frozen state.

### Thawing of Plastic Container

Thaw frozen container at room temperature 20°C to 25°C [68°F to 77°F] or under refrigeration (2°C to 8°C [36°F to 46°F]). Do not force thaw by immersion in water baths or by microwave irradiation.

Check for minute leaks by squeezing container firmly. If leaks are detected, discard solution as sterility may be impaired.

The container should be visually inspected. Components of the solution may precipitate in the frozen state and will dissolve upon reaching room temperature with little or no agitation. Potency is not affected. Agitate after solution has reached room temperature. If after visual inspection, the solution remains cloudy or if an insoluble precipitate is noted or if any seals or outlet ports are not intact, the container should be discarded.

### Administration Instructions for ZOSYN Injection in GALAXY Containers to Adult Patients

Administer by infusion over a period of at least 30 minutes. During the infusion it is desirable to discontinue the primary infusion solution.

### Administration Instruction for ZOSYN Injection in GALAXY Containers to Pediatric Patients Weighing up to 40 kg

If a dose of ZOSYN is required that does not equal 2.25 g, 3.375 g, or 4.5 g, ZOSYN injection in GALAXY Containers is not recommended for use and an alternative formulation of ZOSYN should be considered.

### Storage of ZOSYN Injection

Store in a freezer capable of maintaining a temperature of -20°C (-4°F).

For GALAXY Containers, the thawed solution is stable for 14 days under refrigeration (2°C to 8°C [36°F to 46°F]) or 24 hours at room temperature 20°C to 25°C [68°F to 77°F]. Do not refreeze thawed ZOSYN Injection.

## 2.7 Compatibility With Aminoglycosides

Due to the in vitro inactivation of aminoglycosides by piperacillin, ZOSYN and aminoglycosides are recommended for separate administration. ZOSYN and aminoglycosides should be reconstituted, diluted, and administered separately when concomitant therapy with aminoglycosides is indicated [see *Drug Interactions (7.1)*].

In circumstances where co-administration via Y-site is necessary, ZOSYN formulations containing EDTA are compatible for simultaneous co-administration via Y-site infusion only with the following aminoglycosides under the following conditions:

**Table 3: Compatibility with Aminoglycosides**

<b>Aminoglycoside</b>	<b>ZOSYN Dose (grams)</b>	<b>Aminoglycoside Concentration Range*</b> <b>(mg/mL)</b>	<b>Acceptable Diluents</b>
Amikacin	2.25 3.375 4.5	1.75 - 7.5	0.9% sodium chloride or 5% dextrose
Gentamicin	2.25 3.375† 4.5	0.7 - 3.32	0.9% sodium chloride or 5% dextrose

\* The concentration ranges in Table 3 are based on administration of the aminoglycoside in divided doses (10–15 mg/kg/day in two daily doses for amikacin and 3–5 mg/kg/day in three daily doses for gentamicin). Administration of amikacin or gentamicin in a single daily dose or in doses exceeding those stated above via Y-site with ZOSYN containing EDTA has not been evaluated. See package insert for each aminoglycoside for complete Dosage and Administration instructions.

† ZOSYN 3.375 g per 50 mL GALAXY Containers are NOT compatible with gentamicin for co-administration via a Y-site due to the higher concentrations of piperacillin and tazobactam.

Only the concentration and diluents for amikacin or gentamicin with the dosages of ZOSYN listed above have been established as compatible for co-administration via Y-site infusion. Simultaneous co-administration via Y-site infusion in any manner other than listed above may result in inactivation of the aminoglycoside by ZOSYN.

ZOSYN is not compatible with tobramycin for simultaneous co-administration via Y-site infusion. Compatibility of ZOSYN with other aminoglycosides has not been established.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

## 3 DOSAGE FORMS AND STRENGTHS

ZOSYN® (piperacillin and tazobactam) Injection is supplied in GALAXY Containers as a frozen, iso-osmotic, sterile, non-pyrogenic solution in single-dose plastic containers:

2.25 g (piperacillin sodium equivalent to 2 g piperacillin and tazobactam sodium equivalent to 0.25 g tazobactam) in 50 mL.

3.375 g (piperacillin sodium equivalent to 3 g piperacillin and tazobactam sodium equivalent to 0.375 g tazobactam) in 50 mL.

4.5 g (piperacillin sodium equivalent to 4 g piperacillin and tazobactam sodium equivalent to 0.5 g tazobactam) in 100 mL.

## **4 CONTRAINDICATIONS**

ZOSYN is contraindicated in patients with a history of allergic reactions to any of the penicillins, cephalosporins, or beta-lactamase inhibitors.

## **5 WARNINGS AND PRECAUTIONS**

### **5.1 Hypersensitivity Adverse Reactions**

Serious and occasionally fatal hypersensitivity (anaphylactic/anaphylactoid) reactions (including shock) have been reported in patients receiving therapy with ZOSYN. These reactions are more likely to occur in individuals with a history of penicillin, cephalosporin, or carbapenem hypersensitivity or a history of sensitivity to multiple allergens. Before initiating therapy with ZOSYN, careful inquiry should be made concerning previous hypersensitivity reactions. If an allergic reaction occurs, ZOSYN should be discontinued and appropriate therapy instituted.

### **5.2 Severe Cutaneous Adverse Reactions**

ZOSYN may cause severe cutaneous adverse reactions, such as Stevens-Johnson syndrome, toxic epidermal necrolysis, drug reaction with eosinophilia and systemic symptoms, and acute generalized exanthematous pustulosis. If patients develop a skin rash they should be monitored closely and ZOSYN discontinued if lesions progress.

### **5.3 Hemophagocytic Lymphohistiocytosis**

Cases of hemophagocytic lymphohistiocytosis (HLH) have been reported in pediatric and adult patients treated with ZOSYN. Signs and symptoms of HLH may include fever, rash, lymphadenopathy, hepatosplenomegaly and cytopenia. If HLH is suspected, discontinue ZOSYN immediately and institute appropriate management.

### **5.4 Hematologic Adverse Reactions**

Bleeding manifestations have occurred in some patients receiving beta-lactam drugs, including piperacillin. These reactions have sometimes been associated with abnormalities of coagulation tests such as clotting time, platelet aggregation and prothrombin time, and are more likely to occur in patients with renal failure. If bleeding manifestations occur, ZOSYN should be discontinued and appropriate therapy instituted.

The leukopenia/neutropenia associated with ZOSYN administration appears to be reversible and most frequently associated with prolonged administration.

Periodic assessment of hematopoietic function should be performed, especially with prolonged therapy, i.e.,  $\geq 21$  days [see *Adverse Reactions (6.1)*].

## 5.5 Central Nervous System Adverse Reactions

As with other penicillins, ZOSYN may cause neuromuscular excitability or seizures. Patients receiving higher doses, especially patients with renal impairment may be at greater risk for central nervous system adverse reactions. Closely monitor patients with renal impairment or seizure disorders for signs and symptoms of neuromuscular excitability or seizures [see *Adverse Reactions (6.2)*].

## 5.6 Nephrotoxicity in Critically Ill Patients

The use of ZOSYN was found to be an independent risk factor for renal failure and was associated with delayed recovery of renal function as compared to other beta-lactam antibacterial drugs in a randomized, multicenter, controlled trial in critically ill patients [see *Adverse Reactions (6.1)*]. Based on this study, alternative treatment options should be considered in the critically ill population. If alternative treatment options are inadequate or unavailable, monitor renal function during treatment with ZOSYN [see *Dosage and Administration (2.4)*].

Combined use of piperacillin and tazobactam and vancomycin may be associated with an increased incidence of acute kidney injury [see *Drug Interactions (7.3)*].

## 5.7 Electrolyte Effects

ZOSYN contains a total of 2.84 mEq (65 mg) of Na<sup>+</sup> (sodium) per gram of piperacillin in the combination product. This should be considered when treating patients requiring restricted salt intake. Periodic electrolyte determinations should be performed in patients with low potassium reserves, and the possibility of hypokalemia should be kept in mind with patients who have potentially low potassium reserves and who are receiving cytotoxic therapy or diuretics.

## 5.8 *Clostridioides difficile*-Associated Diarrhea

*Clostridioides difficile*-associated diarrhea (CDAD) has been reported with use of nearly all antibacterial agents, including ZOSYN, and may range in severity from mild diarrhea to fatal colitis. Treatment with antibacterial agents alters the normal flora of the colon leading to overgrowth of *C. difficile*.

*C. difficile* produces toxins A and B which contribute to the development of CDAD. Hypertoxin producing strains of *C. difficile* cause increased morbidity and mortality, as these infections can be refractory to antimicrobial therapy and may require colectomy. CDAD must be considered in all patients who present with diarrhea following antibacterial drug use. Careful medical history is necessary since CDAD has been reported to occur over two months after the administration of antibacterial agents.

If CDAD is suspected or confirmed, ongoing antibacterial drug use not directed against *C. difficile* may need to be discontinued. Appropriate fluid and electrolyte management, protein supplementation, antibacterial treatment of *C. difficile*, and surgical evaluation should be instituted as clinically indicated.

## 5.9 Development of Drug-Resistant Bacteria

Prescribing ZOSYN in the absence of a proven or strongly suspected bacterial infection or a prophylactic indication is unlikely to provide benefit to the patient and increases the risk of development of drug-resistant bacteria.

## 6 ADVERSE REACTIONS

The following clinically significant adverse reactions are described elsewhere in the labeling:

- Hypersensitivity Adverse Reactions [see Warnings and Precautions (5.1)]
- Severe Cutaneous Adverse Reactions [see Warnings and Precautions (5.2)]
- Hemophagocytic Lymphohistiocytosis [see Warnings and Precautions (5.3)]
- Hematologic Adverse Reactions [see Warnings and Precautions (5.4)]
- Central Nervous System Adverse Reactions [see Warnings and Precautions (5.5)]
- Nephrotoxicity in Critically Ill Patients [see Warnings and Precautions (5.6)]
- *Clostridioides difficile*-Associated Diarrhea [see Warnings and Precautions (5.8)]

### 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

#### Clinical Trials in Adult Patients

During the initial clinical investigations, 2621 patients worldwide were treated with ZOSYN in phase 3 trials. In the key North American monotherapy clinical trials (n=830 patients), 90% of the adverse events reported were mild to moderate in severity and transient in nature. However, in 3.2% of the patients treated worldwide, ZOSYN was discontinued because of adverse events primarily involving the skin (1.3%), including rash and pruritus; the gastrointestinal system (0.9%), including diarrhea, nausea, and vomiting; and allergic reactions (0.5%).

**Table 4: Adverse Reactions from ZOSYN Monotherapy Clinical Trials**

---

#### **System Organ Class**

Adverse Reaction

---

#### **Gastrointestinal disorders**

Diarrhea (11.3%)

Constipation (7.7%)

Nausea (6.9%)

Vomiting (3.3%)

Dyspepsia (3.3%)

Abdominal pain (1.3%)

#### **General disorders and administration site conditions**

Fever (2.4%)

Injection site reaction ( $\leq 1\%$ )

Rigors ( $\leq 1\%$ )

#### **Immune system disorders**

Anaphylaxis ( $\leq 1\%$ )

#### **Infections and infestations**

Candidiasis (1.6%)

Pseudomembranous colitis ( $\leq 1\%$ )

**Metabolism and nutrition disorders**

Hypoglycemia ( $\leq 1\%$ )

**Musculoskeletal and connective tissue disorders**

Myalgia ( $\leq 1\%$ )

Arthralgia ( $\leq 1\%$ )

**Nervous system disorders**

Headache (7.7%)

**Psychiatric disorders**

Insomnia (6.6%)

**Skin and subcutaneous tissue disorders**

Rash (4.2%, including maculopapular, bullous, and urticarial)

Pruritus (3.1%)

Purpura ( $\leq 1\%$ )

**Vascular disorders**

Phlebitis (1.3%)

Thrombophlebitis ( $\leq 1\%$ )

Hypotension ( $\leq 1\%$ )

Flushing ( $\leq 1\%$ )

**Respiratory, thoracic and mediastinal disorders**

Epistaxis ( $\leq 1\%$ )

---

Nosocomial Pneumonia Trials

Two trials of nosocomial lower respiratory tract infections were conducted. In one study, 222 patients were treated with ZOSYN in a dosing regimen of 4.5 g every 6 hours in combination with an aminoglycoside and 215 patients were treated with imipenem/cilastatin (500 mg/500 mg every 6 hours) in combination with an aminoglycoside. In this trial, treatment-emergent adverse events were reported by 402 patients, 204 (91.9%) in the piperacillin and tazobactam group and 198 (92.1%) in the imipenem/cilastatin group. Twenty-five (11.0%) patients in the piperacillin and tazobactam group and 14 (6.5%) in the imipenem/cilastatin group ( $p > 0.05$ ) discontinued treatment due to an adverse event.

The second trial used a dosing regimen of 3.375 g given every 4 hours with an aminoglycoside.

**Table 5: Adverse Reactions from ZOSYN Plus Aminoglycoside Clinical Trials\***

---

**System Organ Class**

Adverse Reaction

---

**Blood and lymphatic system disorders**

Thrombocythemia (1.4%)

Anemia ( $\leq 1\%$ )

Thrombocytopenia ( $\leq 1\%$ )

Eosinophilia ( $\leq 1\%$ )

**Gastrointestinal disorders**

Diarrhea (20%)  
Constipation (8.4%)  
Nausea (5.8%)  
Vomiting (2.7%)  
Dyspepsia (1.9%)  
Abdominal pain (1.8%)  
Stomatitis ( $\leq 1\%$ )

#### **General disorders and administration site conditions**

Fever (3.2%)  
Injection site reaction ( $\leq 1\%$ )

#### **Infections and infestations**

Oral candidiasis (3.9%)  
Candidiasis (1.8%)

#### **Investigations**

BUN increased (1.8%)  
Blood creatinine increased (1.8%)  
Liver function test abnormal (1.4%)  
Alkaline phosphatase increased ( $\leq 1\%$ )  
Aspartate aminotransferase increased ( $\leq 1\%$ )  
Alanine aminotransferase increased ( $\leq 1\%$ )

#### **Metabolism and nutrition disorders**

Hypoglycemia ( $\leq 1\%$ )  
Hypokalemia ( $\leq 1\%$ )

#### **Nervous system disorders**

Headache (4.5%)

#### **Psychiatric disorders**

Insomnia (4.5%)

#### **Renal and urinary disorders**

Renal failure ( $\leq 1\%$ )

#### **Skin and subcutaneous tissue disorders**

Rash (3.9%)  
Pruritus (3.2%)

#### **Vascular disorders**

Thrombophlebitis (1.3%)  
Hypotension (1.3%)

---

\* For adverse drug reactions that appeared in both studies the higher frequency is presented.

#### Other Trials: Nephrotoxicity

In a randomized, multicenter, controlled trial in 1200 adult critically ill patients, piperacillin and tazobactam was found to be a risk factor for renal failure (odds ratio 1.7, 95% CI 1.18 to 2.43), and associated with delayed recovery of renal function as compared to other beta-lactam antibacterial drugs<sup>1</sup> [see *Warnings and Precautions* (5.6)].

Adverse Laboratory Changes (Seen During Clinical Trials)

Of the trials reported, including that of nosocomial lower respiratory tract infections in

which a higher dose of ZOSYN was used in combination with an aminoglycoside, changes in laboratory parameters include:

*Hematologic*—decreases in hemoglobin and hematocrit, thrombocytopenia, increases in platelet count, eosinophilia, leukopenia, neutropenia. These patients were withdrawn from therapy; some had accompanying systemic symptoms (e.g., fever, rigors, chills)

*Coagulation*—positive direct Coombs' test, prolonged prothrombin time, prolonged partial thromboplastin time

*Hepatic*—transient elevations of AST (SGOT), ALT (SGPT), alkaline phosphatase, bilirubin

*Renal*—increases in serum creatinine, blood urea nitrogen

Additional laboratory events include abnormalities in electrolytes (i.e., increases and decreases in sodium, potassium, and calcium), hyperglycemia, decreases in total protein or albumin, blood glucose decreased, gamma-glutamyltransferase increased, hypokalemia, and bleeding time prolonged.

### Clinical Trials in Pediatric Patients

Clinical studies of ZOSYN in pediatric patients suggest a similar safety profile to that seen in adults.

In a prospective, randomized, comparative, open-label clinical trial of pediatric patients, 2 to 12 years of age, with intra-abdominal infections (including appendicitis and/or peritonitis), 273 patients were treated with ZOSYN 112.5 mg/kg given IV every 8 hours and 269 patients were treated with cefotaxime (50 mg/kg) plus metronidazole (7.5 mg/kg) every 8 hours. In this trial, treatment-emergent adverse events were reported by 146 patients, 73 (26.7%) in the ZOSYN group and 73 (27.1%) in the cefotaxime/metronidazole group. Six patients (2.2%) in the ZOSYN group and 5 patients (1.9%) in the cefotaxime/metronidazole group discontinued due to an adverse event.

In a retrospective, cohort study, 140 pediatric patients 2 months to less than 18 years of age with nosocomial pneumonia were treated with ZOSYN and 267 patients were treated with comparators (which included ticarcillin-clavulanate, carbapenems, ceftazidime, cefepime, or ciprofloxacin). The rates of serious adverse reactions were generally similar between the ZOSYN and comparator groups, including patients aged 2 months to 9 months treated with ZOSYN 90 mg/kg IV every 6 hours and patients older than 9 months and less than 18 years of age treated with ZOSYN 112.5 mg/kg IV every 6 hours.

## **6.2 Postmarketing Experience**

In addition to the adverse drug reactions identified in clinical trials in Table 4 and Table 5, the following adverse reactions have been identified during post-approval use of ZOSYN. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

*Hepatobiliary*—hepatitis, jaundice

*Hematologic*—hemolytic anemia, agranulocytosis, pancytopenia

*Immune*—hypersensitivity reactions, anaphylactic/anaphylactoid reactions (including shock), hemophagocytic lymphohistiocytosis (HLH)

*Renal*—interstitial nephritis

*Nervous system disorders*—seizures

*Psychiatric disorders*—delirium

*Respiratory*—eosinophilic pneumonia

*Skin and Appendages*—erythema multiforme, Stevens-Johnson syndrome, toxic epidermal necrolysis, drug reaction with eosinophilia and systemic symptoms, (DRESS), acute generalized exanthematous pustulosis (AGEP), dermatitis exfoliative

Postmarketing experience with ZOSYN in pediatric patients suggests a similar safety profile to that seen in adults.

### **6.3 Additional Experience with Piperacillin**

The following adverse reaction has also been reported for piperacillin for injection:

*Skeletal*—prolonged neuromuscular blockade [see *Drug Interactions (7.5)*].

## **7 DRUG INTERACTIONS**

### **7.1 Aminoglycosides**

Piperacillin may inactivate aminoglycosides by converting them to microbiologically inert amides.

#### *In vivo* inactivation:

When aminoglycosides are administered in conjunction with piperacillin to patients with end-stage renal disease requiring hemodialysis, the concentrations of the aminoglycosides (especially tobramycin) may be significantly reduced and should be monitored.

Sequential administration of ZOSYN and tobramycin to patients with either normal renal function or mild to moderate renal impairment has been shown to modestly decrease serum concentrations of tobramycin but no dosage adjustment is considered necessary.

#### *In vitro* inactivation:

Due to the *in vitro* inactivation of aminoglycosides by piperacillin, ZOSYN and aminoglycosides are recommended for separate administration. ZOSYN and aminoglycosides should be reconstituted, diluted, and administered separately when concomitant therapy with aminoglycosides is indicated. ZOSYN, which contains EDTA, is compatible with amikacin and gentamicin for simultaneous Y-site infusion in certain diluents and at specific concentrations. ZOSYN is not compatible with tobramycin for simultaneous Y-site infusion [see *Dosage and Administration (2.7)*].

### **7.2 Probenecid**

Probenecid administered concomitantly with ZOSYN prolongs the half-life of piperacillin by 21% and that of tazobactam by 71% because probenecid inhibits tubular renal secretion of both piperacillin and tazobactam. Probenecid should not be co-administered with ZOSYN unless the benefit outweighs the risk.

### **7.3 Vancomycin**

Studies have detected an increased incidence of acute kidney injury in patients concomitantly administered piperacillin and tazobactam and vancomycin as compared to vancomycin alone [*see Warnings and Precautions (5.6)*].

Monitor kidney function in patients concomitantly administered with piperacillin and tazobactam and vancomycin.

No pharmacokinetic interactions have been noted between piperacillin and tazobactam and vancomycin.

### **7.4 Anticoagulants**

Coagulation parameters should be tested more frequently and monitored regularly during simultaneous administration of high doses of heparin, oral anticoagulants, or other drugs that may affect the blood coagulation system or the thrombocyte function [*see Warnings and Precautions (5.4)*].

### **7.5 Vecuronium**

Piperacillin when used concomitantly with vecuronium has been implicated in the prolongation of the neuromuscular blockade of vecuronium. ZOSYN could produce the same phenomenon if given along with vecuronium. Due to their similar mechanism of action, it is expected that the neuromuscular blockade produced by any of the non-depolarizing neuromuscular blockers could be prolonged in the presence of piperacillin. Monitor for adverse reactions related to neuromuscular blockade (see package insert for vecuronium bromide).

### **7.6 Methotrexate**

Limited data suggests that co-administration of methotrexate and piperacillin may reduce the clearance of methotrexate due to competition for renal secretion. The impact of tazobactam on the elimination of methotrexate has not been evaluated. If concurrent therapy is necessary, serum concentrations of methotrexate as well as the signs and symptoms of methotrexate toxicity should be frequently monitored.

### **7.7 Effects on Laboratory Tests**

There have been reports of positive test results using the Bio-Rad Laboratories Platelia Aspergillus EIA test in patients receiving piperacillin and tazobactam injection who were subsequently found to be free of Aspergillus infection. Cross-reactions with non-Aspergillus polysaccharides and polyfuranoses with the Bio-Rad Laboratories Platelia Aspergillus EIA test have been reported. Therefore, positive test results in patients receiving piperacillin and tazobactam should be interpreted cautiously and confirmed by other diagnostic methods.

As with other penicillins, the administration of ZOSYN may result in a false-positive reaction for glucose in the urine using a copper-reduction method (CLINITEST®). It is recommended that glucose tests based on enzymatic glucose oxidase reactions be used.

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

#### Risk Summary

Piperacillin and tazobactam cross the placenta in humans. However, there are insufficient data with piperacillin and/or tazobactam in pregnant women to inform a drug-associated risk for major birth defects and miscarriage. No fetal structural abnormalities were observed in rats or mice when piperacillin and tazobactam was administered intravenously during organogenesis at doses 1 to 2 times and 2 to 3 times the human dose of piperacillin and tazobactam, respectively, based on body-surface area ( $\text{mg}/\text{m}^2$ ). However, fetotoxicity in the presence of maternal toxicity was observed in developmental toxicity and peri/postnatal studies conducted in rats (intraperitoneal administration prior to mating and throughout gestation or from gestation day 17 through lactation day 21) at doses less than the maximum recommended human daily dose based on body-surface area ( $\text{mg}/\text{m}^2$ ) (see *Data*).

The background risk of major birth defects and miscarriage for the indicated population is unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2–4% and 15–20%, respectively.

#### Data

##### *Animal Data*

In embryo-fetal development studies in mice and rats, pregnant animals received intravenous doses of piperacillin and tazobactam up to 3000/750  $\text{mg}/\text{kg}/\text{day}$  during the period of organogenesis. There was no evidence of teratogenicity up to the highest dose evaluated, which is 1 to 2 times and 2 to 3 times the human dose of piperacillin and tazobactam, in mice and rats respectively, based on body-surface area ( $\text{mg}/\text{m}^2$ ). Fetal body weights were reduced in rats at maternally toxic doses at or above 500/62.5  $\text{mg}/\text{kg}/\text{day}$ , minimally representing 0.4 times the human dose of both piperacillin and tazobactam based on body-surface area ( $\text{mg}/\text{m}^2$ ).

A fertility and general reproduction study in rats using intraperitoneal administration of tazobactam or the combination piperacillin and tazobactam prior to mating and through the end of gestation, reported a decrease in litter size in the presence of maternal toxicity at 640  $\text{mg}/\text{kg}/\text{day}$  tazobactam (4 times the human dose of tazobactam based on body-surface area), and decreased litter size and an increase in fetuses with ossification delays and variations of ribs, concurrent with maternal toxicity at  $\geq 640/160$   $\text{mg}/\text{kg}/\text{day}$  piperacillin and tazobactam (0.5 times and 1 times the human dose of piperacillin and tazobactam, respectively, based on body-surface area).

Peri/postnatal development in rats was impaired with reduced pup weights, increased stillbirths, and increased pup mortality concurrent with maternal toxicity after intraperitoneal administration of tazobactam alone at doses  $\geq 320$   $\text{mg}/\text{kg}/\text{day}$  (2 times the human dose based on body surface area) or of the combination piperacillin and tazobactam at doses  $\geq 640/160$   $\text{mg}/\text{kg}/\text{day}$  (0.5 times and 1 times the human dose of piperacillin and tazobactam, respectively, based on body-surface area) from gestation day 17 through lactation day 21.

## 8.2 Lactation

### Risk Summary

Piperacillin is excreted in human milk; tazobactam concentrations in human milk have not been studied. No information is available on the effects of piperacillin and tazobactam on the breast-fed child or on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for ZOSYN and any potential adverse effects on the breastfed child from ZOSYN or from the underlying maternal condition.

## 8.4 Pediatric Use

The safety and effectiveness of ZOSYN for intra-abdominal infections, and nosocomial pneumonia have been established in pediatric patients 2 months of age and older.

Use of ZOSYN in pediatric patients 2 months of age and older with intra-abdominal infections including appendicitis and/or peritonitis is supported by evidence from well-controlled studies and pharmacokinetic studies in adults and in pediatric patients. This includes a prospective, randomized, comparative, open-label clinical trial with 542 pediatric patients 2 to 12 years of age with intra-abdominal infections (including appendicitis and/or peritonitis), in which 273 pediatric patients received piperacillin and tazobactam [see *Adverse Reactions (6.1) and Clinical Pharmacology (12.3)*].

Use of ZOSYN in pediatric patients 2 months of age and older with nosocomial pneumonia is supported by evidence from well-controlled studies in adults with nosocomial pneumonia, a simulation study performed with a population pharmacokinetic model, and a retrospective, cohort study of pediatric patients with nosocomial pneumonia in which 140 pediatric patients were treated with ZOSYN and 267 patients treated with comparators (which included ticarcillin-clavulanate, carbapenems, ceftazidime, cefepime, or ciprofloxacin) [see *Adverse Reactions (6.1) and Clinical Pharmacology (12.3)*].

Because of the limitations of the available strengths and administration requirements (i.e., administration of fractional doses is not recommended) of ZOSYN Injection supplied in GALAXY Containers, and to avoid unintentional overdose, this product is not recommended for use if a dose of ZOSYN Injection in GALAXY Containers that does not equal 2.25 g, 3.375 g, or 4.5 g is required and an alternative formulation of ZOSYN should be considered [see *Dosage and Administration (2.1, 2.5, and 2.6)*].

The safety and effectiveness of ZOSYN have not been established in pediatric patients less than 2 months of age [see *Clinical Pharmacology (12) and Dosage and Administration (2)*].

Dosage of ZOSYN in pediatric patients with renal impairment has not been determined.

## 8.5 Geriatric Use

Patients over 65 years are not at an increased risk of developing adverse effects solely because of age. However, dosage should be adjusted in the presence of renal impairment [see *Dosage and Administration (2)*].

In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic,

renal, or cardiac function, and of concomitant disease or other drug therapy.

ZOSYN contains 65 mg (2.84 mEq) of sodium per gram of piperacillin in the combination product. At the usual recommended doses, patients would receive between 780 and 1040 mg/day (34.1 and 45.5 mEq) of sodium. The geriatric population may respond with a blunted natriuresis to salt loading. This may be clinically important with regard to such diseases as congestive heart failure.

This drug is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

## **8.6 Renal Impairment**

In patients with creatinine clearance  $\leq$  40 mL/min and dialysis patients (hemodialysis and CAPD), the intravenous dose of ZOSYN should be reduced to the degree of renal function impairment [see *Dosage and Administration* (2)].

## **8.7 Hepatic Impairment**

Dosage adjustment of ZOSYN is not warranted in patients with hepatic cirrhosis [see *Clinical Pharmacology* (12.3)].

## **8.8 Patients with Cystic Fibrosis**

As with other semisynthetic penicillins, piperacillin therapy has been associated with an increased incidence of fever and rash in cystic fibrosis patients.

## **10 OVERDOSAGE**

There have been postmarketing reports of overdose with piperacillin and tazobactam. The majority of those events experienced, including nausea, vomiting, and diarrhea, have also been reported with the usual recommended dosages. Patients may experience neuromuscular excitability or seizures if higher than recommended doses are given intravenously (particularly in the presence of renal failure) [see *Warnings and Precautions* (5.5)].


Treatment should be supportive and symptomatic according to the patient's clinical presentation. Excessive serum concentrations of either piperacillin or tazobactam may be reduced by hemodialysis. Following a single 3.375 g dose of piperacillin and tazobactam, the percentage of the piperacillin and tazobactam dose removed by hemodialysis was approximately 31% and 39%, respectively [see *Clinical Pharmacology* (12)].

## **11 DESCRIPTION**


ZOSYN (piperacillin and tazobactam) Injection is an injectable antibacterial combination product consisting of the semisynthetic antibacterial piperacillin sodium and the beta-lactamase inhibitor tazobactam sodium for intravenous administration.

Piperacillin sodium is derived from D(-)- $\alpha$ -aminobenzyl-penicillin. The chemical name of piperacillin sodium is sodium (2S,5R,6R)-6-[(R)-2-(4-ethyl-2,3-dioxo-1-piperazine-

carboxamido)-2-phenylacetamido]-3,3-dimethyl-7-oxo-4-thia-1-azabicyclo[3.2.0]heptane-2-carboxylate. The chemical formula is  $C_{23}H_{26}N_5NaO_7S$  and the molecular weight is 539.5. The chemical structure of piperacillin sodium is:



Tazobactam sodium, a derivative of the penicillin nucleus, is a penicillanic acid sulfone. Its chemical name is sodium (2*S*,3*S*,5*R*)-3-methyl-7-oxo-3-(1*H*-1,2,3-triazol-1-ylmethyl)-4-thia-1-azabicyclo[3.2.0]heptane-2-carboxylate-4,4-dioxide. The chemical formula is  $C_{10}H_{11}N_4NaO_5S$  and the molecular weight is 322.3. The chemical structure of tazobactam sodium is:



ZOSYN Injection in the GALAXY Container is a frozen iso-osmotic sterile non-pyrogenic premixed solution. The components and dosage formulations are given in the table below:

**Table 6: ZOSYN In GALAXY Containers Premixed Frozen Solution**

Component*	Function	Dosage Formulations		
		2.25 g/50 mL	3.375 g/50 mL	4.5 g/100 mL
Piperacillin	active ingredient	2 g	3 g	4 g
Tazobactam	beta-lactamase inhibitor	250 mg	375 mg	500 mg
Dextrose Hydrus	osmolality adjusting agent	1 g	350 mg	2 g
Sodium Citrate Dihydrate	buffering agent	100 mg	150 mg	200 mg
Edetate Disodium	metal chelator	0.5 mg	0.75 mg	1 mg

Dihydrate				
Water for Injection	solvent	q.s. 50 mL	q.s. 50 mL	q.s. 100 mL

\* Piperacillin and tazobactam are present in the formulation as sodium salts. Dextrose hydrous, sodium citrate dihydrate, and edetate disodium dihydrate amounts are approximate.

ZOSYN contains a total of 2.84 mEq (65 mg) of sodium (Na<sup>+</sup>) per gram of piperacillin in the combination product.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

ZOSYN is an antibacterial drug [see *Microbiology (12.4)*].

### 12.2 Pharmacodynamics

The pharmacodynamic parameter for piperacillin and tazobactam that is most predictive of clinical and microbiological efficacy is time above MIC.

### 12.3 Pharmacokinetics

The mean and coefficients of variation (CV%) for the pharmacokinetic parameters of piperacillin and tazobactam after multiple intravenous doses are summarized in Table 7.

**Table 7: Mean (CV%) Piperacillin and Tazobactam PK Parameters**

Piperacillin						
Piperacillin and Tazobactam Dose*	C <sub>max</sub> (mcg/mL)	AUC <sup>†</sup> (mcg•h/mL)	CL (mL/min)	V (L)	T <sub>1/2</sub> (h)	CL <sub>R</sub> (mL/min)
2.25 g	134	131 [14]	257	17.4	0.79	--
3.375 g	242	242 [10]	207	15.1	0.84	140
4.5 g	298	322 [16]	210	15.4	0.84	--
Tazobactam						
Piperacillin and Tazobactam Dose*	C <sub>max</sub> (mcg/mL)	AUC <sup>†</sup> (mcg•h/mL)	CL (mL/min)	V (L)	T <sub>1/2</sub> (h)	CL <sub>R</sub> (mL/min)
2.25 g	15	16.0 [21]	258	17.0	0.77	--
3.375 g	24	25.0 [8]	251	14.8	0.68	166
4.5 g	34	39.8 [15]	206	14.7	0.82	--

C<sub>max</sub> : maximum observed concentration, AUC: Area under the curve, CL=clearance, CL<sub>R</sub>= Renal clearance

V=volume of distribution, T<sub>1/2</sub> = elimination half-life

\* Piperacillin and tazobactam were given in combination, infused over 30 minutes.

† Numbers in []parentheses are coefficients of variation [CV%].

Peak plasma concentrations of piperacillin and tazobactam are attained immediately after completion of an intravenous infusion of ZOSYN. Piperacillin plasma concentrations, following a 30-minute infusion of ZOSYN, were similar to those attained when equivalent doses of piperacillin were administered alone. Steady-state plasma concentrations of piperacillin and tazobactam were similar to those attained after the first dose due to the short half-lives of piperacillin and tazobactam.

### Distribution

Both piperacillin and tazobactam are approximately 30% bound to plasma proteins. The protein binding of either piperacillin or tazobactam is unaffected by the presence of the other compound. Protein binding of the tazobactam metabolite is negligible.

Piperacillin and tazobactam are widely distributed into tissues and body fluids including intestinal mucosa, gallbladder, lung, female reproductive tissues (uterus, ovary, and fallopian tube), interstitial fluid, and bile. Mean tissue concentrations are generally 50% to 100% of those in plasma. Distribution of piperacillin and tazobactam into cerebrospinal fluid is low in subjects with non-inflamed meninges, as with other penicillins (see Table 8).

**Table 8: Piperacillin and Tazobactam Concentrations in Selected Tissues and Fluids after Single 4 g/0.5 g 30-min IV Infusion of ZOSYN**

Tissue or Fluid	N*	Sampling period† (h)	Mean PIP Concentration Range (mg/L)	Tissue:Plasma Range	Tazo Concentration Range (mg/L)	Tazo Tissue:Plasma Range
Skin	35	0.5 - 4.5	34.8 - 94.2	0.60 - 1.1	4.0 - 7.7	0.49 - 0.93
Fatty Tissue	37	0.5 - 4.5	4.0 - 10.1	0.097 - 0.115	0.7 - 1.5	0.10 - 0.13
Muscle	36	0.5 - 4.5	9.4 - 23.3	0.29 - 0.18	1.4 - 2.7	0.18 - 0.30
Proximal Intestinal Mucosa	7	1.5 - 2.5	31.4	0.55	10.3	1.15
Distal Intestinal Mucosa	7	1.5 - 2.5	31.2	0.59	14.5	2.1
Appendix	22	0.5 - 2.5	26.5 - 64.1	0.43 - 0.53	9.1 - 18.6	0.80 - 1.35

\* Each subject provided a single sample.

† Time from the start of the infusion

### Metabolism

Piperacillin is metabolized to a minor microbiologically active desethyl metabolite. Tazobactam is metabolized to a single metabolite that lacks pharmacological and antibacterial activities.

### Excretion

Following single or multiple ZOSYN doses to healthy subjects, the plasma half-life of piperacillin and of tazobactam ranged from 0.7 to 1.2 hours and was unaffected by

dose or duration of infusion.

Both piperacillin and tazobactam are eliminated via the kidney by glomerular filtration and tubular secretion. Piperacillin is excreted rapidly as unchanged drug with 68% of the administered dose excreted in the urine. Tazobactam and its metabolite are eliminated primarily by renal excretion with 80% of the administered dose excreted as unchanged drug and the remainder as the single metabolite. Piperacillin, tazobactam and desethyl piperacillin are also secreted into the bile.

## Specific Populations

### *Renal Impairment*

After the administration of single doses of piperacillin and tazobactam to subjects with renal impairment, the half-life of piperacillin and of tazobactam increases with decreasing creatinine clearance. At creatinine clearance below 20 mL/min, the increase in half-life is twofold for piperacillin and fourfold for tazobactam compared to subjects with normal renal function. Dosage adjustments for ZOSYN are recommended when creatinine clearance is below 40 mL/min in patients receiving the usual recommended daily dose of ZOSYN. See *Dosage and Administration (2)* for specific recommendations for the treatment of patients with renal-impairment.

Hemodialysis removes 30% to 40% of a piperacillin and tazobactam dose with an additional 5% of the tazobactam dose removed as the tazobactam metabolite. Peritoneal dialysis removes approximately 6% and 21% of the piperacillin and tazobactam doses, respectively, with up to 16% of the tazobactam dose removed as the tazobactam metabolite. For dosage recommendations for patients undergoing hemodialysis [see *Dosage and Administration (2)*].

### *Hepatic Impairment*

The half-life of piperacillin and of tazobactam increases by approximately 25% and 18%, respectively, in patients with hepatic cirrhosis compared to healthy subjects. However, this difference does not warrant dosage adjustment of ZOSYN due to hepatic cirrhosis.

### *Pediatrics*

Piperacillin and tazobactam pharmacokinetics were studied in pediatric patients 2 months of age and older. The clearance of both compounds is slower in the younger patients compared to older children and adults.

In a population PK analysis, estimated clearance for 9 month-old to 12 year-old patients was comparable to adults, with a population mean (SE) value of 5.64 (0.34) mL/min/kg. The piperacillin clearance estimate is 80% of this value for pediatric patients 2 - 9 months old. In patients younger than 2 months of age, clearance of piperacillin is slower compared to older children; however, it is not adequately characterized for dosing recommendations. The population mean (SE) for piperacillin volume of distribution is 0.243 (0.011) L/kg and is independent of age.

### *Geriatrics*

The impact of age on the pharmacokinetics of piperacillin and tazobactam was evaluated in healthy male subjects, aged 18 - 35 years (n=6) and aged 65 to 80 years (n=12). Mean half-life for piperacillin and tazobactam was 32% and 55% higher, respectively, in the elderly compared to the younger subjects. This difference may be due to age-related changes in creatinine clearance.

## Race

The effect of race on piperacillin and tazobactam was evaluated in healthy male volunteers. No difference in piperacillin or tazobactam pharmacokinetics was observed between Asian (n=9) and Caucasian (n=9) healthy volunteers who received single 4/0.5 g doses.

## Drug Interactions

The potential for pharmacokinetic drug interactions between ZOSYN and aminoglycosides, probenecid, vancomycin, heparin, vecuronium, and methotrexate has been evaluated [see *Drug Interactions (7)*].

## 12.4 Microbiology

### Mechanism of Action

Piperacillin sodium exerts bactericidal activity by inhibiting septum formation and cell wall synthesis of susceptible bacteria. In vitro, piperacillin is active against a variety of gram-positive and gram-negative aerobic and anaerobic bacteria. Tazobactam sodium has little clinically relevant in vitro activity against bacteria due to its reduced affinity to penicillin-binding proteins. It is, however, a beta-lactamase inhibitor of the Molecular class A enzymes, including Richmond-Sykes class III (Bush class 2b & 2b') penicillinases and cephalosporinases. It varies in its ability to inhibit class II and IV (2a & 4) penicillinases. Tazobactam does not induce chromosomally-mediated beta-lactamases at tazobactam concentrations achieved with the recommended dosage regimen.

### Antimicrobial Activity

ZOSYN has been shown to be active against most isolates of the following microorganisms, both in vitro and in clinical infections [see *Indications and Usage (1)*]:

#### Aerobic bacteria

Gram-positive bacteria

*Staphylococcus aureus* (methicillin susceptible isolates only)

Gram-negative bacteria

*Acinetobacter baumannii*

*Escherichia coli*

*Haemophilus influenzae* (excluding beta-lactamase negative, ampicillin-resistant isolates)

*Klebsiella pneumoniae*

*Pseudomonas aeruginosa* (given in combination with an aminoglycoside to which the isolate is susceptible)

#### Anaerobic bacteria

*Bacteroides fragilis* group (*B. fragilis*, *B. ovatus*, *B. thetaiotaomicron*, and *B. vulgatus*)

The following in vitro data are available, but their clinical significance is unknown. At least 90 percent of the following bacteria exhibit an in vitro minimum inhibitory concentration (MIC) less than or equal to the susceptible breakpoint for piperacillin and tazobactam against isolates of similar genus or organism group. However, the efficacy of ZOSYN in treating clinical infections caused by these bacteria has not been established in adequate and well-controlled clinical trials.

## Aerobic bacteria

### Gram-positive bacteria

*Enterococcus faecalis* (ampicillin or penicillin-susceptible isolates only)

*Staphylococcus epidermidis* (methicillin susceptible isolates only)

*Streptococcus agalactiae*<sup>1</sup>

*Streptococcus pneumoniae*<sup>1</sup> (penicillin-susceptible isolates only)

*Streptococcus pyogenes*<sup>1</sup>

Viridans group streptococci<sup>1</sup>

### Gram-negative bacteria

*Citrobacter koseri*

*Moraxella catarrhalis*

*Morganella morganii*

*Neisseria gonorrhoeae*

*Proteus mirabilis*

*Proteus vulgaris*

*Serratia marcescens*

*Providencia stuartii*

*Providencia rettgeri*

*Salmonella enterica*

## Anaerobic bacteria

*Clostridium perfringens*

*Bacteroides distasonis*

*Prevotella melaninogenica*

## Susceptibility Testing

For specific information regarding susceptibility test interpretive criteria, and associated test methods and quality control standards recognized by FDA for this drug, please see: <https://www.fda.gov/STIC>.

---

<sup>1</sup> These are not beta-lactamase producing bacteria and, therefore, are susceptible to piperacillin alone.

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

#### Carcinogenesis

Long-term carcinogenicity studies in animals have not been conducted with piperacillin and tazobactam, piperacillin, or tazobactam.

#### Mutagenesis

Piperacillin and tazobactam was negative in microbial mutagenicity assays, the unscheduled DNA synthesis (UDS) test, a mammalian point mutation (Chinese hamster ovary cell HPRT) assay, and a mammalian cell (BALB/c-3T3) transformation assay. *In vivo*, piperacillin and tazobactam did not induce chromosomal aberrations in rats.

## Fertility

Reproduction studies have been performed in rats and have revealed no evidence of impaired fertility when piperacillin and tazobactam is administered intravenously up to a dose of 1280/320 mg/kg piperacillin and tazobactam, which is similar to the maximum recommended human daily dose based on body-surface area (mg/m<sup>2</sup>).

## **15 REFERENCES**

1. Jensen J-US, Hein L, Lundgren B, et al. BMJ Open 2012; 2:e000635. doi:10.1136.

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

ZOSYN<sup>®</sup> (piperacillin and tazobactam) Injection in GALAXY Containers are supplied as a frozen, iso-osmotic, sterile, nonpyrogenic solution in single-dose plastic containers as follows:

- 2.25 g (piperacillin sodium equivalent to 2 g piperacillin and tazobactam sodium equivalent to 0.25 g tazobactam) in 50 mL. Each container has 5.58 mEq (128 mg) of sodium. Supplied 24/box—NDC 0206-8860-02
- 3.375 g (piperacillin sodium equivalent to 3 g piperacillin and tazobactam sodium equivalent to 0.375 g tazobactam) in 50 mL. Each container has 8.38 mEq (192 mg) of sodium. Supplied 24/box—NDC 0206-8861-02
- 4.5 g (piperacillin sodium equivalent to 4 g piperacillin and tazobactam sodium equivalent to 0.5 g tazobactam) in 100 mL. Each container has 11.17 mEq (256 mg) of sodium. Supplied 12/box—NDC 0206-8862-02

ZOSYN<sup>®</sup> Injection in GALAXY Containers should be stored at or below -20°C (-4°F) [*see Dosage and Administration (2.6)*].

Handle frozen product containers with care. Product containers may be fragile in the frozen state.

## **17 PATIENT COUNSELING INFORMATION**

### Serious Hypersensitivity Reactions

Advise patients, their families, or caregivers that serious hypersensitivity reactions, including serious allergic cutaneous reactions, could occur with use of ZOSYN that require immediate treatment. Ask them about any previous hypersensitivity reactions to ZOSYN, other beta-lactams (including cephalosporins), or other allergens [*see Warnings and Precautions (5.2)*].

### Hemophagocytic Lymphohistiocytosis

Prior to initiation of treatment with ZOSYN, inform patients that excessive immune activation may occur with ZOSYN and that they should report signs or symptoms such as fever, rash, or lymphadenopathy to a healthcare provider immediately [*see Warnings and Precautions (5.3)*].

### Diarrhea

Advise patients, their families, or caregivers that diarrhea is a common problem caused by antibacterial drugs, including ZOSYN, which usually ends when the drug is discontinued. Sometimes after starting treatment with antibacterial drugs, patients can develop watery and bloody stools (with or without stomach cramps and fever) even as late as two or more months after having taken the last dose of the drug. If this occurs, patients should contact their physician as soon as possible [see *Warnings and Precautions (5.8)*].

### Antibacterial Resistance

Patients should be counseled that antibacterial drugs including ZOSYN should only be used to treat bacterial infections. They do not treat viral infections (e.g., the common cold). When ZOSYN is prescribed to treat a bacterial infection, patients should be told that although it is common to feel better early in the course of therapy, the medication should be taken exactly as directed. Skipping doses or not completing the full course of therapy may (1) decrease the effectiveness of the immediate treatment and (2) increase the likelihood that bacteria will develop resistance and will not be treatable by ZOSYN or other antibacterial drugs in the future.

### Pregnancy and Lactation

Patients should be counseled that ZOSYN can cross the placenta in humans and is excreted in human milk [see *Use in Specific Populations (8.1, 8.2)*].



This product's labeling may have been updated.  
For the most recent prescribing information,  
please visit <http://www.pfizer.com>.



Distributed by  
Wyeth Pharmaceuticals LLC  
A subsidiary of Pfizer Inc.  
Philadelphia, PA 19101

CLINITEST<sup>®</sup> is a registered trademark of Siemens Healthcare Diagnostics Inc.

GALAXY is a registered trademark of Baxter International Inc.

LAB-1527-1.0

## **PRINCIPAL DISPLAY PANEL - 2.25 g Container**

**ZOSYN<sup>®</sup>**  
**(Piperacillin and Tazobactam**  
**Injection)**

**2.25 g**

**Rx only**

**GALAXY**

**Single Dose**

**Container**

**50 mL**

**Iso-osmotic**

**NDC 0206-8860-01**

**Code 2G3575**

**Sterile Nonpyrogenic**

**Store at or below -20°C/-4°F.** Thaw at room temperature 20° to 25°C (68° to 77°F) or under refrigeration 2° to 8°C (36° to 46°F). **DO NOT FORCE THAW BY IMMERSION**

**IN WATER BATHS OR BY MICROWAVE IRRADIATION.** Thawed solution remains stable for 14 days under refrigeration or 24 hours at room temperature.

**Do not refreeze. Handle frozen product containers with care. Product containers may be fragile in the frozen state.**

**Dosage and Use:** For intravenous use only. As directed by a physician. See Package Insert.

**Cautions: Do not add supplementary medication or additives.** Must not be used in series connections. Check for minute leaks and solution clarity.

Each 50 mL contains: Piperacillin Sodium equivalent to 2 g Piperacillin and Tazobactam Sodium equivalent to 0.25 g Tazobactam in Water for Injection.

Approximately 1 g Dextrose Hydrous added to adjust osmolality. Approximately 100 mg Sodium Citrate Dihydrate added as a buffer. pH adjusted with sodium bicarbonate and hydrochloric acid. pH 5.5 - 6.8

Each 50 mL also contains approximately 0.5 mg Edetate Disodium Dihydrate added as a metal chelator.

GALAXY is a registered trademark of Baxter International Inc.

Distributed by Wyeth Pharmaceuticals LLC

A subsidiary of Pfizer Inc

Philadelphia, PA 19101

PL 2040 Plastic

14715600

0734001604

**PfizerHospital**

**ZOSYN<sup>®</sup>**  
*(Piperacillin and Tazobactam  
Injection)*

**2.25 g**

Rx only

**GALAXY**  
Single Dose  
Container

**50 mL**  
**Iso-osmotic**

NDC 0206-8860-01  
**Code 2G3575**  
Sterile Nonpyrogenic

Store at or below -20°C/-4°F. Thaw at room temperature 20° to 25°C (68° to 77°F) or under refrigeration 2° to 8°C (36° to 46°F). **DO NOT FORCE THAW BY IMMERSION IN WATER BATHS OR BY MICROWAVE IRRADIATION.** Thawed solution remains stable for 14 days under refrigeration or 24 hours at room temperature.

**Do not refreeze. Handle frozen product containers with care. Product containers may be fragile in the frozen state.**

**Dosage and Use:** For intravenous use only. As directed by a physician. See Package Insert.

**Cautions:** **Do not add supplementary medication or additives.** Must not be used in series connections. Check for minute leaks and solution clarity.

Each 50 mL contains: Piperacillin Sodium equivalent to 2 g Piperacillin and Tazobactam Sodium equivalent to 0.25 g Tazobactam In Water for Injection.

Approximately 1 g Dextrose Hydrated added to adjust osmolality. Approximately 100 mg Sodium Citrate Dihydrate added as a buffer. pH adjusted with sodium bicarbonate and hydrochloric acid. pH 5.5 - 6.8

Each 50 mL also contains approximately 0.5 mg Edetate Disodium Dihydrate added as a metal chelator.

GALAXY is a registered trademark of Baxter International Inc.

PL 2040 Plastic

Distributed by Wyeth Pharmaceuticals LLC  
A subsidiary of Pfizer Inc.  
Philadelphia, PA 19101



14715600  
0734001604

FPO UPC © 100%





## PRINCIPAL DISPLAY PANEL - 2.25 g Container Box Label

### ZOSYN® (Piperacillin and Tazobactam Injection)

Rx only

2.25 g

2 x 12 x 50 mL Single Dose Containers  
Contains 2 boxes of 12 of NDC 0206-8860-01

Iso-osmotic

Store at or  
below -20°C/-4°F.

Do not refreeze.

NDC 0206-8860-02

Contains 24 of NDC 0206-8860-01

Code 2G3575

**PfizerHospital**



## PRINCIPAL DISPLAY PANEL - 3.375 g Container

### ZOSYN® (Piperacillin and Tazobactam Injection)

3.375 g

Rx only

## **GALAXY**

**Single Dose  
Container**

**50 mL  
Iso-osmotic**

**NDC 0206-8861-01**

**Code 2G3576**

**Sterile Nonpyrogenic**

**Store at or below -20°C/-4°F.** Thaw at room temperature 20° to 25°C (68° to 77°F) or under refrigeration 2° to 8°C (36° to 46°F). **DO NOT FORCE THAW BY IMMERSION**

**IN WATER BATHS OR BY MICROWAVE IRRADIATION.** Thawed solution remains stable for 14 days under refrigeration or 24 hours at room temperature.

**Do not refreeze. Handle frozen product containers with care. Product containers may be fragile in the frozen state.**

**Dosage and Use:** For intravenous use only. As directed by a physician. See Package Insert.

**Cautions: Do not add supplementary medication or additives.** Must not be used in series connections. Check for minute leaks and solution clarity.

Each 50 mL contains: Piperacillin Sodium equivalent to 3 g Piperacillin and Tazobactam Sodium equivalent to 0.375 g Tazobactam in Water for Injection.

Approximately 350 mg Dextrose Hydrous added to adjust osmolality. Approximately 150 mg Sodium Citrate Dihydrate added as a buffer. pH adjusted with sodium bicarbonate and hydrochloric acid. pH 5.5 - 6.8

Each 50 mL also contains approximately 0.75 mg Edetate Disodium Dihydrate added as a metal chelator.

GALAXY is a registered trademark of Baxter International Inc.

Distributed by Wyeth Pharmaceuticals LLC

A subsidiary of Pfizer Inc

Philadelphia, PA 19101

PL 2040 Plastic

14716000

0734001603

**PfizerHospital**

**ZOSYN<sup>®</sup>**  
*(Piperacillin and Tazobactam  
Injection)*

**3.375 g**

Rx only

**GALAXY**  
Single Dose  
Container

**50 mL**  
**Iso-osmotic**

NDC 0206-8861-01  
**Code 2G3576**  
Sterile Nonpyrogenic

Store at or below -20°C/-4°F. Thaw at room temperature 20° to 25°C (68° to 77° F) or under refrigeration 2° to 8°C (36° to 46°F). **DO NOT FORCE THAW BY IMMERSION IN WATER BATHS OR BY MICROWAVE IRRADIATION.** Thawed solution remains stable for 14 days under refrigeration or 24 hours at room temperature.

**Do not refreeze. Handle frozen product containers with care. Product containers may be fragile in the frozen state.**

**Dosage and Use:** For intravenous use only. As directed by a physician. See Package Insert.  
**Cautions:** Do not add supplementary medication or additives. Must not be used in series connections. Check for minute leaks and solution clarity.

Each 50 mL contains: Piperacillin Sodium equivalent to 3 g Piperacillin and Tazobactam Sodium equivalent to 0.375 g Tazobactam in Water for Injection. Approximately 350 mg Dextrose Hydrated added to adjust osmolality. Approximately 150 mg Sodium Citrate Dihydrate added as a buffer. pH adjusted with sodium bicarbonate and hydrochloric acid. pH 5.5 - 6.8

Each 50 mL also contains approximately 0.75 mg Edetate Disodium Dihydrate added as a metal chelator.

GALAXY is a registered trademark of Baxter International Inc.

PL 2040 Plastic

Distributed by Wyeth Pharmaceuticals LLC  
A subsidiary of Pfizer Inc.  
Philadelphia, PA 19101



14716000  
0734001603

PO UPC © 100%





## PRINCIPAL DISPLAY PANEL - 3.375 g Container Box Label

### ZOSYN® (Piperacillin and Tazobactam Injection)

Rx only

3.375g

2 x 12 x 50 mL Single Dose Containers  
Contains 2 boxes of 12 of NDC 0206-8861-01  
Iso-osmotic  
Store at or  
below -20°C/-4°F.  
Do not refreeze.

NDC 0206-8861-02  
Contains 24 of NDC 0206-8861-01  
Code 2G3576

**PfizerHospital**



## PRINCIPAL DISPLAY PANEL - 4.5 g Container

### ZOSYN® (Piperacillin and Tazobactam Injection)

4.5 g

**Rx only**

**GALAXY**

**Single Dose  
Container**

**100 mL  
Iso-osmotic**

**NDC 0206-8862-01**

**Code 2G3577**

**Sterile Nonpyrogenic**

**Store at or below -20°C/-4°F.** Thaw at room temperature 20° to 25°C (68° to 77°F) or under refrigeration 2° to 8°C (36° to 46°F). **DO NOT FORCE THAW BY IMMERSION**

**IN WATER BATHS OR BY MICROWAVE IRRADIATION.** Thawed solution remains stable for 14 days under refrigeration or 24 hours at room temperature.

**Do not refreeze. Handle frozen product containers with care. Product containers may be fragile in the frozen state.**

**Dosage and Use:** For intravenous use only. As directed by a physician. See Package Insert.

**Cautions: Do not add supplementary medication or additives.** Must not be used in series connections. Check for minute leaks and solution clarity.

Each 100 mL contains: Piperacillin Sodium equivalent to 4 g Piperacillin and Tazobactam Sodium equivalent to 0.5 g Tazobactam in Water for Injection.

Approximately 2 g Dextrose Hydrous added to adjust osmolality. Approximately 200 mg Sodium Citrate Dihydrate added as a buffer. pH adjusted with sodium bicarbonate and hydrochloric acid. pH 5.5 - 6.8

Each 100 mL also contains approximately 1 mg Edetate Disodium Dihydrate added as a metal chelator.

GALAXY is a registered trademark of Baxter International Inc.

Distributed by  
Wyeth Pharmaceuticals LLC  
A subsidiary of Pfizer Inc  
Philadelphia, PA 19101

PL 2040 Plastic

14715700  
0734001602

**PfizerHospital**

**ZOSYN<sup>®</sup>**  
(Piperacillin and Tazobactam  
Injection)

**4.5 g**

Rx only

**GALAXY**  
Single Dose  
Container

**100 mL**  
**Iso-osmotic**

NDC 0206-8862-01  
**Code 2G3577**  
Sterile Nonpyrogenic

Store at or below -20°C/-4°F. Thaw at room temperature 20° to 25°C (68° to 77°F) or under refrigeration 2° to 8°C (36° to 46°F). **DO NOT FORCE THAW BY IMMERSION IN WATER BATHS OR BY MICROWAVE IRRADIATION.** Thawed solution remains stable for 14 days under refrigeration or 24 hours at room temperature.

**Do not refreeze. Handle frozen product containers with care. Product containers may be fragile in the frozen state.**

**Dosage and Use:** For intravenous use only. As directed by a physician. See Package Insert.

**Cautions:** Do not add supplementary medication or additives. Must not be used in series connections. Check for minute leaks and solution clarity.

Each 100 mL contains: Piperacillin Sodium equivalent to 4 g Piperacillin and Tazobactam Sodium equivalent to 0.5 g Tazobactam in Water for Injection.

Approximately 2 g Dextrose Hydrated added to adjust osmolality. Approximately 200 mg Sodium Citrate Dihydrate added as a buffer. pH adjusted with sodium bicarbonate and hydrochloric acid. pH 5.5 - 6.8

Each 100 mL also contains approximately 1 mg Edetate Disodium Dihydrate added as a metal chelator.


GALAXY is a registered trademark of Baxter International Inc.

Distributed by  
Wyeth Pharmaceuticals LLC  
A subsidiary of Pfizer Inc.  
Philadelphia, PA 19101



PL 2040 Plastic  
14715700  
0734001602

FPO UPC © 100%





## PRINCIPAL DISPLAY PANEL - 4.5 g Container Box Label

### ZOSYN® (Piperacillin and Tazobactam Injection)


Rx only

4.5 g

2 x 6 x 100 mL Single Dose Containers  
Contains 2 boxes of 6 of NDC 0206-8862-01  
Iso-osmotic  
Store at or  
below -20°C/-4°F.  
Do not refreeze.

NDC 0206-8862-02  
Contains 12 of NDC 0206-8862-01  
Code 2G3577

**PfizerHospital**



## ZOSYN IN GALAXY CONTAINERS

tazobactam sodium and piperacillin sodium injection, solution

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0206-8860
<b>Route of Administration</b>	INTRAVENOUS		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>TAZOBACTAM SODIUM</b> (UNII: UXA545ABTT) (TAZ OBACTAM - UNII:SE10G96M8W)	TAZ OBACTAM	250 mg in 50 mL
<b>PIPERACILLIN SODIUM</b> (UNII: M98T69Q7HP) (PIPERACILLIN ANHYDROUS - UNII:9I628532GX)	PIPERACILLIN ANHYDROUS	2 g in 50 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>DEXTROSE MONOHYDRATE</b> (UNII: LX22YL083G)	1 g in 50 mL
<b>SODIUM CITRATE, UNSPECIFIED FORM</b> (UNII: 1Q73Q2JULR)	100 mg in 50 mL
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	0.5 mg in 50 mL
<b>SODIUM BICARBONATE</b> (UNII: 8MDF5V39QO)	
<b>HYDROCHLORIC ACID</b> (UNII: QTT17582CB)	
<b>WATER</b> (UNII: 059QF0KOOR)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0206-8860-02	24 in 1 BOX	03/01/1998	
1	NDC:0206-8860-01	50 mL in 1 BAG; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA050750	03/01/1998	

**ZOSYN IN GALAXY CONTAINERS**

tazobactam sodium and piperacillin sodium injection, solution

**Product Information**

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0206-8861
<b>Route of Administration</b>	INTRAVENOUS		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>TAZOBACTAM SODIUM</b> (UNII: UXA545ABTT) (TAZ OBACTAM - UNII:SE10G96M8W)	TAZ OBACTAM	375 mg in 50 mL
<b>PIPERACILLIN SODIUM</b> (UNII: M98T69Q7HP) (PIPERACILLIN ANHYDROUS -	PIPERACILLIN	2 g in 50 mL

UNII:9I628532GX)

ANHYDROUS

5 g in 50 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>DEXTROSE MONOHYDRATE</b> (UNII: LX22YL083G)	1.5 g in 50 mL
<b>SODIUM CITRATE, UNSPECIFIED FORM</b> (UNII: 1Q73Q2JULR)	150 mg in 50 mL
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	0.75 mg in 50 mL
<b>SODIUM BICARBONATE</b> (UNII: 8MDF5V39QO)	
<b>HYDROCHLORIC ACID</b> (UNII: QTT17582CB)	
<b>WATER</b> (UNII: 059QF0KO0R)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0206-8861-02	24 in 1 BOX	03/01/1998	
1	NDC:0206-8861-01	50 mL in 1 BAG; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA050750	03/01/1998	

**ZOSYN IN GALAXY CONTAINERS**

tazobactam sodium and piperacillin sodium injection, solution

**Product Information**

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0206-8862
<b>Route of Administration</b>	INTRAVENOUS		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>TAZOBACTAM SODIUM</b> (UNII: UXA545ABTT) (TAZOBACTAM - UNII:SE10G96M8W)	TAZOBACTAM	500 mg in 100 mL
<b>PIPERACILLIN SODIUM</b> (UNII: M98T69Q7HP) (PIPERACILLIN ANHYDROUS - UNII:9I628532GX)	PIPERACILLIN ANHYDROUS	4 g in 100 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>DEXTROSE MONOHYDRATE</b> (UNII: LX22YL083G)	2 g in 100 mL

<b>SODIUM CITRATE, UNSPECIFIED FORM</b> (UNII: 1Q73Q2JULR)	200 mg in 100 mL
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	1 mg in 100 mL
<b>SODIUM BICARBONATE</b> (UNII: 8MDF5V39QO)	
<b>HYDROCHLORIC ACID</b> (UNII: QTT17582CB)	
<b>WATER</b> (UNII: 059QF0KO0R)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0206-8862-02	12 in 1 BOX	03/01/1998	
1	NDC:0206-8862-01	100 mL in 1 BAG; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA050750	03/01/1998	

**Labeler** - Wyeth Pharmaceuticals LLC, a subsidiary of Pfizer Inc. (113008515)

### Establishment

Name	Address	ID/FEI	Business Operations
Yuhan Chemical Inc.		687831958	API MANUFACTURE(0206-8860, 0206-8861, 0206-8862)

### Establishment

Name	Address	ID/FEI	Business Operations
Baxter Healthcare Corporation		194684502	ANALYSIS(0206-8860, 0206-8861, 0206-8862) , LABEL(0206-8860, 0206-8861, 0206-8862) , MANUFACTURE(0206-8860, 0206-8861, 0206-8862) , PACK(0206-8860, 0206-8861, 0206-8862)

Revised: 3/2023

Wyeth Pharmaceuticals LLC, a subsidiary of Pfizer Inc.