

## **TOPCARE MILK OF MAGNESIA- magnesium hydroxide liquid**

**Topco Associates LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

### **Topco Associates LLC. Milk of Magnesia Drug Facts**

#### **Active ingredient (in each 15 mL Tablespoon)**

Magnesium hydroxide 1200 mg

#### **Purpose**

Saline laxative

#### **Uses**

- relieves occasional constipation (irregularity)
- generally produces bowel movement in ½ to 6 hours

#### **Warnings**

##### **Ask a doctor before use if you have**

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

##### **Ask a doctor or pharmacist before use if you are**

taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

##### **Stop use and ask a doctor if**

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

##### **If pregnant or breast-feeding,**

ask a health professional before use.

##### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

#### **Directions**

- do not exceed the maximum recommended daily dose in a 24 hour period

- shake well before use
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided
- mL = milliliter; TBSP = Tablespoon

adults and children 12 years and older	30 mL (2 TBSP) to 60 mL (4 TBSP)
children 6 to 11 years	15 mL (1 TBSP) to 30 mL (2 TBSP)
children under 6 years	ask a doctor

### Other information

- each tablespoon (15 mL) contains: calcium 20 mg, magnesium 595 mg, potassium 2 mg and sodium 3 mg
- does not meet USP requirements for preservative effectiveness
- store at 20-25°C (68-77°F)
- close cap tightly after use
- avoid freezing

### Inactive ingredients

anhydrous citric acid, D&C red #28, flavor, glycerin, purified water, sorbitol solution, sucrose, xanthan gum

### Questions or comments?

1-888-423-0139

### Principal Display Panel

STIMULANT FREE

Milk of Magnesia

SALINE LAXATIVE

MAGNESIUM HYDROXIDE 1200 mg

Comfortable, cramp free relief

ALCOHOL 0.4%

CHERRY FLAVOR

COMPARE TO PHILLIPS'® WILD CHERRY active ingredient

12 FL OZ (355 mL)

#1 Doctor recommended saline laxative active ingredient

NDC 36800-949-40

**STIMULANT FREE**

**TopCare<sup>®</sup>**

# Milk of Magnesia

**SALINE LAXATIVE  
MAGNESIUM HYDROXIDE 1200 mg**

- **Comfortable,  
cramp free relief**

**ALCOHOL 0.4%  
CHERRY FLAVOR**



*COMPARE TO PHILLIPS'<sup>®</sup>  
WILD CHERRY  
active ingredient\**

**12 FL OZ  
(355 mL)**

**#1 Doctor  
recommended  
saline laxative  
active ingredient ♦**

: 94940 88 F7

## Drug Facts

**DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING.**

### Active ingredient (in each 15 mL Tablespoon)

### Purpose

Magnesium hydroxide 1200 mg.....Saline laxative

**Uses** ■ relieves occasional constipation (irregularity)  
■ generally produces bowel movement in ½ to 6 hours

### Warnings

**Ask a doctor before use if you have**

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

**Ask a doctor or pharmacist before use if you are taking any other drug.**

Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

**Stop use and ask a doctor if** ■ you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.

- you need to use a laxative for more than 1 week

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### Directions

- do not exceed the maximum recommended daily dose in a 24 hour period
- shake well before use
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided ■ mL = milliliter; TBSP = Tablespoon

adults and children 12 years and older	30 mL (2 TBSP) to 60 mL (4 TBSP)
children 6 to 11 years	15 mL (1 TBSP) to 30 mL (2 TBSP)
children under 6 years	ask a doctor

### Other information

- each tablespoon (15 mL) contains: calcium 20 mg, magnesium 595 mg, potassium 2 mg and sodium 3 mg
- does not meet USP requirements for preservative effectiveness
- store at 20-25°C (68-77°F) ■ close cap tightly after use ■ avoid freezing

**Inactive ingredients** anhydrous citric acid, D&C red #28, flavor, glycerin, purified water, sorbitol solution, sucrose, xanthan gum

**Questions or comments?** 1-888-423-0139

04646 :

\*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Phillips'® Wild Cherry.

◆In the U.S. Source: Symphony Health Solutions.

DISTRIBUTED BY

TORCO ASSOCIATES LLC

**GLUTEN FREE**

38  
B7

TOPCO ASSOCIATES LLC  
 ELK GROVE VILLAGE, IL 60007  
 1-888-423-0139 ©TOPCO  
 PER1114 topcare@topco.com



## TOPCARE MILK OF MAGNESIA

magnesium hydroxide liquid

### Product Information

Product Type	HUMAN OTC DRUG LABEL	Item Code (Source)	NDC:36800-949
Route of Administration	ORAL	DEA Schedule	

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MAGNESIUM HYDROXIDE (MAGNESIUM CATION and HYDROXIDE ION)	MAGNESIUM HYDROXIDE	1200 mg in 15 mL

### Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID	
D&C RED NO. 28	
GLYCERIN	
WATER	
SORBITOL	
SUCROSE	
XANTHAN GUM	

### Product Characteristics

Color	PINK (Opaque Light pink)	Score	
Shape		Size	
Flavor	CHERRY (wild cherry)	Imprint Code	
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-949-40	355 mL in 1 BOTTLE		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	02/20/2006	

