ACID RELIEF- famotidine tablet, film coated Rite Aid Corporation

Rite Aid Corporation Acid Relief Drug Facts

Active ingredient (in each tablet)

Famotidine 20 mg

Purpose

Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating, or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- adults and children 12 years and over:
- to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10
 to 60 minutes before eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture

Inactive ingredients

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc, titanium dioxide

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

FREE FROM | GLUTEN FREE

SOY FREE

Compare to the active ingredient of Maximum Strength Zantac 360°[™]

MAXIMUM STRENGTH

ACID RELIEF

Famotidine Tablets, 20 mg

Acid Reducer

Just one tablet prevents & relieves heartburn due to acid indigestion

ACTUAL SIZE

50 TABLETS



JUST ONE TABLET prevents and relieves heartb um due to acid indigestion brought on by eating and drinking certain foods and beverages. DO NOTUSE IF PRINTED FOILUNDER CAP IS B ROK EN OR MISSING	Tips for Managing Hear foun • D onot lieflat or bend over a fer eating • D onot wear tight-fitting clothing around the stom ach • O onot eat before bedtime • Baise the head of your bed • Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocola te, caffeine, a lochol, and certain fruits and vegetables • Eat slowly and avoid big meals • I foverweight, lose weight	
Drug Facts	Drug Facts (continued)	
Active in gre dient (in each tablet) Purpose Famotidine 20 mg	 heartburn with ligh the adedness, sweating, or dizzinees chest pain or should erp ain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lighthea dedness frequent of hestpain frequent wheezing, particularly with hearburn unexplained weight loss nause a or vomiting stom ach pain kidney disease 	
Warnings Allergyalert: Do not use if you are allergic to fam otidine or other acid reducers	Ask adoctor or pharmacist before use if you a re taking a prescription drug. Acid red ucers may interact with certain prescription drugs.	
Donot use ■ if you have trouble or pain swallowing food, vomiting with blood, orbloody or black stools. The se may be signs of a serious condition. See yourd octor	Stop use a ndask a doctor if yourheartburn continues orwonsens you need to take this product formo re than 14 days If pre gnant or breast-feeding, ask a health	
 with other acid reducers As k adoc tor before use if you have had heartburn over 3 mon ths. This may be a sign of a more serious condition. 	professional before use. Keep out of reach of children. In case of overdose, get me dical help or contact a Poison Con trol Center right away.(1-800-222-1222)	
Drug Facts (continued) Directions adults and children 12 years and over:	Drug Facts (continued) ■ store at 20°-25°C (68°-77°F) ■ protect from moisture	
 to re lie ve symptoms, swall ow 1 tablet with a glass of water. Do not chew. to prevent symptoms, swall ow 1 tablet with a glass of va terat any time from 10 to 60 minutes before eating food or drinking beverages that cause hearburn 	In ac ti ve in gredien ts carnaub a wa x, colloi dal si licon dixxid e, croscarmell ose so diu m, l actose monoh yd rate, magnesium stea rate, microcrysta lline ce llu lose, polyethylene glycol, polyvin yl alcohol, tal c, titani um dixxide	
■ donot use more than 2 tab lets in 24 hours ■ children un der 12 years:ask a doctor	Questions or comments? 1-800-719-9260	
Other information read the directions and warnings before use keep the carton. It contains important information.	*This product is not manu factured or distributed by Chattern, In c, a Sanofi Company, distributor of Maximum Strength Zantac 360 ª™.	

ACID RELIEF

famotidine tablet, film coated

Product Information					
Product Type	HUMAN OTC DRUG	Item Code (Source) NDC:11822		322-6123	
Route of Administration	ORAL				
Active Ingredient/Active	Molety				
Ingredient Name Basis of Strengt			rength	Strength	
FAMOTIDINE (UNII: 5QZ015J2Z8) (FAMOTIDINE - UNII:5QZ015J2Z8) FAMOTIDINE			20 mg		
Inactive Ingredients					
	Ingredient Name				Strength
CARNAUBA WAX (UNII: R12CBM0	EIZ)				
-					
SILICON DIOXIDE (UNII: ETJ7Z6X	BU4)				

LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics	
-------------------------	--

Color	WHITE	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	L194
Contains			

Packaging

#	ltem Code	Package Description	Marketing Start Date	Marketing End Date	
	NDC:11822- 6123-0	1 in 1 CARTON	07/19/2023		
1		50 in 1 BOTTLE; Type 0: Not a Combination Product			
Marketing Information					

Marketing	Application Number or Monograph	Marketing Start	Marketing End
Category	Citation	Date	Date
ANDA	ANDA077351	07/19/2023	

Labeler - Rite Aid Corporation (014578892)

Revised: 8/2023

Rite Aid Corporation