MOTION SICKNESS RELIEF- meclizine hcl tablet L.N.K. International, Inc.

Quality Plus 44-403

Active ingredient (in each tablet)

Meclizine HCl 25 mg

Purpose

Antiemetic

Uses

for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness

Warnings

Do not use

for children under 12 years of age unless directed by a doctor.

Ask a doctor before use if you have

- difficulty in urination due to enlargement of the prostate gland
- glaucoma
- a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers.

When using this product

- drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- take first dose ½ to 1 hour before starting activity
- adults and children 12 years and over: 1 to 2 tablets once daily, or as directed by a doctor

Other information

- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- protect from heat and humidity
- see end flap for expiration date and lot number

Inactive ingredients

colloidal silicon dioxide, D&C yellow #10 aluminum lake, lactose monohydrate, magnesium stearate, pregelatinized starch

Questions or comments?

1-800-426-9391

Principal Display Panel

QUALITY +PLUS

NDC 50844-403-19

*Compare to active ingredient in Dramamine® Less Drowsy

Motion Sickness Relief LESS DROWSY FORMULA Meclizine HCl 25 mg • Antiemetic

HELPS PREVENT NAUSEA, VOMITING AND DIZZINESS DUE TO MOTION SICKNESS FOR UP TO **24 HOURS**

Ages 12 Years and Over

8Tablets

ACTUAL SIZE

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS

ANY SIGNS OF TAMPERING

*This product is not manufactured or distributed by Medtech Products Inc., owner of the registered trademark

Dramamine® All Day Less Drowsy. 50844 ORG042340319

Distributed by **LNK INTERNATIONAL, INC.** 60 Arkay Drive Hauppauge, NY 11788 USA





Motion Sickness Relief LESS DROWSY FORMULA

QUALITY PLUS

NDC 50844-403-19

*Compare to active ingredient in Dramamine® Less Drowsy

Motion Sickness Relief LESS DROWSY FORMULA

Meclizine HCl 25 mg • Antiemetic



HELPS PREVENT NAUSEA, VOMITING AND DIZZINESS DUE TO MOTION SICKNESS FOR UP TO 24 HOURS

8 Tablets

Motion Sick Relief CESS DROWS

Ages 12 Years and Over

ACTUAL



Hauppauge, NY 11788 60 Arkay Drive LNK INTERNATIONAL, INC. Distubuted by

ORG042340319 20844 trademark Dramamine® Less Drowsy. *This product is not manufactured or distributed by Medtech Products Inc., owner of the registered

Questions or comments? 1-800-426-9391

specials, pregelatinized starch yellow #10 aluminum lake, lactose monohydrate, magnesium Inactive ingredients colloidalsificon dioxide, D&C

- see end filsp for expiration date and lot number
 - brosect from heat and humidity

- 12°-30°C (59°-86°F) ■ store at 25°C (77°F); excursions permit led between
- OPENED OR BLISTER IS TORN OR BROKEN

■ TAMPER EVIDENT: DO NOT USE IF OUTER PACK AGE IS попътпоти тяпти

sqn jts sug cyrjquen 15 hears sug over: 1 to 5 tapjets once A MUNICIPA PRINCIPA PRINCIPA INCHI I CIR ZZ GONO YE III GAVEN

daily, or as directed by a doctor

- пае сапрои муви димид в шорт мвриде от обегатий **Growsiness**
 - sicohol, sedatives, and tranquitizers may increase
 - safeuanaq orjectorie proverages
 - quowajuesa jugit occnt

When using this product

sedatives or tranquifizers.

Ask a doctor or pharmacist before use if you are taking


- s preatring problem such as emphysema or chronic
- gland **a** glaucoma quincuity in unnation due to enlargement of the prostate Ask a doctor before use if you have

directed by a doctor.

Do not use for children under 12 years of age unless sgnimsW

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

B-1603-403A-19-R 0RG042340319



MOTION SICKNESS RELIEF

meclizine hcl tablet

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:50844-403

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name

MECLIZINE HYDROCHLORIDE (UNII: HDP7W44CIO) (MECLIZINE - UNII:3L5TQ84570)

Basis of Strength

MECLIZINE HYDROCHLORIDE

MECLIZINE HYDROCHLORIDE

25 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	

LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)
MAGNESIUM STEARATE (UNII: 70097M6I30)

STARCH, CORN (UNII: O8232NY3SJ)

	haract		

Color	yellow	Score	no score
Shape	ROUND	Size	9mm
Flavor		Imprint Code	44;403
Contains			

Pa	ck	aa	ing
	٠.,	~9	9

Warketing Start Marketing End

4	item cou	е гаскаде рессприон	Date	Date
]	NDC:50844-40	1 in 1 CARTON	06/24/2002	
]	L	8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
OTC Monograph Drug	M009	06/24/2002			
3 1 3					

Labeler - L.N.K. International, Inc. (038154464)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(50844-403)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(50844-403) , pack(50844-403)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(50844-403)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(50844-403)

Revised: 7/2023 L.N.K. International, Inc.