

STOOL SOFTENER- marc glassman capsule
Marc Glassman, Inc.

420-Stool Softener

Active ingredient(s)

Docusate sodium 100mg

Purpose

Stool softener

Use(s)

- relieves occasional constipation (irregularity)
- generally produces bowel movement in 12 to 72 hours

Warnings

Do not use

- if you are presently taking mineral oil, unless told to do so by a doctor

Ask a doctor before use if

- stomach pain
- nausea
- vomiting
- noticed a sudden change in bowel habits that lasts over a period of 2 weeks

Stop use and ask a doctor if

- you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.
- you need to use a stool softener laxative for more than 1 week

Pregnancy/Breastfeeding

ask a health professional before use.

Keep out of reach of children

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

Take only by mouth. Doses may be taken as a single daily dose or in divided doses.

Adults and children 12 years and over: take 1-3 capsules daily

Children 2 to under 12 years of age: take 1 capsule daily

Children under 2 years of age: ask a doctor

Other information

- each capsule contains: sodium 5mg
- VERY LOW SODIUM

Storage

store at 25°C (77°F) excursions permitted between 15°- 30°C (59°-86°F)

Inactive ingredients

black edible ink, citric acid, D&C red #33, FD&C blue #1, FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol, propylene glycol, purified water, sorbitol special

Questions

Call 1-888-952-0050 Monday through Friday 9am-5pm

Principal Display Panel

Marc's NDC 68998-420-01
 †Compare to the active ingredient in Colace®

Docosate Sodium

STOOL SOFTENER



100 CAPSULES - 100 MG EACH

- Gentle
- Stimulant Free
- Effective

TAMPER EVIDENT: DO NOT USE IF SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active Ingredient (in each capsule)
 Docosate sodium 100mg..... Stool softener

Uses
 ■ relieves occasional constipation (irregularity)
 ■ generally produces bowel movement in 12 to 72 hours

Warnings
Do not use ■ if you are presently taking mineral oil, unless told to do so by a doctor
Ask a doctor before use if you have ■ stomach pain ■ nausea ■ vomiting ■ noticed a sudden change in bowel habits that lasts over a period of 2 weeks
Stop use and ask a doctor if ■ you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition. ■ you need to use a stool softener laxative for more than 1 week

Continued On Back Of Label
 †This product is not manufactured or distributed by Purdue Products L.P., owner of the registered trademark Colace®.
 Distributed By:
 Marc Glassman, Inc.
 West 130th Street
 Cleveland, OH 44130



181414

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children.
 In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions
 ■ Take only by mouth. Doses may be taken as a single daily dose or in divided doses.
Adults and children 12 years and older take 1-3 capsules daily
Children 2 to under 12 years of age take 1 capsule daily
Children under 2 years of age ask a doctor

Other Information
 ■ each capsule contains sodium 5mg
 ■ VERY LOW SODIUM
 ■ store at 25°C (77°F); excursions permitted between 15° - 30°C (59° - 86°F)

Inactive ingredients
 black edible ink, citric acid, D&G red #33, FD&C blue #1, FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol, propylene glycol, purified water, sorbitol special

Stool Softener

STOOL SOFTENER

marc glassman capsule

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:68998-420 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------|----------|
| DOCUSATE SODIUM (UNII: F05Q2T2JA0) (DOCUSATE - UNII:M7P27195AG) | DOCUSATE SODIUM | 100 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) | |
| D&C RED NO. 33 (UNII: 9DBA0SBB0L) | |
| FD&C BLUE NO. 1 (UNII: H3R47K3TBD) | |
| FD&C RED NO. 40 (UNII: WZB9127XOA) | |
| FD&C YELLOW NO. 6 (UNII: H77VEI93A8) | |
| GELATIN (UNII: 2G86QN327L) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| SORBITOL SOLUTION (UNII: 8KW3E207O2) | |
| WATER (UNII: 059QF0KO0R) | |

Product Characteristics

| | | | |
|-----------------|----------------|---------------------|----------|
| Color | red, white | Score | no score |
| Shape | CAPSULE (oval) | Size | 13mm |
| Flavor | | Imprint Code | PC18 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:68998-420-01 | 100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 09/04/2019 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M007 | 04/10/2015 | |

Labeler - Marc Glassman, Inc. (094487477)