

## **MILK OF MAGNESIA- magnesium hydroxide liquid**

### **H E B**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

### **HEB Milk of Magnesia Drug Facts**

#### **Active ingredient (in each 15 mL Tablespoon)**

Magnesium hydroxide 1200 mg

#### **Purpose**

Saline laxative

#### **Uses**

- relieves occasional constipation (irregularity)
- generally produces bowel movement in ½ to 6 hours

#### **Warnings**

##### **Ask a doctor before use if you have**

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

##### **Ask a doctor or pharmacist before use if you are**

taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

##### **Stop use and ask a doctor if**

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

##### **If pregnant or breast-feeding,**

ask a health professional before use.

##### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

#### **Directions**

- do not exceed the maximum recommended daily dose in a 24 hour period

- shake well before use
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided
- mL = milliliter; TBSP = Tablespoon

adults and children 12 years and older	30 mL (2 TBSP) to 60 mL (4 TBSP)
children 6 to 11 years	15 mL (1 TBSP) to 30 mL (2 TBSP)
children under 6 years	ask a doctor

### Other information

- each tablespoon (15 mL) contains: calcium 20 mg, magnesium 595 mg, potassium 2 mg and sodium 3 mg
- does not meet USP requirements for preservative effectiveness
- store at 20-25°C (68-77°F)
- close cap tightly after use
- avoid freezing

### Inactive ingredients

anhydrous citric acid, D&C red #28, flavor, glycerin, purified water, sorbitol solution, sucrose, xanthan gum

### Questions or comments?

1-800-719-9260

### Principal Display Panel

Compare to Phillips'® Milk of Magnesia Wild Cherry active ingredient

MILK OF MAGNESIA

Magnesium Hydroxide

Saline Laxative

Comfortable, Cramp Free Relief

Stimulant Free

Wild Cherry Flavor

#1 DOCTOR RECOMMENDED saline laxative active ingredient

ALCOHOL 0.4%

GLUTEN FREE

12 FL OZ (355 mL)

Compare to Phillips'®  
Milk of Magnesia Wild Cherry  
active ingredient\*

NDC 37808-949-40



**MILK OF MAGNESIA**

**Magnesium Hydroxide**

**Saline Laxative**

- Comfortable, Cramp Free Relief
- Stimulant Free

**Wild Cherry Flavor**

**ALCOHOL 0.4%**

**GLUTEN FREE**

12 FL OZ (355 mL)

: 94940 1J F3



## Drug Facts

**DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING.**

### Active ingredient (in each 15 mL Tablespoon)

Magnesium hydroxide 1200 mg.....Saline laxative

### Purpose

### Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in ½ to 6 hours

### Warnings

**Ask a doctor before use if you have**

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

**Ask a doctor or pharmacist before use if you are taking any other drug.**

Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

**Stop use and ask a doctor if** ■ you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.

- you need to use a laxative for more than 1 week

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### Directions

- do not exceed the maximum recommended daily dose in a 24 hour period
- shake well before use
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided
- mL = milliliter; TBSP = Tablespoon

adults and children 12 years and older	30 mL (2 TBSP) to 60 mL (4 TBSP)
children 6 to 11 years	15 mL (1 TBSP) to 30 mL (2 TBSP)
children under 6 years	ask a doctor

### Other information

- **each tablespoon (15 mL) contains:** calcium 20 mg, magnesium 595 mg, potassium 2 mg and sodium 3 mg
- does not meet USP requirements for preservative effectiveness
- store at 20-25°C (68-77°F)
- close cap tightly after use
- avoid freezing

**Inactive ingredients** anhydrous citric acid, D&C red #28, flavor, glycerin, purified water, sorbitol solution, sucrose, xanthan gum

**Questions or comments? 1-800-719-9260**

\*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Phillips'® Milk of Magnesia Wild Cherry.

\*\*In the U.S. Source: Symphony Health Solutions.

1410

MADE WITH PRIDE & CARE FOR H-E-B®

1  
2526  
02217  
0

SAN ANTONIO, TX 78204  
100% GUARANTEE PROMISE:  
If you aren't completely pleased with  
this product, we'll be happy to replace  
it or refund your money.  
You have our word on it.

1  
2526  
02217  
0

MILK OF MAGNESIA

magnesium hydroxide liquid

Product Information			
Product Type	HUMAN OTC DRUG LABEL	Item Code (Source)	NDC:37808-949
Route of Administration	ORAL	DEA Schedule	

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
MAGNESIUM HYDRO XIDE (MAGNESIUM CATION and HYDROXIDE ION)	MAGNESIUM HYDROXIDE	1200 mg in 15 mL

Inactive Ingredients	
Ingredient Name	Strength
ANHYDROUS CITRIC ACID	
GLYCERIN	
WATER	
SUCROSE	
XANTHAN GUM	
D&C RED NO. 28	
SORBITOL	

Product Characteristics			
Color	PINK (Opaque Light pink)	Score	
Shape		Size	
Flavor	CHERRY (wild cherry)	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37808-949-40	355 mL in 1 BOTTLE		
2	NDC:37808-949-35	796 mL in 1 BOTTLE		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	10/10/2012	

**Labeler** - HEB (007924756)

Revised: 2/2015

HEB