MILK OF MAGNESIA- magnesium hydroxide liquid H E B

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

HEB Milk of Magnesia Drug Facts

Active ingredient (in each 15 mL TBSP)

Magnesium hydroxide 1200 mg

Purpose

Saline laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in ½ to 6 hours

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

Ask a doctor or pharmacist before use if you are

taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

shake well before use

- do not exceed the maximum recommended daily dose in a 24 hour period
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided
- TBSP = tablespoonful

adults and children 12 years and older	2 to 4 TBSP.
children 6 to 11 years	1 to 2 TBSP.
children under 6 years	ask a doctor

Other information

- each tablespoon (15 mL) contains: calcium 20 mg, magnesium 595 mg, potassium 2 mg and sodium 2 mg
- does not meet USP requirements for preservative effectiveness
- store at 20-25°C (68-77°F)
- keep tightly closed

Inactive ingredients

anhydrous citric acid, D&C red #28, flavor, glycerin, purified water, sorbitol solution, sucrose, xanthan gum

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to Phillips' ® Milk of Magnesia Wild Cherry active ingredient

MILK OF MAGNESIA

Magnesium Hydroxide

Saline Laxative

Comfortable, Cramp Free Relief

Stimulant Free

Wild Cherry Flavor

ALCOHOL 0.4%

GLUTEN FREE

Compare to Phillips'® Milk of Magnesia Wild Cherry active ingredient*

NDC 37808-949-40



MILK OF MAGNESIA

Magnesium Hydroxide


Saline Laxative

- Comfortable, Cramp
 Free Relief
- Stimulant Free

Wild Cherry Flavor

ALCOHOL 0.4%

GLUTEN FREE



12 FL OZ (355 mL)

: 94940 1J F2

Drug Facts

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING.

Active ingredient (in each 15 mL TBSP)

Purpose

Magnesium hydroxide 1200 mg......Saline laxative

- **Uses** relieves occasional constipation (irregularity)
- generally produces bowel movement in 1/2 to 6 hours

Warnings

Ask a doctor before use if you have

- kidney disease
 a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- shake well before use
- do not exceed the maximum recommended daily dose in a 24 hour period
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
 drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided
- TBSP = tablespoonful

adults and children 12 years and older	2 to 4 TBSP.	
children 6 to 11 years	1 to 2 TBSP.	
children under 6 years	ask a doctor	

Other information

■ each tablespoon (15 mL) contains: calcium 20 mg, magnesium 595 mg, potassium 2 mg and sodium 2 mg
 ■ store at 20-25°C (68-77°F)
 ■ keep tightly closed
 ■ does not meet USP requirements for preservative effectiveness

Inactive ingredients anhydrous citric acid, D&C red #28, flavor, glycerin, purified water, sorbitol solution, sucrose, xanthan gum

(Questions or comments? 1-800-719-9260

THIS PRODUCT IS NOT MAINTACTURED or distributed by Bayer HealthCare LLC, distributor of Phillips'® Milk of Magnesia Wild Cherry.

MADE WITH PRIDE & CARE FOR H-E-B SAN ANTONIO, TX 78204



MILK OF MAGNESIA

magnesium hydroxide liquid

Product Information

Product Type HUMAN OTC DRUG LABEL NDC:37808-949 Item Code (Source) ORAL **Route of Administration DEA Schedule**

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength MAGNESIUM HYDRO XIDE (MAGNESIUM CATION and HYDRO XIDE ION) MAGNESIUM CATION $1200\ mg$ in $15\ mL$

Inactive Ingredients

Ingredient Name	Strength
ANHYDRO US CITRIC ACID	
GLYCERIN	
WATER	
SUCROSE	
XANTHAN GUM	
D&C RED NO. 28	
SORBITOL	

Product Characteristics

1 Toutet Omit determines		
Color	PINK (Opaque Light pink)	Score
Shape		Size
Flavor	CHERRY (wild cherry)	Imprint Code
Contains		

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37808-949-40	355 mL in 1 BOTTLE		
2	NDC:37808-949-35	796 mL in 1 BOTTLE		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	10/10/2012	

Labeler - HEB (007924756)

Revised: 1/2013 HE B