

**FOSTER AND THRIVE ANTI DIARRHEAL ANTI GAS- loperamide hydrochloride, simethicone tablet**  
**Strategic Sourcing Services LLC**

-----  
**McKesson Anti-Diarrheal Anti-Gas Drug Facts**

**Active ingredients (in each caplet)**

Loperamide HCl 2 mg

Simethicone 125 mg

**Purposes**

Anti-diarrheal

Anti-gas

**Uses**

relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas

**Warnings**

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

**Heart alert:** Taking more than directed can cause serious heart problems or death

**Do not use**

- if you have bloody or black stool
- if you have difficulty swallowing

**Ask a doctor before use if you have**

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug. Loperamide may interact with certain prescription drugs.

**When using this product**

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

## Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.
- you have difficulty swallowing the caplet

## If pregnant or breast-feeding,

ask a health professional before use.

## Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- take only on an empty stomach (1 hour before or 2 hours after a meal)
- take with a full (8 oz.) glass of water
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

## Other information

- **each caplet contains:** calcium 165 mg, sodium 3 mg
- store between 20°-25°C (68°-77°F). Protect from light.
- **Do not use if printed blister unit is broken or torn**

## Inactive ingredients

acesulfame potassium, croscarmellose sodium, dibasic calcium phosphate, maltodextrin, microcrystalline cellulose, natural and artificial flavor, propylene glycol, stearic acid

## Questions or comments?

Call **833-358-6431** Monday to Friday 9:00am to 7:00pm EST

## Package/Label Principal Display Panel

COMPARE TO IMODIUM® MULTI-SYMPTOM RELIEF ACTIVE INGREDIENTS

Foster & Thrive™

MULTI-SYMPTOM RELIEF CAPLET

Loperamide Hydrochloride and Simethicone tablets, 2 mg/125mg

Anti-Diarrheal

Anti-Gas

ACTUAL SIZE

Relieves symptoms of diarrhea *plus*:

- Cramps & Pressure
- Bloating • Gas

12 CAPLETS\*

\*CAPSULE-SHAPED TABLETS

Foster & Thrive™

MULTI-SYMPTOM RELIEF CAPLET

Loperamide Hydrochloride and Simethicone tablets, 2 mg/125 mg

Anti-Diarrheal/Anti-Gas

NDC 70677-1105-1


COMPARE TO IMODIUM®  
MULTI-SYMPTOM RELIEF  
ACTIVE INGREDIENTS\*\*

Foster & Thrive™

MULTI-SYMPTOM RELIEF CAPLET

Loperamide Hydrochloride and  
Simethicone tablets, 2 mg/125 mg

Anti-Diarrheal  
Anti-Gas



ACTUAL SIZE

Relieves symptoms of diarrhea plus:

- Cramps & Pressure
- Bloating & Gas

12 CAPLETS\*

\*CAPSULE-SHAPED TABLETS

OR 153 KM C-1

### Drug Facts

**Active ingredients (in each caplet)** **Purposes**  
Loperamide HCl 2 mg..... Anti-diarrheal  
Simethicone 125 mg..... Anti-gas

**Uses** relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas

### Warnings

**Allergic alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

**Heart alert:** Taking more than directed can cause serious heart problems or death

### Do not use

- if you have bloody or black stool
- if you have difficulty swallowing

### Ask a doctor before use if you have

- fever ■ mucus in the stool ■ a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

### Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.
- you have difficulty swallowing the caplet

### Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### Directions

- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- take only on an empty stomach (1 hour before or 2 hours after a meal)
- take with a full (8 oz.) glass of water
- find right to dose on chart below. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (40-59 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

### Other information

- each caplet contains: calcium 165 mg sodium 3 mg
- store between 20°-25°C (68°-77°F). Protect from light.
- do not use if printed blister unit is broken or torn

\*\*This product is not manufactured or distributed by  
Johnson & Johnson Consumer Inc., distributor of  
Imodium® Multi-Symptom Relief.

DO NOT USE IF PRINTED BLISTER UNIT IS BROKEN OR TORN

KEEP OUT OF REACH OF CHILDREN. READ WARNING LABEL AND PRODUCT INFORMATION

©2023 McKesson Corporation


Distributed by: McKesson Corp.,  
via Strategic Sourcing Services LLC.

Memphis, TN 38141

Money Back Guarantee

www.fosterandthrive.com

REV. 03/23



## FOSTER AND THRIVE ANTI DIARRHEAL ANTI GAS

loperamide hydrochloride, simethicone tablet

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70677-1105
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>LOPERAMIDE HYDROCHLORIDE</b> (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	125 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>ACESULFAME POTASSIUM</b> (UNII: 23OV73Q5G9)	
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>ANHYDROUS DIBASIC CALCIUM PHOSPHATE</b> (UNII: L11K75P92J)	
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	

### Product Characteristics

Color	WHITE	Score	2 pieces
Shape	OVAL	Size	17mm
Flavor		Imprint Code	LOR1;2;125
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70677-1105-1	12 in 1 CARTON	10/31/2023	

1	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
<b>Marketing Information</b>			
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANDA	ANDA209837	10/31/2023	

**Labeler** - Strategic Sourcing Services LLC (116956644)

Revised: 10/2023

Strategic Sourcing Services LLC