

RANITIDINE - ACID REDUCER- ranitidine hydrochloride tablet, film coated
Chain Drug Consortium, LLC.

Drug Facts

ACTIVE INGREDIENT (IN EACH TABLET)

Ranitidine 75 mg (as ranitidine hydrochloride USP, 84 mg)

PURPOSE

Acid reducer

USES

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

WARNINGS

Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

DIRECTIONS

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water
 - to **prevent** symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn
 - can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor

OTHER INFORMATION

- **TAMPER EVIDENT: DO NOT USE IF THE CARTON OR PRINTED FOIL UNDER CAP IS OPEN OR TORN.**
- store at 20° - 25° C (68° - 77° F)
- avoid excessive heat or humidity
- this product is sodium and sugar free

INACTIVE INGREDIENTS

colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, talc, titanium dioxide

QUESTIONS?

call **1-800-406-7984**

Read the directions, consumer information leaflet and warnings before use.

Keep the carton. It contains important information.

DISTRIBUTED BY

CHAIN DRUG CONSORTIUM

3301 NW BOCA RATON BLVD

SUITE 101, BOCA RATON, FL 33431

PRINCIPAL DISPLAY PANEL

Premier Value[®]

NDC 68016-352-60

Ranitidine Tablets, USP 75 mg

Acid Reducer

Regular Strength

60 Tablets

Prevents & Relieves Heartburn Associated with Acid Indigestion & Sour Stomach

†Compare to the active ingredient of Zantac 75[®]

†This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals Inc., owner of the registered trademark Zantac 75[®].

Drug Facts
Active ingredient (in each tablet)
 Ranitidine 75 mg (as ranitidine hydrochloride USP 84 mg).....Acid reducer

Uses
 ■ relieves heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages
 ■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings
 Do not use if you are allergic to ranitidine or other acid reducers
 ■ If you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools, these may be signs of a serious condition. See your doctor.
 ■ with other acid reducers

NDC 60-81098-00

†Compare to the active ingredient of Zantac 75®

Regular Strength



Ranitidine Tablets, USP 75 mg

Acid Reducer

Prevents & Relieves Heartburn Associated with Acid Indigestion & Sour Stomach




60 Tablets

Read the directions, consumer information leaflet and warnings before use. Keep the carton. It contains important information.

Expiration Date:

Batch No.

NON VARNISH

Regular Strength



Ranitidine Tablets, USP 75 mg

Acid Reducer

60 Tablets

Stop use and ask a doctor if ■ your heartburn continues or worsens ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
 ■ adults and children 12 years and over:
 ■ to relieve symptoms, swallow 1 tablet with a glass of water
 ■ to prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating
 ■ food or drinking beverages that cause heartburn
 ■ can be used up to twice daily (do not take more than 2 tablets in 24 hours)

Other information
 ■ TAMPER EVIDENT: DO NOT USE IF THE CARTON OR PRINTED FOIL UNDER CAP IS OPEN OR TORN.
 ■ store at 20° - 25° C (68° - 77° F)
 ■ avoid excessive heat or humidity
 ■ this product is sodium and sugar free

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, talc, titanium dioxide

Questions? Call 1-800-406-7984

5094939



8 409861 01079 8

†This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals Inc., owner of the registered trademark Zantac 75®.




If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.

DISTRIBUTED BY
 CHAIN DRUG CONSORTIUM
 3301 NW BOCA RATON BLVD
 SUITE 101, BOCA RATON, FL 33431
 MADE IN INDIA

0412



5094939



RANITIDINE - ACID REDUCER

ranitidine hydrochloride tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68016-352
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
RANITIDINE HYDROCHLORIDE (UNII: BK76465IHM) (RANITIDINE - UNII:884KT10YB7)	RANITIDINE	75 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	pink	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	OR;606
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68016-352-60	60 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:68016-352-30	30 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA201745	07/10/2012	

Labeler - Chain Drug Consortium, LLC. (101668460)

Registrant - Ohm Laboratories Inc. (184769029)

Establishment

Name	Address	ID/FEI	Business Operations
Shasun Pharmaceuticals Limited		915786829	MANUFACTURE(68016-352)

Revised: 10/2015

Chain Drug Consortium, LLC.