# COLD AND FLU SEVERE- acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl tablet, film coated Walgreen Company

-----

Walgreens 44-640

# Active ingredients (in each caplet)

Acetaminophen 325 mg Dextromethorphan HBr 10 mg Guaifenesin 200 mg Phenylephrine HCl 5 mg

# **Purpose**

Pain reliever/fever reducer Cough suppressant Expectorant Nasal decongestant

#### Uses

- temporarily relieves common cold and flu symptoms:
  - minor aches and pains
  - sinus congestion and pressure
  - nasal congestion
  - sore throat
  - fever
  - headache
  - cough due to minor throat and bronchial irritation
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- reduces swelling of nasal passages

# Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

blisters

- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

# Ask a doctor before use if you have

- liver disease
- thyroid disease
- diabetes
- heart disease
- high blood pressure
- difficulty in urination due to enlargement of the prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

# Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

# When using this product

do not exceed recommended dosage.

# Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs
  of a serious condition.

# If pregnant or breast-feeding,

ask a health professional before use.

# Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### **Directions**

- do not take more than directed
- do not take more than 8 caplets in 24 hours
- adults and children 12 years and over: take 2 caplets with water every 4 hours
- children under 12 years: ask a doctor

#### Other information

- each caplet contains: sodium 3 mg
- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

# Inactive ingredients

corn starch, crospovidone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polysorbate 80, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

## Questions or comments?

1-800-426-9391

# **Principal Display Panel**

# Walgreens

Compare to the active ingredients in Vicks® DayQuil® Severe Cold & Flu<sup>††</sup>
• WALGREENS • PHARMACIST RECOMMENDED<sup>†</sup>

NDC 0363-0640-08

#### DAY

Severe Cold & Flu

**ACETAMINOPHEN 325 mg** / PAIN RELIEVER / FEVER REDUCER DEXTROMETHORPHAN HBr 10 mg / COUGH SUPPRESSANT GUAIFENESIN 200 mg / EXPECTORANT PHENYLEPHRINE HCI 5 mg / NASAL DECONGESTANT

# **Maximum Strength**

 Relieves headache, fever, sore throat, minor aches & pains, chest congestion, thins & loosens mucus, nasal congestion, sinus pressure & cough

### 24

**CAPLETS** 

**ACTUAL SIZE** 

TAMPER EVIDENT:
DO NOT USE IF
PACKAGE IS OPENED OR IF
BLISTER UNIT IS
TORN, BROKEN
OR SHOWS ANY SIGNS
OF TAMPERING

†Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.
††This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademark Vicks® DayQuil® Severe Cold & Flu.

50844 REV0722A64008

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com
© 2021 Walgreen Co.

#### **PARENTS:**

Learn about teen medicine abuse www.StopMedicineAbuse.org



# Walgreens 44-640

# **COLD AND FLU SEVERE**

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl tablet, film coated

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0640	
Route of Administration	ORAL			

Active Ingredient/Active Moiety					
Ingredient Name	Basis of Strength	Strength			
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg			
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg			
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg			
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg			

# **Inactive Ingredients**

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSPOVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYSORBATE 80 (UNII: 60ZP39ZG8H)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
	_

Product Characteristics				
Color	orange	Score	no score	
Shape	OVAL	Size	19mm	
Flavor		Imprint Code	44;640	
Contains				

Packaging					
# Item Code	Package Description	Marketing Start Date	Marketing End Date		
NDC:0363- 0640-08	2 in 1 CARTON	02/27/2014			
1	12 in 1 BLISTER PACK; Type 0: Not a Combination Product				

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M012	02/27/2014		

# Labeler - Walgreen Company (008965063)

<b>Establishment</b>			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(0363-0640) , pack(0363-0640)

# **Establishment**

Name	Address	ID/FEI	<b>Business Operations</b>
LNK International, Inc.		832867894	manufacture(0363-0640)

Establishment			
Name	Address	ID/FEI	<b>Business Operations</b>
LNK International, Inc.		117025878	manufacture(0363-0640)

Revised: 1/2024 Walgreen Company