

POLLENS - GRASSES, BROME, SMOOTH BROMUS INERMIS- brome, smooth bromus inermis injection, solution
POLLENS - GRASSES, CORN, CULTIVATED ZEA MAYS- corn, cultivated zea mays injection, solution
POLLENS - GRASSES, JOHNSON GRASS SORGHUM HALEPENSE- johnson grass sorghum halepense injection, solution
POLLENS - GRASSES, OATS, COMMON, CULTIVATED AVENA SATIVA- oats, common, cultivated avena sativa injection, solution
POLLENS - GRASSES, SOUTHERN GRASS MIX- pollens - grasses, southern grass mix injection, solution
POLLENS - TREES, ACACIA ACACIA LONGIFOLIA- acacia longifolia injection, solution
POLLENS - TREES, ALDER, RED ALNUS RUBRA- alder, red alnus rubra injection, solution
POLLENS - TREES, ASH, WHITE FRAXINUS AMERICANA- ash, white fraxinus americana injection, solution
POLLENS - TREES, BEECH, AMERICAN FAGUS GRANDIFOLIA- beech, american fagus grandifolia injection, solution
POLLENS - TREES, BIRCH MIX- birch mix injection, solution
POLLENS - TREES, BOTTLEBRUSH, CALLISTEMON SPP.- bottlebrush, callistemon citrinus injection, solution
POLLENS - TREES, BOXELDER/MAPLE MIX- boxelder/maple mix injection, solution
POLLENS - TREES, CEDAR, MOUNTAIN JUNIPERUS ASHEI- cedar, mountain juniperus ashei injection, solution
POLLENS - TREES, CEDAR, RED JUNIPERUS VIRGINIANA- cedar, red juniperus virginiana injection, solution
Jubilant HollisterStier LLC

INSTRUCTIONS ALLERGENIC EXTRACTS FOR SCRATCH, PRICK OR PUNCTURE TESTING

WARNINGS

This product is intended for use only by licensed medical personnel experienced in administering allergenic extracts and trained to provide immediate emergency treatment in the event of a life-threatening reaction. Allergenic extracts may potentially elicit a severe life threatening systemic reaction, rarely resulting in death.⁷ Therefore, emergency measures and personnel trained in their use must be available immediately in the event of such a reaction. Patients should be instructed to recognize adverse reaction symptoms and cautioned to contact the physician's office if symptoms occur. Patients on non-selective beta blockers may be more reactive to allergens given for testing or treatment and may be unresponsive to the usual doses of epinephrine used to treat allergic reactions. This product should never be injected intravenously. Refer also to the WARNINGS, PRECAUTIONS, and ADVERSE REACTIONS Sections for further discussion.

DESCRIPTION

Sterile extracts for scratch, prick or puncture testing are supplied in dropper vials containing, in addition to the extract allergens and antigens, 50% (v/v) glycerin as preservative, 0.5% sodium chloride and 0.275% sodium bicarbonate. The strength of these extracts may be expressed in terms of

- Weight to Volume (w/v)
- Protein Nitrogen Units/mL (PNU/mL)
- Amb a 1 Units/mL (Amb a 1/mL)
- Allergy Units/mL (AU/mL)

- Bioequivalent Allergy Units/mL (BAU/mL)
- Concentrate

1. Weight to volume (w/v).

For regular extracts this describes the extraction ratio, i.e., the amount of crude allergen added to the extracting fluid. A 1:10 extract, therefore, indicates that the solution contains the extracted material from one gram of raw material added to each 10 mL of extracting fluid. The amount and composition of extracted materials will vary with the kind of antigen, the extracting fluid, duration of extraction, pH, temperature, and other variables. APTM (acetone precipitated) extracts, if present, are prepared by reconstituting dry, allergenically active concentrates produced by precipitation process from extracts of raw materials. For those APTM extracts labeled on a weight per volume (w/v) basis, the strength designation indicates the dry weight of finished (acetone) precipitate per volume of reconstituting fluid. For example, 1:50 (w/v) means that each gram of dry precipitate obtained from the original extract is reconstituted in 50 mL of solution.

2. Protein Nitrogen Units per mL (PNU/mL).

One protein nitrogen unit represents 0.00001 mg phosphotungstic acid-precipitable protein nitrogen dissolved in one mL of antigen extract. The PNU content of extracts of the same antigen may vary according to the method of measuring the PNU. Thus, the PNU content of extracts from different manufacturers is not comparable unless the PNU method is known to be the same and reproducible from lot to lot. Also, the amount of protein nitrogen extracted from an antigen is influenced by the same variables as the weight to volume extract. Allergenic materials make up a variable proportion of the total protein of an extract.

3. Amb a 1. Of the many allergens which have been purified and characterized from Short Ragweed (Amb a 1¹², Amb a 2¹³, Ra3¹⁴, Ra4(BPA-R)¹⁵, Ra5¹⁶, Ra6, Ra7, and Ra8¹⁷, and cytochrome C¹⁸), Amb a 1 (also known as Antigen E) is considered the most important and has been selected as the basis for standardization. Extracts of Short Ragweed containing Amb a 1 are diffused in agar against standard anti-serum to Amb a 1, and compared to the diffusion of standard Amb a 1 solutions. The amount of Amb a 1 is expressed as units of Amb a 1 per mL of extract.

If an extract is diluted with a diluent or other allergenic extracts, the Amb a 1 concentration must be determined by calculation.

4. Allergy Units per mL (AU/mL).

The potency of extracts labeled in Allergy Units per mL (AU/mL) is determined by in vitro comparison to a reference standard established by the Center for Biologics Evaluation and Research (CBER) of the Food and Drug Administration (FDA).

5. Bioequivalent Allergy Units per mL (BAU/mL).

When originally licensed, the Reference Preparations for standardized extracts were arbitrarily assigned 100,000 Allergy Units (AU)/mL. Subsequently, quantitative skin testing by the ID₅₀EAL method 13 was used to determine that some Reference Preparations should be assigned 10,000 AU/mL, and others 100,000 AU/mL. To avoid possible confusion about this change in the method of allergy unit assignment, the nomenclature changed for standardized extracts whose allergy units are assigned based on quantitative skin testing, and such products are labeled in Bioequivalent Allergy Units (BAU)/mL. References labeled 10,000 BAU/mL can be diluted one to a half million fold, and references labeled 100,000 BAU/mL can be diluted one to 5 million fold and produce a sum of erythema diameter of 50 mm when Intradermal testing highly reactive subjects.

6. Concentrate.

Concentrate label terminology applies to allergenic extract mixtures, where the individual allergens being combined vary in strength or the designation of strength.

CLINICAL PHARMACOLOGY

Allergenic extracts for scratch, prick or puncture testing, used according to the DOSAGE AND ADMINISTRATION section, produce erythema or erythema and wheal reactions in patients with significant IgE-mediated sensitivity to the relevant allergen. This allergic inflammatory response, although not completely understood, is thought to begin with reaction of antigen with IgE on the surface of basophils or mast cells, which initiates a series of biochemical events resulting in the production of histamine and other mediators. These, in turn, produce the immediate-type "wheal and flare" skin reaction.

INDICATIONS & USAGE

Certain diagnostics carry labeling which states **Allergenic Extract for Diagnostic Use Only**. Data to support the therapeutic use of products labeled with this statement have not been established. 14 In addition to a carefully taken history, the use of glycerin-containing extracts in scratch, prick or puncture testing is an accepted method in the diagnosis of allergic conditions. 1, 2, 3 Extracts of all allergens do not produce equivalent results in scratch, prick or puncture tests. The intensity of the skin reactions produced will be determined by two factors: the degree of sensitivity of the patient, and the nature of the allergenic extract applied.

Scratch, prick or puncture tests are not as sensitive as the intradermal test, but are safer and cause less discomfort. They may, therefore, be the method of choice when a large number of tests are needed, or when testing the pediatric patient. In some cases, where the relatively insensitive scratch, prick or puncture tests are negative or do not confirm the allergic history, follow-up intradermal tests may be positive. However, ANTIGENS PRODUCING LARGE 3 to 4+ SCRATCH, PRICK OR PUNCTURE TESTS SHOULD NOT BE TESTED INTRADERMALLY.

CONTRAINDICATIONS

There are no known absolute contraindications to allergy skin testing. Patients with cardiovascular diseases or pulmonary diseases such as symptomatic asthma, and/or who are receiving cardiovascular drugs such as beta blockers, may be at higher risk for severe adverse reactions. These patients may also be more refractory to the normal anaphylaxis treatment regimen.

WARNINGS

Excessively large local reactions or systemic reactions are more likely to occur if the patient is skin tested shortly after exposure to large amounts of antigen to which s/he is sensitive. Use caution when skin testing patients during a season when pollen is present. Refer to boxed WARNINGS Section.

PRECAUTIONS

1. General

Always have injectable epinephrine and a tourniquet available when tests are being made. (See ADVERSE REACTIONS section.) Generally 50 to 60 scratch, prick or puncture tests can be applied safely at one sitting. Patients whose history suggests severe sensitivity should have only 5 to 10 tests applied at a time and these tests applied to the volar surface of one arm. These tests should not all be of the same type of antigen; that is, all grass pollens, all weed pollens, all danders, etc. One or two tests from several classes of antigens should be applied at a time. As soon as a large wheal begins to develop, wipe the antigen from it with a damp cotton sponge. After 10 minutes wipe off all the antigens with a damp cotton sponge, followed by a dry cotton sponge. Be careful not to wipe antigen from a positive reaction onto an adjacent test site.

2 Information for Patients

Patients should be instructed in the recognition of adverse reactions to diagnostic testing. Patients should be made to understand the importance of a 30 minute observation period, and be warned to return to the office promptly if symptoms occur after leaving.

3. Drug Interactions

Patients on non-selective beta blockers may be more reactive to allergens given for testing or treatment and may be unresponsive to the usual doses of epinephrine used to treat allergic reactions.⁶ Certain medications may lessen the skin test wheal and erythema responses elicited by allergens and histamine for varying time periods. Conventional antihistamines should be discontinued at least 5 days before skin testing. Long acting antihistamines should be discontinued for at least 3 weeks prior to skin testing.⁹ Topical steroids should be discontinued at the skin test site for at least 2-3 weeks before skin testing.^{9, 10} Tricyclic antidepressants such as Doxepin should be withheld for at least 7 days before skin testing.¹¹ Topical local anesthetics may suppress the flare responses and should be avoided in skin test sites.¹²

4. Carcinogenesis, mutagenesis, Impairment of Fertility

Long-term studies in animals have not been conducted with allergenic extracts to determine their potential for carcinogenicity, mutagenicity or impairment of fertility.

5. Pregnancy

4,5
Animal reproduction studies have not been conducted with allergenic extracts. It is also not known whether allergenic extracts can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Allergenic extracts should be given to a pregnant woman only if clearly needed.

6. Nursing Mothers

There are no current studies on secretion of the allergenic extract components in human milk or effect on the nursing infant. Because many drugs are excreted in human milk, caution should be exercised when allergenic extracts are administered to a nursing woman.

7. Pediatric Use

Wheal sizes in response to allergen skin testing can be smaller in infants than in adults. The skin response to histamine parallels that for allergens; therefore, appropriate positive control skin tests should always be performed.¹

8. Geriatric Use

Skin test wheal size decreases with age. The decrease in allergen-induced skin test reaction parallels that to histamine; therefore, appropriate positive skin test controls should always be performed.¹

ADVERSE REACTIONS

1. Local Reactions

If a severe local reaction occurs during scratch, prick or puncture testing, WIPE OFF test antigen. Large, persistent local reactions or minor exacerbations of the patient's allergic symptoms may be treated by local cold applications and/or the use of oral antihistamines, but they should be considered a warning of possible severe systemic reactions.

2. Systemic Reactions

With careful attention to dosage and administration, such reactions occur infrequently, but it must be

remembered that allergenic extracts are highly potent in sensitive individuals and OVERDOSE could result in anaphylactic symptoms. Therefore, it is imperative that physicians administering allergenic extracts understand and be prepared for the treatment of severe reactions.

Adverse reaction frequency data for allergenic extract administration for testing and treatment show that risk is low.^{7, 8}

It cannot be overemphasized that, under certain unpredictable combinations of circumstances, anaphylactic shock is a possibility. Other possible systemic reaction symptoms include fainting, pallor, bradycardia, hypotension, angioedema, cough, wheezing, conjunctivitis, rhinitis and urticaria. **If a systemic or anaphylactic reaction does occur, WIPE OFF test antigen, apply a tourniquet above the site of injection, if tests are performed on the arms, and inject the 1:1,000 epinephrine-hydrochloride intramuscularly or subcutaneously into the opposite arm. Loosen the tourniquet at least every 10 minutes. Do not obstruct arterial blood flow with the tourniquet.**

EPINEPHRINE:

ADULT DOSAGE: 0.3 to 0.5 mL should be injected. Repeat in 5 to 10 minutes if necessary.

PEDIATRIC DOSAGE: The usual initial dose is 0.01 mg (mL) per kg body weight or 0.3 mg (mL) per square meter of body surface area. Suggested dosage for infants to 2 years of age is 0.05 mL to 0.1 mL; for children 2 to 6 years, 0.15 mL; and children 6 to 12 years, 0.2 mL. Single pediatric doses should not exceed 0.3 mg (mL). Doses may be repeated as frequently as every 20 minutes, depending on the severity of the condition and the response of the patient. After administration of epinephrine, profound shock or vasomotor collapse should be treated with intravenous fluids, and other appropriate drugs. Oxygen should be given by mask. Intravenous antihistamine, theophylline or corticosteroids may be used if necessary after adequate epinephrine and circulatory support have been given. Emergency resuscitation measures and personnel trained in their use should be available immediately in the event of a serious systemic or anaphylactic reaction not responsive to the above measures (Ref. J. Allergy Clin. Immunol. 77 (2): 271-273, 1986). Rarely are all of the above measures necessary; the tourniquet and epinephrine usually produce prompt responses. However, the physician should be prepared in advance for all contingencies. Promptness in beginning emergency treatment measures is of utmost importance.

3. Adverse Event Reporting

Report all adverse events to Jubilant HollisterStier LLC Customer Technical Services Department at 1 (800) 992-1120. A voluntary adverse event reporting system for health professionals is available through the FDA MEDWATCH program. Preprinted forms (FDA Form 3500) are available from the FDA by calling 1 (800) FDA-1088. Completed forms should be mailed to MEDWATCH, 5600 Fisher Lane, Rockville, MD 20852-9787 or Fax to: 1 (800) FDA-0178.

OVERDOSAGE

See ADVERSE REACTIONS Section.

DOSAGE & ADMINISTRATION

1. General

Parenteral Drug Products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

2. Scratch, Prick or Puncture Testing Methods

There are two general methods of skin testing. (1) The skin is scarified first, and the extract is then applied. (2) A drop of extract is put onto the skin, and a prick or puncture is made through the drop. Avoid touching tip of dropper to skin. Either method is satisfactory, but the second requires that the instrument be cleansed between tests or that separate needles be used.

The extracts for scratch, prick or puncture testing are supplied in dropper vials and should be kept in a rack or box in rows of 10 vials corresponding to the rows of tests to be applied to the skin.

All skin tests should be validated by appropriate positive control tests (e.g., histamine) and negative control tests (e.g., Glycerin, Albumin Saline with Phenol (0.4%), or Buffered Saline with Phenol). The negative control test should be the same material as is used as a diluting fluid in the tested extracts. Diluting fluid is used in the same way as an active test extract.

Test sites should be examined at 15 and 30 minutes. To prevent excessive absorption, wipe off antigens producing large reactions as soon as the wheal appears. Record the size of the reaction. Delayed reactions may rarely occur from tests, so it may be helpful to examine the test sites in 24 hours.

Use of Scarifiers and Spacing. Make scarifications at least 2.5 cm apart. Use more space between pollen tests to prevent smearing into adjacent sites. Hold the scarifier between the thumb and index finger, press the sharp edge of the instrument against the skin and twirl instrument rapidly. The scratch should disrupt only the outer layers of epidermis but should not produce immediate oozing of blood. The amount of pressure needed to produce a satisfactory scratch will vary between patients according to the thickness or fragility of their skin. Experience will indicate the proper amount of pressure to exert in making the scratch. If the scarifier is kept sharp and the scratch made quickly, discomfort to the patient is minimized.

Use of Prick Test Needles. The skin is cleaned and single drops of each extract applied to the properly identified test sites. A small, sterile disposable needle, such as a 1/2-inch 26 gauge needle (with the bevel up), a bifurcated vaccinating needle, or a Prick Lancetter™ is inserted through the drop superficially into the skin, the skin lifted slightly and the needle withdrawn. No bleeding should be produced. After about 1 minute the extract may be wiped away.

3. Most Satisfactory Sites for Testing

Prior to testing, clean the skin area to be tested with ether or alcohol and allow to dry. Use a sterile instrument for each patient. The back or the volar surface of the arms are the most satisfactory sites for testing. Skin of the posterior thighs or abdomen may be used if necessary. Avoid very hairy areas where possible, since the reactions will be smaller and more difficult to interpret. The most satisfactory areas of the back are from the posterior axillary fold to 2.5 cm from the spinal column, and from the top of the scapula to the lower rib margins. The best areas of the arms are the volar surfaces from the axilla to 2.5 or 5 cm above the wrist, skipping the anti-cubital space.

4. Use of Antigen Mixes

The use of complicated mixes of unrelated pollens for testing is not recommended since in the case of a positive reaction, it does not indicate which pollen(s) are responsible, and, in the case of a negative reaction, it fails to indicate whether the individual pollens at full concentration would give a positive reaction.

5. Reading Skin Test Reactions

A positive reaction consists of an urticarial wheal with surrounding erythema (resembling somewhat a mosquito bite reaction) larger than the control site. The smallest reaction considered positive is erythema with a central papule at least 5 mm in diameter. In some instances with no reaction at the control site, erythema may be considered an indication of sensitivity. In general, the size of wheal and erythema response correlates directly with the patient's sensitivity to that allergen.

Standardized Products

(a) Mites:

The skin test concentration of 30,000 AU/mL in dropper vials is used for scratch, prick or puncture testing. Puncture tests performed on 12 highly sensitive subjects showed the following:

Species	Mean Sum of Wheal ± Std. Dev. (mm)	Mean Sum of Erythema ± Std. Dev. (mm)
D. farinae	22.4 ± 10.7	82.2 ± 21.7
D. pteronyssinus	24.0 ± 9.9	89.3 ± 24.5

The sum of a skin response is the sum of the longest diameter and the mid-point orthogonal diameter.

(b) Cat Hair and Cat Pelt: The skin test concentration of 10,000 BAU/mL (10-19.9 Fel d 1 Units/mL) in dropper vials is used for prick or puncture testing. Puncture tests performed on 15 highly sensitive subjects showed the following:

Product	Mean Sum of Wheal ± Std. Dev (mm)	Mean Sum of Erythema ± Std. Dev (mm)
Standardized Cat Hair	15.1 ± 3.8	73.3 ± 14.3
Standardized Cat Pelt	13.9 ± 4.3	67.3 ± 13.3

The sum of a skin response is the sum of the longest diameter and the mid-point orthogonal diameter.

(c) Ragweed pollen (Short Ragweed or Giant and Short Ragweed Mixture) Antigen E Assayed: Short Ragweed extract at 1:20 w/v in 50% glycerin containing approximately 100 to 300 units of Amb a 1/mL or Giant and Short Ragweed Mix at 1:20 w/v in 50% glycerin containing approximately 50 to 150 units of Amb a 1/mL are usually used for scratch, prick or puncture testing.

Refer to the following table to determine the skin test sensitivity grade. The corresponding ΣE (sum of the longest diameter and the mid-point orthogonal diameters of erythema) is also presented.

Grade	Erythema mm	Papule or Wheal mm	Corresponding mm ΣE
0	<5	<5	<10
±	5-10	5-10	10-20
1+	11-20	5-10	20-40
2+	21-30	5-10	40-60
3+	31-40	10-15 (a)	60-80
4+	>40	>15 (b)	>80

(a) or with pseudopods (b) or with many pseudopods

A positive skin reaction to any allergen must be interpreted in light of the patient's history of symptoms, time of the year, and known exposures.

THE SKIN TESTS ARE IN NO WAY A SUBSTITUTE FOR A CAREFUL ALLERGIC HISTORY. THEY SERVE AS ADDITIONAL INFORMATION TO AID IN IDENTIFYING CAUSATIVE ALLERGENS IN PATIENTS WITH ALLERGIC DISORDERS.

6. Geriatric Use

The dose is the same in patients of all age groups. Because the wheal size in response to allergen skin testing decreases with age, appropriate histamine positive control skin tests must be performed.¹

7. Pediatric Use

The dose is the same in patients of all age groups. Wheal size in response to allergen skin testing can be smaller in infants than in adults. Appropriate histamine positive control skin tests must be performed.¹

HOW SUPPLIED

In 5 mL dropper bottles of extract at 1:10 w/v except pollen at 1:20, APTM extracts at 1:50 w/v, except

AP™ Dog Hair-Dander at 1:100 w/v, AP™ House Dust at 20,000 PNU/mL, some mixes as Concentrate, and Standardized products at AU/mL (Mite extracts at 30,000 AU/mL) or BAU/mL (Cat Hair and Pelt extracts at 10,000 BAU/mL) value. Strengths are listed on product labels.

STORAGE

The expiration date of the diagnostic extracts is listed on the container label. The extract should be stored at 2° - 8°C and kept at this temperature range during office use.

LIMITED WARRANTY

A number of factors beyond our control could reduce the efficacy of this product or even result in an ill effect following its use. These include storage and handling of the product after it leaves our hands, diagnosis, dosage, method of administration, and biological differences in individual patients. Because of these factors, it is important that this product be stored properly, and that the directions be followed carefully during use. No warranty, express or implied, including any warranty of merchantability or fitness, is made. Representatives of the Company are not authorized to vary the terms or the contents of any printed labeling, including the package insert, for this product except by printed notice from the Company's headquarters. The prescriber and user of this product must accept the terms hereof.

REFERENCES

1. Middleton, Elliott, Jr., C.E. Reed, E.F. Ellis (ed.) Allergy Principles and Practice. Fourth Edition, Vol. 1. C.V. Mosby. 1993.
2. Sheldon, J.M., R.G. Lovell, K.P. Mathews. A Manual of Clinical Allergy. W.B. Saunders. 1967.
3. Tuft, L., H.L. Mueller. Allergy in Children. W.B. Saunders. 1970.
4. DuBuske, L.M., C.J. Ling, A.L. Sheffer. Special problems regarding allergen immunotherapy. Immunol. Allergy Clin. North Am. (USA). 12(1): 145-175, 1992.
5. Weinstein, A.M., B.D. Dubin, W.K. Podleski, S.L. Spector, R.S. Farr. Asthma and pregnancy. JAMA. 124(11): 1161-1165, 1979.
6. Jacobs, Robert L., Geoffrey W. Rake, Jr., et.al. Potentiated anaphylaxis in patients with drug-induced beta-adrenergic blockade. J. Allergy Clin. Immunol. 68(2): 125-127, August 1981.
7. Lockey, Richard F., Linda M. Benedict, Paul C. Turkeltaub, Samuel C. Bukantz. Fatalities from immunotherapy (IT) and skin testing (ST). J. Allergy Clin. Immunol. 79(4): 660-677, 1987.
8. Turkeltaub, Paul C., Peter J. Gergen. The risk of adverse reactions from percutaneous prick-puncture allergen skin testing, venipuncture, and body measurements: Data from the second National Health and Nutrition Examination Survey 1976-80 (NHANES II). J. Allergy Clin. Immunol. 84(6): 886-890, Dec. 1989.
9. Pipkorn, Ulf. Pharmacological influence of anti-allergic medication on *In Vivo* allergen testing. Allergy. 43: 81-86, 1988.
10. Andersson, M., U. Pipkorn. Inhibition of the dermal immediate allergic reaction through prolonged treatment with topical glucocorticosteroids. J. Allergy Clin. Immunol. 79 (2): 345-349, February 1987.
11. Rao, Kamineni S., et al. Duration of suppressive effect of tricyclic anti-depressants on histamine induced wheal and flare reactions on human skin. J. Allergy Clin. Immunol. 82: 752-757, November 1988.
12. Pipkorn, Ulf, M. Andersson. Topical dermal anesthesia inhibits the flare but not the wheal response to allergen and histamine in the skin prick test. Clin. Allergy. 17: 307-311, 1987.
13. Turkeltaub, Paul C., Suresh, C. Rastogi, Harold Baer. Office of Biologics Research and Review skin test method for evaluation of subject sensitivity to standardized allergenic extracts and for assignment of allergy units to reference preparations using the ID₅₀EAL method (Intradermal Dilution for 50 mm Sum of Erythema Determines the Allergy Unit). Methods of the Allergenic Products Branch Office of Biologics Research and Review, FDA, Bethesda, MD 20892. Revised May 9, 1986.
14. Food and Drug Administration. Biological products; Allergenic extracts classified in Category

Acacia, Golden, 5 mL 1:20 w/v Carton Label

ALLERGENIC EXTRACT

ACACIA, GOLDEN

Preservative:
50% Glycerin v/v

Inactive Ingredients:
0.5% Sodium chloride
0.275% Sodium bicarbonate

A. longifolia



(01)00365044100711
(17)201101
(10)T0000182
(21)000000020001

1:20 W/V

Item: 1007ED No U.S. Standard of Potency
Lot: T0000182 Dose/Route: 1 drop topically
Exp: 2020Nov01 5 mL Item: 1007ED

Non-Returnable Store at 2-8°C NDC: 65044-1007-1
RX Only Sterile Until Opened US License No. 1272
50000000649 - H01

50000000099 - H01

Acacia, Golden, 5 mL 1:20 w/v Vial Label

ALLERGENIC EXTRACT

ACACIA, GOLDEN
ACACIA LONGIFOLIA
A. longifolia

5 mL **1:20 W/V** RX Only Sterile
Dose/Route: 1 drop topically Until Opened
Store at 2-8°C

Preservative 50% Glycerin v/v Item: 1007ED
U.S. License No. 1272 Lot: T0000182
Exp: 2020Nov01

Jubilant HollisterStier LLC Spokane, WA 99207

50000002619 - H01
5000000143-H01

Alder, Red, 5 mL 1:20 w/v Carton Label

ALLERGENIC EXTRACT

ALDER, RED

Preservative:
50% Glycerin v/v

Inactive Ingredients:
0.5% Sodium chloride
0.275% Sodium
bicarbonate

A. rubra



(01)00365044101916
(17)201101
(10)T0000184
(21)000000020001

1:20 W/V

Item: 1019ED
Lot: T0000184
Exp: 2020Nov01

No U.S. Standard of Potency
Dose/Route: 1 drop topically

NDC: 65044-1019-1
US License No. 1272

5 mL Item: 1019ED

50000000649 - H01

Store at 2-8°C
RX Only Sterile Until Opened

Non-Returnable

50000000099 - H01

Alder, Red, 5 mL 1:20 w/v Vial Label

ALLERGENIC EXTRACT

ALDER, RED
ALNUS RUBRA (OREGONA)
A. rubra

5 mL **1:20 W/V**
Dose/Route: 1 drop topically

RX Only Sterile
Until Opened
Store at 2-8°C

Preservative 50% Glycerin v/v
U.S. License No. 1272

Item: 1019ED
Lot: T0000184
Exp: 2020Nov01

Jubilant HollisterStier LLC Spokane, WA 99207

50000002619 - H01
5000000143-H01

Ash, White, 5 mL 1:20 w/v Carton Label

ALLERGENIC EXTRACT

ASH, WHITE



Preservative:
50% Glycerin w/v

Inactive Ingredients:
0.5% Sodium chloride
0.275% Sodium
bicarbonate

F. americana

(01)00365044106119
(17)201101
(10)T0000187
(21)000000020001

1:20 W/V

Item: 1061ED
Lot: T0000187
Exp: 2020Nov01

No U.S. Standard of Potency
Dose/Route: 1 drop topically

NDC: 65044-1061-1
US License No. 1272

5 mL Item: 1061ED

50000000649 - H01

Non-Returnable

Store at 2-8°C
RX Only Sterile Until Opened

50000000099 - H01

Ash, White, 5 mL 1:20 w/v Vial Label

ALLERGENIC EXTRACT

**ASH, WHITE
FRAXINUS AMERICANA
F. americana**

5 mL **1:20 W/V**
Dose/Route: 1 drop topically

RX Only Sterile
Until Opened
Store at 2-8°C

Preservative 50% Glycerin w/v
U.S. License No. 1272

Item: 1061ED
Lot: T0000187
Exp: 2020Nov01

Jubilant HollisterStier LLC Spokane, WA 99207

50000002619 - H01

5000000143-H01

Beech, American, 5 mL 1:20 w/v Carton Label

ALLERGENIC EXTRACT

BEECH, AMERICAN

Preservative:
50% Glycerin v/v

Inactive Ingredients:
0.5% Sodium chloride
0.275% Sodium
bicarbonate

F. grandifolia



(01)00365044112110
(17)201101
(10)T0000192
(21)000000020001

1:20 W/V

Item: 1121ED
Lot: T0000192
Exp: 2020Nov01

No U.S. Standard of Potency
Dose/Route: 1 drop topically

NDC: 65044-1121-1
US License No. 1272

5 mL Item: 1121ED

50000000649 - H01

Store at 2-8°C
RX Only Sterile Until Opened

Non-Returnable

50000000099 - H01

Beech, American, 5 mL 1:20 w/v Vial Label

ALLERGENIC EXTRACT

**BEECH, AMERICAN
FAGUS GRANDIFOLIA
F. grandifolia**

5 mL **1:20 W/V**
Dose/Route: 1 drop topically

RX Only Sterile
Until Opened
Store at 2-8°C

Preservative 50% Glycerin v/v
U.S. License No. 1272

Item: 1121ED
Lot: T0000192
Exp: 2020Nov01

Jubilant HollisterStier LLC Spokane, WA 99207

50000002619 - H01

10H-F3100000000

Birch Mix PRW, 5 mL 1:20 w/v Carton Label

ALLERGENIC EXTRACT

BOTTLEBRUSH TREE

Preservative:
50% Glycerin w/v

Inactive Ingredients:
0.5% Sodium chloride
0.275% Sodium
bicarbonate

M. citrina



(01)00365044120818
(17)201101
(10)T0000197
(21)000000020001

1:20 W/V

Item: 1208ED
Lot: T0000197
Exp: 2020Nov01

No U.S. Standard of Potency
Dose/Route: 1 drop topically

NDC: 65044-1208-1
US License No. 1272

5 mL Item: 1208ED

50000000649 - H01

Store at 2-8°C
RX Only Sterile Until Opened

Non-Returnable

50000000099 - H01

Bottlebrush Tree, 5 mL 1:20 w/v Vial Label

ALLERGENIC EXTRACT

BOTTLEBRUSH TREE
CALLISTEMON CITRINUS
M. citrina

5 mL **1:20 W/V**
Dose/Route: 1 drop topically

RX Only Sterile
Until Opened
Store at 2-8°C

Preservative 50% Glycerin w/v
U.S. License No. 1272

Item: 1208ED
Lot: T0000197
Exp: 2020Nov01

Jubilant HollisterStier LLC Spokane, WA 99207

50000002619 - H01

50000001434401

Boxelder-Maple Mix, 5 mL 1:20 w/v Carton Label

ALLERGENIC EXTRACT

BOXELDER-MAPLE MIX

Preservative:
50% Glycerin v/v

Inactive Ingredients:
0.5% Sodium chloride
0.275% Sodium
bicarbonate

**33.3% each of A.
negundo, A.
saccharum, A. rubrum**



(01)00365044121419
(17)201101
(10)T0000199
(21)000000020001

1:20 W/V

Item: 1214ED
Lot: T0000199
Exp: 2020Nov01

No U.S. Standard of Potency
Dose/Route: 1 drop topically

NDC: 65044-1214-1
US License No. 1272

5 mL Item: 1214ED

50000000649 - H01

Store at 2-8°C
RX Only Sterile Until Opened

Non-Returnable

50000000099 - H01

Boxelder-Maple Mix, 5 mL 1:20 w/v Vial Label

ALLERGENIC EXTRACT

BOXELDER-MAPLE MIX BHR (BOXELDER, HARD, RED)

**33.3% each of A. negundo, A. saccharum,
A. rubrum**

5 mL **1:20 W/V**
Dose/Route: 1 drop topically

RX Only Sterile
Until Opened
Store at 2-8°C

Preservative 50% Glycerin v/v
U.S. License No. 1272

Item: 1214ED
Lot: T0000199
Exp: 2020Nov01

Jubilant HollisterStier LLC Spokane, WA 99207

50000000649 - H01

101-43100000005

Brome, Smooth, 5 mL 1:20 w/v Carton Label

ALLERGENIC EXTRACT

BROME, SMOOTH

Preservative:
50% Glycerin w/v

Inactive Ingredients:
0.5% Sodium chloride
0.275% Sodium
bicarbonate

B. inermis



(01)00365044123819
(17)201101
(10)T0000202
(21)000000020001

1:20 W/V

Item: 1238ED
Lot: T0000202
Exp: 2020Nov01

No U.S. Standard of Potency
Dose/Route: 1 drop topically

NDC: 65044-1238-1
US License No. 1272

5 mL Item: 1238ED

50000000649 - H01

Non-Returnable

Store at 2-8°C
RX Only Sterile Until Opened

50000000099 - H01

Brome, Smooth, 5 mL 1:20 w/v Vial Label

ALLERGENIC EXTRACT

BROME, SMOOTH
BROMUS INERMIS
B. inermis

5 mL **1:20 W/V**
Dose/Route: 1 drop topically

RX Only Sterile
Until Opened
Store at 2-8°C

Preservative 50% Glycerin w/v
U.S. License No. 1272

Item: 1238ED
Lot: T0000202
Exp: 2020Nov01

Jubilant HollisterStier LLC Spokane, WA 99207

50000002619 - H01

5000000143-H01

Cedar, Mountain, 5 mL 1:20 w/v Carton Label

ALLERGENIC EXTRACT

CEDAR, MOUNTAIN

Preservative:
50% Glycerin v/v

Inactive Ingredients:
0.5% Sodium chloride
0.275% Sodium
bicarbonate

J. ashei



(01)00365044133719
(17)201101
(10)T0000208
(21)000000020001

1:20 W/V

Item: 1337ED
Lot: T0000208
Exp: 2020Nov01

Dose/Route: 1 drop topically

NDC: 65044-1337-1
US License No. 1272

5 mL

Item: 1337ED

50000002595 - H01

Store at 2-8°C

RX Only Sterile Until Opened

Non-Returnable

50000000099-H01

Cedar, Mountain, 5 mL 1:20 w/v Vial Label

ALLERGENIC EXTRACT

**CEDAR, MOUNTAIN
JUNIPERUS ASHEI
J. ashei**

5 mL **1:20 W/V**
Dose/Route: 1 drop topically

RX Only Sterile
Until Opened
Store at 2-8°C

Preservative 50% Glycerin v/v
U.S. License No. 1272

Item: 1337ED
Lot: T0000208
Exp: 2020Nov01

Jubilant HollisterStier LLC Spokane, WA 99207

50000002619 - H01

5000000143-H01

Cedar, Red, 5 mL 1:20 w/v Carton Label

ALLERGENIC EXTRACT

CEDAR, RED

Preservative:
50% Glycerin w/v

Inactive Ingredients:
0.5% Sodium chloride
0.275% Sodium
bicarbonate

J. virginiana



(01)00365044134013
(17)201101
(10)T0000210
(21)000000020001

1:20 W/V

Item: 1340ED
Lot: T0000210
Exp: 2020Nov01

No U.S. Standard of Potency
Dose/Route: 1 drop topically

NDC: 65044-1340-1
US License No. 1272

5 mL Item: 1340ED

50000000649 - H01

Store at 2-8°C
RX Only Sterile Until Opened

Non-Returnable

50000000099 - H01

Cedar, Red, 5 mL 1:20 w/v Vial Label

ALLERGENIC EXTRACT

CEDAR, RED
JUNIPERUS VIRGINIANA
J. virginiana

5 mL **1:20 W/V**
Dose/Route: 1 drop topically

RX Only Sterile
Until Opened
Store at 2-8°C

Preservative 50% Glycerin v/v
U.S. License No. 1272

Item: 1340ED
Lot: T0000210
Exp: 2020Nov01

Jubilant HollisterStier LLC Spokane, WA 99207

50000002619 - H01

5000000143-H01

Corn, Cultivated Pollen, 5 mL 1:20 w/v Carton Label

ALLERGENIC EXTRACT

**CORN, CULTIVATED
(POLLEN)**



Preservative:
50% Glycerin w/v

Inactive Ingredients:
0.5% Sodium chloride
0.275% Sodium
bicarbonate

Z. mays

(01)00365044141516
(17)201101
(10)T0000215
(21)000000020051

1:20 W/V

Item: 1415ED
Lot: T0000215
Exp: 2020Nov01

No U.S. Standard of Potency
Dose/Route: 1 drop topically

NDC: 65044-1415-1
US License No. 1272

5 mL Item: 1415ED

50000000649 - H01

Non-Returnable

Store at 2-8°C
RX Only Sterile Until Opened

50000000099 - H01

Corn, Cultivated Pollen, 5 mL 1:20 w/v Vial Label

ALLERGENIC EXTRACT

**CORN, CULTIVATED (POLLEN)
ZEA MAYS
Z. mays**

5000000143-H01
50000002619 - H01

5 mL **1:20 W/V**
Dose/Route: 1 drop topically

RX Only Sterile
Until Opened
Store at 2-8°C

Preservative 50% Glycerin w/v
U.S. License No. 1272

Item: 1415ED
Lot: T0000215
Exp: 2020Nov01

Jubilant HollisterStier LLC Spokane, WA 99207

Johnson Grass, 5 mL 1:20 w/v Carton Label

ALLERGENIC EXTRACT

JOHNSON GRASS

Preservative:
50% Glycerin w/v

Inactive Ingredients:
0.5% Sodium chloride
0.275% Sodium
bicarbonate

S. halepense



(01)00365044174514
(17)201101
(10)T0000241
(21)000000020051

1:20 W/V

Item: 1745ED
Lot: T0000241
Exp: 2020Nov01

No U.S. Standard of Potency
Dose/Route: 1 drop topically

NDC: 65044-1745-1
US License No. 1272

5 mL Item: 1745ED

50000000649 - H01

Store at 2-8°C
RX Only Sterile Until Opened

Non-Returnable

50000000099 - H01

Johnson Grass, 5 mL 1:20 w/v Vial Label

ALLERGENIC EXTRACT

**JOHNSON GRASS
SORGHUM HALEPENSE
S. halepense**

5 mL **1:20 W/V**
Dose/Route: 1 drop topically

RX Only Sterile
Until Opened
Store at 2-8°C

Preservative 50% Glycerin w/v
U.S. License No. 1272

Item: 1745ED
Lot: T0000241
Exp: 2020Nov01

Jubilant HollisterStier LLC Spokane, WA 99207

50000002619 - H01

5000000143 - H01

Oats, Common Cultivated Pollen, 5 mL 1:20 w/v Carton Label

ALLERGENIC EXTRACT

OATS, COMMON CULTIVATED (POLLEN)



Preservative:
50% Glycerin w/v

Inactive Ingredients:
0.5% Sodium chloride
0.275% Sodium
bicarbonate

A. sativa

(01)00365044204211
(17)201101
(10)T0000267
(21)000000020151

1:20 W/V

Item: 2042ED
Lot: T0000267
Exp: 2020Nov01

No U.S. Standard of Potency
Dose/Route: 1 drop topically

NDC: 65044-2042-1
US License No. 1272

5 mL Item: 2042ED

50000000649 - H01

Store at 2-8°C
RX Only Sterile Until Opened

Non-Returnable

50000000099 - H01

Oats, Common Cultivated Pollen, 5 mL 1:20 w/v Vial Label

ALLERGENIC EXTRACT

OATS, COMMON CULTIVATED (POLLEN) AVENA SATIVA A. sativa

50000000649 - H01

5 mL **1:20 W/V**
Dose/Route: 1 drop topically

RX Only Sterile
Until Opened
Store at 2-8°C

Preservative 50% Glycerin v/v
U.S. License No. 1272

Item: 2042ED
Lot: T0000267
Exp: 2020Nov01

101-43-100000005

Jubilant HollisterStier LLC Spokane, WA 99207

POLLENS - GRASSES, BROME, SMOOTH BROMUS INERMIS

brome, smooth bromus inermis injection, solution

Product Information

Product Type	NON-STANDARDIZED ALLERGENIC	Item Code (Source)	NDC:65044-1238
Route of Administration	PERCUTANEOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BROMUS INERMIS POLLEN (UNII: 766QT72BK6) (BROMUS INERMIS POLLEN - UNII:766QT72BK6)	BROMUS INERMIS POLLEN	0.05 g in 1 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65044-1238-1	5 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA103888	04/19/1941	

POLLENS - GRASSES, CORN, CULTIVATED ZEA MAYS

corn, cultivated zea mays injection, solution

Product Information

Product Type	NON-STANDARDIZED ALLERGENIC	Item Code (Source)	NDC:65044-1415
Route of Administration	PERCUTANEOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ZEA MAYS POLLEN (UNII: 74PD8J616H) (ZEA MAYS POLLEN - UNII:74PD8J616H)	ZEA MAYS POLLEN	0.05 g in 1 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65044-1415-1	5 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA103888	04/19/1941	

POLLENS - GRASSES, JOHNSON GRASS SORGHUM HALEPENSE				
johnson grass sorghum halepense injection, solution				
Product Information				
Product Type	NON-STANDARDIZED ALLERGENIC	Item Code (Source)	NDC:65044-1745	
Route of Administration	PERCUTANEOUS			
Active Ingredient/Active Moiety				
	Ingredient Name	Basis of Strength	Strength	
	SORGHUM HALEPENSE POLLEN (UNII: 577VA5B4HP) (SORGHUM HALEPENSE POLLEN - UNII:577VA5B4HP)	SORGHUM HALEPENSE POLLEN	0.05 g in 1 mL	
Inactive Ingredients				
	Ingredient Name	Strength		
	GLYCERIN (UNII: PDC6A3C0OX)			
	SODIUM CHLORIDE (UNII: 451W47IQ8X)			
	SODIUM BICARBONATE (UNII: 8MDF5V39QO)			
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65044-1745-1	5 mL in 1 VIAL; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
BLA	BLA103888	04/19/1941		

POLLENS - GRASSES, OATS, COMMON, CULTIVATED AVENA SATIVA			
oats, common, cultivated avena sativa injection, solution			
Product Information			
Product Type	NON-STANDARDIZED ALLERGENIC	Item Code (Source)	NDC:65044-2042

Route of Administration	PERCUTANEOUS			
Active Ingredient/Active Moiety				
	Ingredient Name	Basis of Strength	Strength	
	AVENA SATIVA POLLEN (UNII: A7IKY24TR7) (AVENA SATIVA POLLEN - UNII:A7IKY24TR7)	AVENA SATIVA POLLEN	0.05 g in 1 mL	
Inactive Ingredients				
	Ingredient Name	Strength		
	GLYCERIN (UNII: PDC6A3C0OX)			
	SODIUM CHLORIDE (UNII: 451W47IQ8X)			
	SODIUM BICARBONATE (UNII: 8MDF5V39QO)			
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65044-2042-1	5 mL in 1 VIAL; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
BLA	BLA103888	04/19/1941		

POLLENS - GRASSES, SOUTHERN GRASS MIX			
pollens - grasses, southern grass mix injection, solution			
Product Information			
Product Type	NON-STANDARDIZED ALLERGENIC	Item Code (Source)	NDC:65044-0855
Route of Administration	PERCUTANEOUS		
Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength
	POA PRATENSIS POLLEN (UNII: SCB8J7LS3T) (POA PRATENSIS POLLEN - UNII:SCB8J7LS3T)	POA PRATENSIS POLLEN	100000 [BAU] in 1 mL
	DACTYLIS GLOMERATA POLLEN (UNII: 83N78IDA7P) (DACTYLIS GLOMERATA POLLEN - UNII:83N78IDA7P)	DACTYLIS GLOMERATA POLLEN	100000 [BAU] in 1 mL
	AGROSTIS GIGANTEA POLLEN (UNII: HU8V6E7HOA) (AGROSTIS GIGANTEA POLLEN - UNII:HU8V6E7HOA)	AGROSTIS GIGANTEA POLLEN	100000 [BAU] in 1 mL
	PHLEUM PRATENSE POLLEN (UNII: 65M88RW2EG) (PHLEUM PRATENSE POLLEN - UNII:65M88RW2EG)	PHLEUM PRATENSE POLLEN	100000 [BAU] in 1 mL
	ANTHOXANTHUM ODORATUM POLLEN (UNII: 2KIK19R45Y) (ANTHOXANTHUM ODORATUM POLLEN - UNII:2KIK19R45Y)	ANTHOXANTHUM ODORATUM POLLEN	100000 [BAU] in 1 mL
	SORGHUM HALEPENSE POLLEN (UNII: 577VA5B4HP) (SORGHUM HALEPENSE POLLEN - UNII:577VA5B4HP)	SORGHUM HALEPENSE POLLEN	0.05 g in 1 mL

CYNODON DACTYLON POLLEN (UNII: 175F461W10) (CYNODON DACTYLON POLLEN - UNII:175F461W10)	CYNODON DACTYLON POLLEN	10000 [BAU] in 1 mL
--	-------------------------	------------------------

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65044-0855-1	5 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA103888	04/19/1941	

POLLENS - GRASSES, SOUTHERN GRASS MIX

pollens - grasses, southern grass mix injection, solution

Product Information

Product Type	NON-STANDARDIZED ALLERGENIC	Item Code (Source)	NDC:65044-0857
Route of Administration	PERCUTANEOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
POA PRATENSIS POLLEN (UNII: SCB8J7LS3T) (POA PRATENSIS POLLEN - UNII:SCB8J7LS3T)	POA PRATENSIS POLLEN	10000 [BAU] in 1 mL
DACTYLIS GLOMERATA POLLEN (UNII: 83N78IDA7P) (DACTYLIS GLOMERATA POLLEN - UNII:83N78IDA7P)	DACTYLIS GLOMERATA POLLEN	10000 [BAU] in 1 mL
AGROSTIS GIGANTEA POLLEN (UNII: HU8V6E7HOA) (AGROSTIS GIGANTEA POLLEN - UNII:HU8V6E7HOA)	AGROSTIS GIGANTEA POLLEN	10000 [BAU] in 1 mL
PHLEUM PRATENSE POLLEN (UNII: 65M88RW2EG) (PHLEUM PRATENSE POLLEN - UNII:65M88RW2EG)	PHLEUM PRATENSE POLLEN	10000 [BAU] in 1 mL
ANTHOXANTHUM ODORATUM POLLEN (UNII: 2KIK19R45Y) (ANTHOXANTHUM ODORATUM POLLEN - UNII:2KIK19R45Y)	ANTHOXANTHUM ODORATUM POLLEN	10000 [BAU] in 1 mL
SORGHUM HALEPENSE POLLEN (UNII: 577VA5B4HP) (SORGHUM HALEPENSE POLLEN - UNII:577VA5B4HP)	SORGHUM HALEPENSE POLLEN	0.005 g in 1 mL
CYNODON DACTYLON POLLEN (UNII: 175F461W10) (CYNODON DACTYLON POLLEN - UNII:175F461W10)	CYNODON DACTYLON POLLEN	1000 [BAU] in 1 mL

Inactive Ingredients

Ingredient Name	Strength
-----------------	----------

GLYCERIN (UNII: PDC6A3C0OX)				
SODIUM CHLORIDE (UNII: 451W47IQ8X)				
SODIUM BICARBONATE (UNII: 8MDF5V39QO)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65044-0857-1	5 mL in 1 VIAL; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
BLA	BLA103888	04/19/1941		

POLLENS - TREES, ACACIA ACACIA LONGIFOLIA				
acacia longifolia injection, solution				
Product Information				
Product Type	NON-STANDARDIZED ALLERGENIC	Item Code (Source)	NDC:65044-1007	
Route of Administration	PERCUTANEOUS			
Active Ingredient/Active Moiety				
	Ingredient Name	Basis of Strength	Strength	
	ACACIA LONGIFOLIA POLLEN (UNII: 24SO2J296O) (ACACIA LONGIFOLIA POLLEN - UNII:24SO2J296O)	ACACIA LONGIFOLIA POLLEN	0.05 g in 1 mL	
Inactive Ingredients				
	Ingredient Name	Strength		
	GLYCERIN (UNII: PDC6A3C0OX)			
	SODIUM CHLORIDE (UNII: 451W47IQ8X)			
	SODIUM BICARBONATE (UNII: 8MDF5V39QO)			
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65044-1007-1	5 mL in 1 VIAL; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
BLA	BLA103888	04/19/1941		

POLLENS - TREES, ALDER, RED ALNUS RUBRA

alder, red alnus rubra injection, solution

Product Information

Product Type	NON-STANDARDIZED ALLERGENIC	Item Code (Source)	NDC:65044-1019
Route of Administration	PERCUTANEOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALNUS RUBRA POLLEN (UNII: Z0F2YK1B7H) (ALNUS RUBRA POLLEN - UNII:Z0F2YK1B7H)	ALNUS RUBRA POLLEN	0.05 g in 1 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65044-1019-1	5 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA103888	04/19/1941	

POLLENS - TREES, ASH, WHITE FRAXINUS AMERICANA

ash, white fraxinus americana injection, solution

Product Information

Product Type	NON-STANDARDIZED ALLERGENIC	Item Code (Source)	NDC:65044-1061
Route of Administration	PERCUTANEOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FRAXINUS AMERICANA POLLEN (UNII: G684LX721Q) (FRAXINUS AMERICANA POLLEN - UNII:G684LX721Q)	FRAXINUS AMERICANA POLLEN	0.05 g in 1 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65044-1061-1	5 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA103888	04/19/1941	

POLLENS - TREES, BEECH, AMERICAN FAGUS GRANDIFOLIA

beech, american fagus grandifolia injection, solution

Product Information

Product Type	NON-STANDARDIZED ALLERGENIC	Item Code (Source)	NDC:65044-1121
Route of Administration	PERCUTANEOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FAGUS GRANDIFOLIA POLLEN (UNII: 34X886W1H4) (FAGUS GRANDIFOLIA POLLEN - UNII:34X886W1H4)	FAGUS GRANDIFOLIA POLLEN	0.05 g in 1 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65044-1121-1	5 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

BLA	BLA103888	04/19/1941	
-----	-----------	------------	--

POLLENS - TREES, BIRCH MIX

birch mix injection, solution

Product Information

Product Type	NON-STANDARDIZED ALLERGENIC	Item Code (Source)	NDC:65044-1169
Route of Administration	PERCUTANEOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BETULA POPYRIFERA POLLEN (UNII: 3538FNV8AY) (BETULA POPYRIFERA POLLEN - UNII:3538FNV8AY)	BETULA POPYRIFERA POLLEN	0.05 g in 1 mL
BETULA PENDULA POLLEN (UNII: ZL5TV40C5Y) (BETULA PENDULA POLLEN - UNII:ZL5TV40C5Y)	BETULA PENDULA POLLEN	0.05 g in 1 mL
BETULA NIGRA POLLEN (UNII: 93963RFO1P) (BETULA NIGRA POLLEN - UNII:93963RFO1P)	BETULA NIGRA POLLEN	0.05 g in 1 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65044-1169-1	5 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA103888	04/19/1941	

POLLENS - TREES, BOTTLEBRUSH, CALLISTEMON SPP.

bottlebrush, callistemon citrinus injection, solution

Product Information

Product Type	NON-STANDARDIZED ALLERGENIC	Item Code (Source)	NDC:65044-1208
Route of Administration	PERCUTANEOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MELALEUCA CITRINA POLLEN (UNII: 620II98F1T) (MELALEUCA CITRINA POLLEN - UNII:620II98F1T)	MELALEUCA CITRINA POLLEN	0.05 g in 1 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65044-1208-1	5 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA103888	04/19/1941	

POLLENS - TREES, BOXELDER/MAPLE MIX

boxelder/maple mix injection, solution

Product Information

Product Type	NON-STANDARDIZED ALLERGENIC	Item Code (Source)	NDC:65044-1214
Route of Administration	PERCUTANEOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACER NEGUNDO POLLEN (UNII: P6K070AR8V) (ACER NEGUNDO POLLEN - UNII:P6K070AR8V)	ACER NEGUNDO POLLEN	0.05 g in 1 mL
ACER SACCHARUM POLLEN (UNII: V38QUQ7861) (ACER SACCHARUM POLLEN - UNII:V38QUQ7861)	ACER SACCHARUM POLLEN	0.05 g in 1 mL
ACER RUBRUM POLLEN (UNII: 700NK45C76) (ACER RUBRUM POLLEN - UNII:700NK45C76)	ACER RUBRUM POLLEN	0.05 g in 1 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65044-1214-1	5 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA103888	04/19/1941	

POLLENS - TREES, CEDAR, MOUNTAIN JUNIPERUS ASHEI

cedar, mountain juniperus ashei injection, solution

Product Information

Product Type	NON-STANDARDIZED ALLERGENIC	Item Code (Source)	NDC:65044-1337
Route of Administration	PERCUTANEOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
JUNIPERUS ASHEI POLLEN (UNII: 544F8MEY0Y) (JUNIPERUS ASHEI POLLEN - UNII:544F8MEY0Y)	JUNIPERUS ASHEI POLLEN	0.05 g in 1 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65044-1337-1	5 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA103888	04/19/1941	

POLLENS - TREES, CEDAR, RED JUNIPERUS VIRGINIANA

cedar, red juniperus virginiana injection, solution

Product Information

Product Type	NON-STANDARDIZED ALLERGENIC	Item Code (Source)	NDC:65044-1340
Route of Administration	PERCUTANEOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
JUNIPERUS VIRGINIANA POLLEN (UNII: PY0JA16R2G) (JUNIPERUS VIRGINIANA POLLEN - UNII:PY0JA16R2G)	JUNIPERUS VIRGINIANA POLLEN	0.05 g in 1 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65044-1340-1	5 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA103888	04/19/1941	

Labeler - Jubilant HollisterStier LLC (069263643)**Registrant** - Jubilant HollisterStier LLC (069263643)