MOTION SICKNESS- dimenhydrinate tablet Strategic Sourcing Services LLC

SunMark 44-198 Delisted

Active ingredient (in each tablet)

Dimenhydrinate 50 mg

Purpose

Antiemetic

Uses

for prevention and treatment of these symptoms associated with motion sickness:

- nausea
- vomiting
- dizziness

Warnings

Do not use

for children under 2 years of age unless directed by a doctor.

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers.

When using this product

- alcohol, sedatives, and tranquilizers may increase drowsiness
- marked drowsiness may occur
- use caution when driving a motor vehicle or operating machinery
- avoid alcoholic beverages

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

 to prevent motion sickness, the first dose should be taken one-half to one hour before starting activity

adults and children 12 years and over	1 to 2 tablets every 4-6 hours; do not exceed 8 tablets in 24 hours, or as directed by a doctor
children 6 to under 12 years	$\frac{1}{2}$ to 1 tablet every 6-8 hours; do not exceed 3 tablets in 24 hours, or as directed by a doctor
children 2 to under 6 years	$\frac{1}{2}$ tablet every 6-8 hours; do not exceed $1\frac{1}{2}$ tablets in 24 hours, or as directed by a doctor

Other information

- each tablet contains: calcium 35 mg
- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- protect from moisture
- see end flap for expiration date and lot number

Inactive ingredients

croscarmellose sodium, dibasic calcium phosphate dihydrate, magnesium stearate, microcrystalline cellulose, silicon dioxide, stearic acid

Questions or comments?

1-800-426-9391

Principal Display Panel

sunmark®

*COMPARE TO DRAMAMINE®
ORIGINAL FORMULA ACTIVE INGREDIENT
NDC 70677-0022-1

motion sickness DIMENHYDRINATE 50 mg

Antiemetic

Prevents Nausea, Vomiting & Dizziness for Children & Adults

12 TABLETS

actual size

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

*This product is not manufactured or distributed by Medtech Products Inc., owner of the registered trademark Dramamine® Original Formula. 50844 REV0518A19802

McKESSON

Distributed by McKesson Corporation 6555 State Highway 161 Irving, TX 75039 www.mckesson.com sun mark[®]

*COMPARE TO DRAMAMINE® **ORIGINAL FORMULA ACTIVE INGREDIENT**

motion sickness

DIMENHYDRINATE 50 mg Antiemetic

actual

12 TABLETS

supmark® *compare to dramamine® original formula active ingredient

motion sickness

DIMENHYDRINATE 50 mg **Antiemetic**

Prevents Nausea, **Vomiting & Dizziness** for Children & Adults

12 TABLETS

actual size

ммм.тскеѕѕоп.сот 17 July 1X 75039 6555 State Highway 167 Distributed by McKesson Corporation

Wckesson

REV0518A19802 Dramamine® Original Formula. Medtech Products Inc., owner of the registered trademark I pie broduct is not manufactured or distributed by

B-1242-198-02 REV0518A19802

O

9

 ∞

6

6

MUCSTIONS OF COMMENTS? 1-800-426-9391

microcrystalline cellulose, silicon dioxide, stearic acid calcium phosphate dihydrate, magnesium stearate, INACTIVE INGREDIENTS croscarmellose sodium, dibasic

- see end flap for expiration date and lot number ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ protect from moisture
- OPENED OR BLISTER IS TORN OR BROKEN ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS ■ each tablet contains: calcium 35 mg

Other information

hildren 2 to under years	% tablet every 6-8 hours; do not exceed 1% tablets in 24 hours, or as directed by a doctor
hildren 6 to under 2 years	% to 1 tablet every 6-8 hours; do not exceed 3 tablets in 24 hours, or as directed by a doctor
2 years and over	do not exceed 8 tablets in 24 hours, or as directed by a doctor

adults and children 1 to 2 tablets every 4-6 hours; חוב-וומוו נס חווב ווחחו חבוחוב פומו ווווח מכווגווא

medical help or contact a Poison Control Center right keep out of reach of children. In case of overdose, get

If pregnant or breast-feeding, ask a health professional

- svoid alcoholic beverages ■ nze cantion when driving a motor vehicle or operating qrowsiness ■ marked drowsiness may occur ■ alcohol, sedatives, and tranquilizers may increase
 - When using this product

sedatives or tranquilizers.

Ask a doctor or pharmacist before use if you are taking

- qitticulty in urination due to enlargement of the prostate bronchitis

 glaucoma ■ a breathing problem such as emphysema or chronic
 - Ask a doctor before use if you have

ру а doctor.

Do not use for children under 2 years of age unless directed Warnings

■ nausea ■ vomiting ■ dizziness SSSOCISTED WITH MOTION SICKNESS: USes for prevention and treatment of these symptoms



TAMPER EVIDENT: DO NOT OPENED OR IF BLISTER UNIOR SHOWS ANY SIGNS O NOT USE IF PACKAGE IS ER UNIT IS TORN, BROKEN SIGNS OF TAMPERING

MOTION SICKNESS

dimenhydrinate tablet

Product Information Product Type HUMAN OTC DRUG Item Code (Source)

Route of Administration ORAL

Active	Ingredi	ient/Active	Moiety
--------	---------	-------------	--------

Active ingredient/Active Plotety				
Ingredient Name	Basis of Strength	Strength		
DIMENHYDRINATE (UNII: JB937PER5C) (DIPHENHYDRAMINE - UNII:8GTS82S83M, 8-CHLOROTHEOPHYLLINE - UNII:GE2UA340FM)	DIMENHYDRINATE	50 mg		

NDC:70677-0022

Inactive Ingredients	
Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics				
Color	white	Score	2 pieces	
Shape	ROUND	Size	9mm	
Flavor		Imprint Code	44;198	
Contains				

P	Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:70677- 0022-1	2 in 1 CARTON	12/01/1992	05/08/2025		
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product				

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
OTC Monograph Drug	M009	12/01/1992	05/08/2025		
OTC Monograph Drug	M009	12/01/1992	05/08/2025		

Labeler - Strategic Sourcing Services LLC (116956644)

Establishment				
Name	Address	ID/FEI	Business Operations	
LNK International, Inc.		038154464	pack(70677-0022)	

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(70677-0022) , pack(70677-0022)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(70677-0022)

Revised: 10/2023 Strategic Sourcing Services LLC