

**ACETAMINOPHEN- acetaminophen tablet, coated**  
**Bonita Pharmaceuticals LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Extra Strength Acetaminophen 500mg Caplets**

***Active Ingredient***

***(in each caplet)***

Acetaminophen 500mg

***Purposes***

Pain reliever/Fever reducer

***Uses***

For the temporary relief of minor aches and pains due to:

- Headache
- Muscular aches
- Backache
- Minor pain of arthritis
- The common cold
- Toothache
- Premenstrual and menstrual cramps

Temporarily reduces fever.

***Warnings***

**☐Liver warning:**

This product contains acetaminophen. The maximum daily dose of this product is 6 caplets (3,000mg) in 24 hours. Severe liver damage may occur if you take

- more than 4,000mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have**

liver disease.

**Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin.

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.****Overdose warning:**

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

***Directions***

- **Do not take more than directed (see overdose warning)**

**Adults and children 12 years and over:**

- take 2 caplets every 6 hours while symptoms last
- do not take more than 6 caplets in 24 hours unless directed by a doctor
- do not take for more than 10 days unless directed by a doctor

**Children under 12 years:** ask a doctor

***Other Information***

- **Do not use if imprinted safety seal under cap is broken or missing**
- Store at room temperature

***Inactive Ingredients***

- Povidone
- Pregelatinized Starch
- Sodium Starch Glycolate
- Stearic Acid

***Questions?***

If you have any questions or comments, or to report an adverse event, please contact (855) 729-7200.

**PRINCIPAL DISPLAY PANEL**

SEE NEW WARNINGS INFORMATION NDC# 53598-003-10

**Drug Facts**

**Active ingredient (in each caplet)** **Purposes**

Acetaminophen 500 mg ..... Pain reliever/fever reducer

**Uses**

- for the temporary relief of minor aches and pains due to:
  - headache • muscular aches • backache • minor pain of arthritis
  - the common cold • toothache • premenstrual and menstrual cramps
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. The maximum daily dose of this product is 6 caplets (3,000 mg) in 24 hours. Severe liver damage may occur if you take • more than 4,000 mg of acetaminophen in 24 hours • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

**Do not use** • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

• if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have liver disease.**

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.**

**Stop use and ask a doctor if** • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • new symptoms occur • redness or swelling is present

These could be signs of a serious condition.

Lot# / Exp Date:



**EXTRA STRENGTH ACETAMINOPHEN**

**Pain Reliever/Fever Reducer**

**500 MG Each**

Compare to the Active Ingredient in Extra Strength Tylenol®

\*This product is not manufactured or distributed by McNeil Healthcare LLC, distributor of Extra Strength Tylenol®.

**1000 Caplets**

**Drug Facts** (continued)

**if pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In the case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- Do not take more than directed (see overdose warning)
- Adults and children 12 years and over:** • take 2 caplets every 6 hours while symptoms last • do not take more than 6 caplets in 24 hours unless directed by a doctor • do not take for more than 10 days unless directed by a doctor
- Children under 12 years:** ask a doctor

**Other information** • Do not use if imprinted safety seal under cap is broken or missing. • Store at room temperature

**Inactive ingredients** Povidone, Pregelatinized Starch, Sodium Starch Glycolate, Stearic Acid.

**Questions?** If you have any questions or comments, or to report an adverse event, please contact (855) 729-7200.

Rev 03/13

**Distributed By:**  
Bonita Pharmaceuticals LLC  
6380 Commerce Dr. Westland, MI 48185



3 53598 00310 7

**Image 1 : Extra Strength Acetaminophen 500mg, 1000ct caplets**  
**Front Top**

SEE NEW WARNINGS INFORMATION NDC# 53598-003-01

**Drug Facts**

**Active ingredient (in each caplet)** **Purposes**

Acetaminophen 500 mg ..... Pain reliever/fever reducer

**Uses**

- for the temporary relief of minor aches and pains due to:
  - headache • muscular aches • backache
  - minor pain of arthritis • the common cold
  - toothache • premenstrual and menstrual cramps
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. The maximum daily dose of this product is 6 caplets (3,000 mg) in 24 hours. Severe liver damage may occur if you take • more than 4,000 mg of acetaminophen in 24 hours • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

**Do not use** • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Rev 03/13

**Do not use if imprinted safety seal under cap is broken or missing**



**EXTRA STRENGTH ACETAMINOPHEN**

**Pain Reliever/Fever Reducer**

**500 MG Each**

Compare to the Active Ingredient in Extra Strength Tylenol®

\*This product is not manufactured or distributed by McNeil Healthcare LLC, distributor of Extra Strength Tylenol®.

**100 Caplets**

**Drug Facts** (continued on back)

**Do not use if imprinted safety seal under cap is broken or missing**

Rev 03/13

**Do not use if imprinted safety seal under cap is broken or missing**

Lot#

Exp.

PEEL HERE



3 53598 00301 5

FOLD PROOF ALONG DOTTED LINE to simulate an actual printed label layout

**Drug Facts** (continued)

- if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have liver disease.**

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.**

**Stop use and ask a doctor if** • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • new symptoms occur • redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In the case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

**Do not take more than directed (see overdose warning)**

**Adults and children 12 years and over:**

- take 2 caplets every 6 hours while symptoms last
- do not take more than 6 caplets in 24 hours unless directed by a doctor
- do not take for more than 10 days unless directed by a doctor

**Children under 12 years:** ask a doctor

**Other information**

• Store at room temperature

**Inactive ingredients** Povidone, Pregelatinized Starch, Sodium Starch Glycolate, Stearic Acid.

**Questions?** If you have any questions or comments, or to report an adverse event, please contact (855) 729-7200.

**Distributed By:** Bonita Pharmaceuticals LLC, 6380 Commerce Dr., Westland, MI 48185

**Drug Facts** (continued)

**Do not use if imprinted safety seal under cap is broken or missing**

Rev 03/13

**Do not use if imprinted safety seal under cap is broken or missing**

**Back Top**

**Image 2 : Extra Strength Acetaminophen 500mg, 100ct caplets**

**ACETAMINOPHEN**  
acetaminophen tablet, coated

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:53598-003
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
POVIDONE (UNII: FZ989GH94E)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

**Product Characteristics**

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	OVAL ((Capsule Shaped Tablets))	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	GPI;A5
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53598-003-01	100 in 1 BOTTLE, PLASTIC		
2	NDC:53598-003-10	1000 in 1 BOTTLE, PLASTIC		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	04/02/2013	

**Labeler** - Bonita Pharmaceuticals LLC (004219442)**Registrant** - Bonita Pharmaceuticals LLC (004219442)**Establishment**

Name	Address	ID/FEI	Business Operations
Bonita Pharmaceuticals LLC		004219442	label(53598-003)