

NICOTINE- nicotine polacrilex gum, chewing
McKesson Corporation dba SKY Packaging

Drug Facts

Active ingredient (in each chewing piece)

Nicotine polacrilex 2 mg (nicotine)

Purpose

Stop smoking aid

Use

- reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding,

only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- stomach ulcer or diabetes
- history of seizures

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth, teeth, or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness,

diarrhea, weakness and rapid heartbeat

- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets.

Pieces of nicotine gum may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- **if you are under 18 years of age, ask a doctor before use. No studies have been done to show if this product will work for you.**
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- **if you smoke your first cigarette within 30 minutes of waking up,** use 4 mg nicotine gum
- **if you smoke your first cigarette more than 30 minutes after waking up,** use 2 mg nicotine gum according to the following 12 week schedule

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat the process until most of the tingle is gone (about 30 minutes)
- do not eat or drink 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- It is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Other information

- **each piece contains:** calcium 115 mg, sodium 11 mg
- store at 20-25°C (68-77°F)
- protect from light

Inactive ingredients

acesulfame potassium, butylated hydroxytoluene, carnauba wax, flavors, gum base, sodium bicarbonate, sodium carbonate, sorbitol, talc

Questions or comments?

Call **1-877-440-7867**

Principal Display Panel

*Compare to the active ingredient in Nicorette® Gum

Nicotine Gum

Nicotine Polacrilex gum USP, 2 mg (nicotine)

Stop Smoking Aid

Sugar free

ORIGINAL FLAVOR

PIECES 2 MG EACH

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES OF WAKING UP.

If you smoke your first cigarette **WITHIN 30 MINUTES** after waking up, use Nicotine Polacrilex Gum USP, 4 mg

TO INCREASE YOUR SUCCESS IN QUITTING:

- 1.** You must be motivated to quit.
- 2.** Use Enough - Use **at least 9** Nicotine Polacrilex Gum per day during the first six weeks.
- 3.** Use Long Enough - Use Nicotine Polacrilex Gum for the full 12 weeks.
- 4. Use With a Support Program** as directed in the enclosed User's Guide.

To remove the gum, tear off single unit.

Peel off backing starting at corner with loose edge.

Push gum through foil

- **NOT FOR SALE TO THOSE UNDER 18 YEARS OF AGE**
- **PROOF OF AGE REQUIRED**
- **NOT FOR SALE IN VENDING MACHINES OR FROM ANY SOURCE WHERE PROOF OF AGE CANNOT BE VERIFIED**

*Nicorette® is a trademark of GlaxoSmithKline Consumer Healthcare, L.P.

This product is not affiliated with GlaxoSmithKline Consumer Healthcare, L.P.

See inside for more details or visit www.nitthehabit.com

BLISTER PACKAGED FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN

Distributed By:

McKesson Corporation dba Sky Packaging

4971 Southridge Blvd. Suite 101, Memphis, TN 38141

Package Label

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. **Use Enough** - Chew at least 9 pieces of Nicotine Polacrilex Gum per day during the first six weeks.
3. **Use Long Enough** - Use Nicotine Polacrilex Gum for the full 12 weeks.
4. **Use With a Support Program** as directed in the enclosed User's Guide.



To remove the gum,
tear off single unit.



Peel off backing
starting at corner
with loose edge.



Push gum through
foil.

NDC 63739-371-63

*Compare to the active ingredient in Nicorette® Gum



Nicotine Gum

Nicotine polacrilex gum USP, 2 mg (nicotine)
Stop Smoking Aid

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE
THAN 30 MINUTES AFTER WAKING UP

If you smoke your first cigarette
WITHIN 30 MINUTES of waking up,
use Nicotine Polacrilex Gum USP, 4 mg

ORIGINAL FLAVOR

110 Pieces

Sugar Free

2
mg

Actual Size

Nicotine Gum

Nicotine polacrilex gum USP, 2 mg (nicotine), Stop Smoking Aid

ORIGINAL FLAVOR

110 Pieces

Sugar Free

2
mg

Color		white	Score		no score
Shape		SQUARE	Size		14mm
Flavor			Imprint Code		
Contains					
Packaging					
#	Item Code	Package Description		Marketing Start Date	Marketing End Date
1	NDC:63739-371-63	110 in 1 CARTON		04/27/2020	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product			
Marketing Information					
Marketing Category		Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANDA		ANDA074507		04/27/2020	

Labeler - McKesson Corporation dba SKY Packaging (140529962)

Revised: 11/2022

McKesson Corporation dba SKY Packaging