

EQUALINE IBUPROFEN- ibuprofen tablet, coated
Supervalu Inc

SuperValu Inc. Ibuprofen Caplets Drug Facts

Active ingredient (in each caplet)

Ibuprofen 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- minor pain of arthritis
- toothache
- backache
- the common cold
- menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chances are higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen,

or others]

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- do not take more than directed
- the smallest effective dose should be used

adults and children 12 years and older	<ul style="list-style-type: none">• take 1 caplet every 4 to 6 hours while symptoms persist• if pain or fever does not respond to 1 caplet, 2 caplets may be used• do not exceed 6 caplets in 24 hours, unless directed by a doctor
children under 12 years	<ul style="list-style-type: none">• ask a doctor

Other information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)
- avoid high humidity and excessive heat above 40°C (104°F)
- see end panel for lot number and expiration date

Inactive ingredients

colloidal silicon dioxide, corn starch, croscarmellose sodium, hypromellose, iron oxide red, iron oxide yellow, microcrystalline cellulose, polyethylene glycol, polysorbate 80, stearic acid, titanium dioxide

Questions or comments?

1-877-932-7948

Principal Display Panel

compare to Advil® active ingredient

ibuprofen caplets**

ibuprofen tablets 200mg

pain reliever / fever reducer

(NSAID)

actual size

50 coated caplets**

**capsule-shaped tablets



EQUALINE

compare to Advil®
active ingredient***
NDC 41163-647-71

ibuprofen caplets**
ibuprofen tablets 200mg
pain reliever/fever reducer
(NSAID)



actual size




50 coated caplets**
**capsule-shaped tablets

Actual Size

THIS LABEL DOES NOT CONTAIN FULL PRODUCT INFORMATION. SEE CARTON FOR COMPLETE INFORMATION. READ ALL WARNINGS.

EQUALINE
NDC 41163-647-71
ibuprofen caplets**
ibuprofen tablets 200mg
pain reliever/fever reducer
(NSAID)
50 coated caplets**
**capsule-shaped tablets

Compare to Advil
DO NOT
GLU
Dial
SUP
Ede
MIN




EXP. LOT NO.

7

853323



DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING



Drug Facts

Active ingredient (in each caplet) **Purposes**
Ibuprofen 200 mg (NSAID)*.....Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to:
■ headache ■ muscular aches ■ minor pain of arthritis
■ toothache ■ backache ■ the common cold ■ menstrual cramps
■ temporarily reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives ■ facial swelling ■ asthma (wheezing)
■ shock ■ skin reddening ■ rash ■ blisters
If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chances are higher if you
■ are age 60 or older ■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

Ask a doctor before use if ■ you have problems or serious side effects from taking pain relievers or fever reducers
■ the stomach bleeding warning applies to you
■ you have a history of stomach problems, such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you have asthma ■ you are taking a diuretic ▶

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug

When using this product ■ take with food or milk if stomach upset occurs ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if
■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood ■ have bloody or black stools
■ have stomach pain that does not get better
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present in the painful area
■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions ■ do not take more than directed ■ the smallest effective dose should be used

adults and children 12 years and older	■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor
children under 12 years	■ ask a doctor ▼

Drug Facts (continued)

Other information ■ read a streptococcal throat culture report
■ avoid high humidity and exercise
■ see end panel for lot number

Inactive ingredients colic cross-carmellose sodium, hypromellose, microcrystalline cellulose, stearic acid, titanium dioxide

Questions or comments:

KEEP CARTON

GLUTEN FREE

***This product is not manufactured or registered in the United States.
Distributed by S
Eden Prairie, MN
Contact us at 1-800-368-2777
www.supervali.com

Read all warnings and directions before use.

Store at controlled room temperature (20° to 25°C) (68° to 77°F) and excursions permitted to 15° to 30° (59° to 86°).

Contains: silicon dioxide, corn starch, croscarmellose sodium, hypromellose, ferric oxide red, ferric oxide yellow, polyethylene glycol, poly sorbate 80, titanium dioxide.

1-877-932-7948

FOR REFERENCE

Not distributed by Wyeth LLC,
a Division of
SUPERVALU INC.
MN 55344 USA
1-877-932-7948, or
www.superbrands.com

EQUALINE IBUPROFEN

ibuprofen tablet, coated

Product Information

Product Type	HUMAN OTC DRUG LABEL	Item Code (Source)	NDC:41163-647
Route of Administration	ORAL	DEA Schedule	

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (IBUPROFEN)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE	
STARCH, CORN	
CROSCARMELOSE SODIUM	
HYPROMELLOSES	
FERRIC OXIDE RED	
FERRIC OXIDE YELLOW	
CELLULOSE, MICROCRYSTALLINE	
STEARIC ACID	
TITANIUM DIOXIDE	
POLYETHYLENE GLYCOLS	
POLYSORBATE 80	

Product Characteristics

Color	BROWN	Score	no score
Shape	CAPSULE	Size	15mm
Flavor		Imprint Code	I2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
---	-----------	---------------------	----------------------	--------------------

1	NDC:41163-647-71	1 in 1 CARTON		
1		50 in 1 BOTTLE		
2	NDC:41163-647-78	1 in 1 CARTON		
2		100 in 1 BOTTLE		
3	NDC:41163-647-76	1 in 1 CARTON		
3		120 in 1 BOTTLE		
4	NDC:41163-647-72	1 in 1 CARTON		
4		60 in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA072096	08/17/2004	

Labeler - Supervalu Inc (006961411)

Revised: 12/2014

Supervalu Inc