SUNMARK COUGH DM- dextromethorphan hydrobromide suspension McKesson

McKesson Cough DM Drug Facts

Active ingredient (in each 5 mL)

Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide, USP

Purpose

Cough suppressant

Uses

temporarily relieves

- cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the impulse to cough to help you get to sleep

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Allergy Alert: Contains sodium metabisulfite, a sulfite that may cause allergic-type reactions.

Ask a doctor before use if you have

- chronic cough that lasts as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

Stop use and ask a doctor if

- side effects occur. You may report side effects to FDA at 1-800-FDA-1088.
- cough lasts more than 7 days, cough comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away 1-800-222-1222.

Directions

- shake bottle well before use
- measure only with dosing cup provided. Do not use dosing cup with other products.
- dose as follows or as directed by doctor
- mL = milliliter

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24 hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24 hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24 hours
children under 4 years of age	do not use

Other information

- each 5 mL contains: sodium 5 mg
- store at 20° to 25°C (68° to 77°F)
- dosing cup provided

Inactive Ingredients

D&C Red #30 aluminum lake, D&C Yellow #10 aluminum lake, flavor, glycerin, high fructose corn syrup, methylparaben, polysorbate 80, polyvinyl acetate, povidone, propylparaben, purified water, sodium metabisulfite, sodium polystyrene sulfonate, sucrose, tartaric acid, tragacanth gum, triacetin, xanthan gum

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

COMPARE TO DELSYM® ACTIVE INGREDIENT

cough DM

Dextromethorphan Polistirex

Extended-Release Oral Suspension

Cough Suppressant

Contains sodium metabisulfite, a sulfite that may cause allergic-type reactions

Dosing cup included

Orange-flavored liquid

Alcohol-Free

12 HOUR COUGH RELIEF

GLUTEN FREE

sun mark[®]

TAMPER EVIDENT:

Do not use if carton is opened or neckband imprinted with "sealed for your protection" is broken or missing.



COMPARE TO DELSYM® ACTIVE INGREDIENT* NDC 49348-315-84

cough DM

Dextromethorphan Polistirex
Extended-Release Oral Suspension
Cough Suppressant

Contains sodium metabisulfite, a sulfite that may cause allergic-type reactions Dosing cup included Orange-flavored liquid Alcohol-Free

12 HOUR COUGH RELIEF

GLUTEN FREE

3 FL OZ (89 mL)



ACTUAL SIZE





NDC 49348-315-84

cough DM

Dextromethorphan Polistirex Extended-Release Oral Suspension Cough Suppressant

> Contains no fever reducer or pain reliever Dosing cup included Orange-flavored liquid Alcohol-Free

12 HOUR COUGH RELIEF

3 FL OZ (89 mL)



MACHINECON




MEKESSUN

Another Quality Product
Distributed By McKesson
One Post Street, San Francisco, CA 94104
Money Back Guarantee
Please visit us at www.sunmarkbrand.com

LOT NO.

EXP.

: 38451 21 (1



Drug Facts

Active ingredient (in each 5 mL)

Purpose

Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide, USP.....

...Cough suppressant

Uses temporarily relieves

- cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the impulse to cough to help you get to sleep

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Allergy Alert: Contains sodium metabisulfite, a sulfite that may cause allergic-type reactions.

Ask a doctor before use if you have

chronic cough that lasts as occurs with smoking, asthma or emphysema cough that occurs with too much phiegm (mucus)

Stop use and ask a doctor if

- side effects occur. You may report side effects to FDA at 1-800-FDA-1088.
- cough lasts more than 7 days, cough comes back, or occurs with fever, rash
 or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away 1-800-222-1222.

Directions shake bottle well before use

- measure only with dosing cup provided. Do not use dosing cup with other products.
- dose as follows or as directed by doctor
- mL = milliliter

adults and children	10 mL every 12 hours,
12 years of age and over	not to exceed 20 mL in 24 hours
children 6 to under	5 mL every 12 hours,
12 years of age	not to exceed 10 mL in 24 hours
children 4 to under	2.5 mL every 12 hours,
6 years of age	not to exceed 5 mL in 24 hours
children under 4 years of age	do not use

Other information

■ store at 20° to 25°C (68° to 77°F)

meach 5 mL contains: sodium 5 mg

dosing cup provided

Drug Facts (continued)

Inactive ingredients D&C Red #30 aluminum lake, D&C Yeliow #10 aluminum lake, flavor, glycerin, high fructose com syrup, methylparaben, polysorbate 80, polyvinyl acetate, povidone, propylparaben, purified water, sodium metabisulfite, sodium polystyrene sulfonate, sucrose, tartaric acid, tragacanth gum, triacetin, xanthan gum

Questions or comments? 1-800-719-9260

*This product is not manufactured or distributed by Reckitt Benckiser Inc., distributor of Delsym®.

DOSING

SHAKE WELL BEFORE USE.

Measure only with dosing cup provided.

Do not use dosing cup with other products.

mL = milliliter

Age (yr)	Dose		
12 years	10 mL		
to adult	EVERY 12 HOURS		
6 to	5 mL		
under 12	EVERY 12 HOURS		
4 to	2.5 mL		
under 6	EVERY 12 HOURS		
Under 4	Do not use		

See back panel for full dosing directions.




PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org







SUNMARK COUGH DM

dextromethorphan polistirex suspension

Product Information

Product TypeHUMAN OTC DRUG LABELItem Code (Source)NDC:49348-315

Route of Administration ORAL DEA Schedule

Active Ingredient/Active Moiety

Ingredient NameBasis of StrengthStrengthDEXTROMETHORPHAN HYDROBROMIDE (DEXTROMETHORPHAN)DEXTROMETHORPHAN HYDROBROMIDE30 mg in 5 mL

Inactive Ingredients			
Ingredient Name	Strength		
POLISTIREX			
HIGH FRUCTO SE CORN SYRUP			
GLYCERIN			
METHYLPARABEN			
POLYSORBATE 80			
POVIDONES			
PROPYLPARABEN			
WATER			
SO DIUM METABISULFITE			
SODIUM POLYSTYRENE SULFONATE			
SUCROSE			
TARTARIC ACID			
TRAGACANTH			
TRIACETIN			
XANTHAN GUM			

Product Characteristics			
Color	ORANGE	Score	
Shape		Size	
Flavor	ORANGE	Imprint Code	
Contains			

Packaging			
# Item Code	Package Description	Marketing Start Date	Marketing End Date
1 NDC:49348-315-84	1 in 1 CARTON		
1	89 mL in 1 BOTTLE		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091135	08/28/2012	

Labeler - McKesson (177667227)

Revised: 11/2012 McKesson