

UP AND UP CHILDRENS ACETAMINOPHEN- acetaminophen suspension **Target Corporation**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Target Corporation Children's Acetaminophen Drug Facts

Active ingredient (in each 5 mL)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to:
 - the common cold
 - flu
 - headache
 - sore throat
 - toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has

liver disease

Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin

When using this product

do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed (see overdose warning)**
- **shake well before using**
- mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL

36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL
72-95	11 years	15 mL

*or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- **each 5 mL contains:** sodium 3 mg
- store at 20-25°C (68-77°F)
- **do not use if printed neckband is broken or missing**

Inactive ingredients

anhydrous citric acid, butylparaben, calcium sulfate, carrageenan, D&C red #33, FD&C red #40, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate

Questions?

Call 1-888-547-7400

Principal Display Panel

Compare to active ingredient in Children's Tylenol® Oral Suspension for ages 2 to 11 years

children's acetaminophen 160 mg per 5 mL

oral suspension

fever reducer/pain reliever

ibuprofen free

alcohol free

aspirin free

paraben free

up & up™

BUBBLE GUM FLAVOR

DOSAGE CUP INCLUDED

AGES 2-11 YEARS

4 FL OZ (118 mL)



Drug Facts

Active ingredient (in each 5 mL)
Acetaminophen 160 mg.....Pain reliever/fever reducer

Purpose

Uses temporarily:

- reduces fever
- relieves minor aches and pains due to:
 - the common cold
 - flu
 - headache
 - sore throat
 - toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes more than 5 doses in 24 hours, which is the maximum daily amount

- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has liver disease

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

When using this product do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see overdose warning)
- shake well before using
- mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL
72-95	11 years	15 mL

*or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- each 5 mL contains sodium 3 mg
- store at 20-25°C (68-77°F)
- do not use if printed neckband is broken or missing

Inactive ingredients

anhydrous citric acid, calcium sulfate, carrageenan, D&C red #33, FD&C red #40, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate

Questions? Call 1-888-547-7400

**This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Children's Tylenol® Oral Suspension.

007 00 0121 R02
Distributed by Target Corporation
Minneapolis, MN 55403
©2022 Target Brands, Inc. Shop Target.com

GLUTEN FREE

children's
acetaminophen
160 mg/5 mL
oral suspension
fever reducer/pain reliever



NDC 11673-105-26

Compare to active ingredient
in Children's Tylenol®
Oral Suspension**
for ages 2 to 11 years

children's
acetaminophen
160 mg per 5 mL
oral suspension
fever reducer/pain reliever

ibuprofen free
alcohol free
aspirin free
paraben free



4 FL OZ (118 mL)



children's
acetaminophen
160 mg per 5 mL
oral suspension
fever reducer/pain reliever



DIRECTIONS:
STEP 1.
Shake well
before using



STEP 2.
Remove child
protective
cap



STEP 3.
Squeeze your
child's dose
into the
dosing cup



IMPORTANT:
Please see dosing chart or ask
a doctor for your child's correct dose.



AD457 UW C2

UP AND UP CHILDRENS ACETAMINOPHEN

acetaminophen suspension

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:11673-105

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CALCIUM SULFATE, UNSPECIFIED FORM (UNII: WAT0DDB505)	
CARRAGEENAN (UNII: 5C69YCD2YJ)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TWG28D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K679OBS311)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0K00R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SODIUM PHOSPHATE, TRIBASIC (UNII: A752Q30A6X)	

Product Characteristics

Color	RED (opaque)	Score	
Shape		Size	
Flavor	BUBBLE GUM	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-105-26	1 in 1 CARTON	07/08/2009	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:11673-105-28	1 in 1 CARTON	08/24/2012	07/14/2014
2		148 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

OTC monograph not
final

part343

07/08/2009

Labeler - Target Corporation (006961700)

Revised: 9/2023

Target Corporation