

**LEUCOVORIN- leucovorin calcium injection, powder, for solution**  
**Sagent Pharmaceuticals**

-----  
**Leucovorin Calcium for Injection**  
**(For IV or IM Use)**


SAGENT™

R<sub>x</sub> only

**DESCRIPTION**

Leucovorin is one of several active, chemically reduced derivatives of folic acid. It is useful as an antidote to drugs which act as folic acid antagonists.

Also known as folinic acid, Citrovorum factor, or 5-formyl-5,6,7,8-tetrahydrofolic acid, this compound has the chemical designation of Calcium *N*-[*p*-[[[(6*RS*)-2-amino-5-formyl-5,6,7,8-tetrahydro-4-hydroxy-6-pteridinyl]methyl]amino]benzoyl]-*L*-glutamate (1:1). The structural formula of leucovorin calcium is:



C<sub>20</sub>H<sub>21</sub>CaN<sub>7</sub>O<sub>7</sub>

M.W.=511.51

Leucovorin calcium for injection is a sterile product indicated for intramuscular (IM) or intravenous (IV) administration and is supplied in 50 mg, 100 mg, 200 mg, and 350 mg vials.

Each 50 mg vial of leucovorin calcium for injection, when reconstituted with 5 mL of sterile diluent, contains leucovorin (as the calcium salt) 10 mg/mL.

Each 100 mg vial of leucovorin calcium for injection, when reconstituted with 10 mL of sterile diluent, contains leucovorin (as the calcium salt) 10 mg/mL.

Each 200 mg vial of leucovorin calcium for injection, when reconstituted with 20 mL of sterile diluent, contains leucovorin (as the calcium salt) 10 mg/mL.

Each 350 mg vial of leucovorin calcium for injection, when reconstituted with 17.5 mL of sterile diluent, contains leucovorin (as the calcium salt) 20 mg/mL.

In each dosage form, one milligram of leucovorin calcium contains 0.002 mmol of leucovorin and 0.002 mmol of calcium.

These lyophilized products contain no preservative. The inactive ingredient is Sodium Chloride, USP, added to adjust tonicity. Sodium hydroxide and / or hydrochloric acid may be used to adjust the pH. The pH is between 6.5 and 8.5. Reconstitute with Bacteriostatic Water for Injection, USP, which contains benzyl alcohol (see **WARNINGS** section), or with Sterile Water for Injection, USP.

## CLINICAL PHARMACOLOGY

Leucovorin is a mixture of the diastereoisomers of the 5-formyl derivative of tetrahydrofolic acid (THF). The biologically active compound of the mixture is the (-)-*l*-isomer, known as Citrovorum factor, or (-)-folinic acid. Leucovorin does not require reduction by the enzyme dihydrofolate reductase in order to participate in reactions utilizing folates as a source of "one-carbon" moieties. *l*-Leucovorin (*l*-5-formyltetrahydrofolate) is rapidly metabolized (via 5,10-methenyltetrahydrofolate then 5,10-methylenetetrahydrofolate) to *l*-5-methyltetrahydrofolate. *l*-5-Methyltetrahydrofolate can in turn be metabolized via other pathways back to 5,10-methylenetetrahydrofolate, which is converted to 5-methyltetrahydrofolate by an irreversible, enzyme catalyzed reduction using the cofactors FADH<sub>2</sub> and NADPH.

Administration of leucovorin can counteract the therapeutic and toxic effects of folic acid antagonists such as methotrexate, which act by inhibiting dihydrofolate reductase.

In contrast, leucovorin can enhance the therapeutic and toxic effects of fluoropyrimidines used in cancer therapy, such as 5-fluorouracil. Concurrent administration of leucovorin does not appear to alter the plasma pharmacokinetics of 5-fluorouracil. 5-Fluorouracil is metabolized to fluorodeoxyuridylic acid, which binds to and inhibits the enzyme thymidylate synthase (an enzyme important in DNA repair and replication).

Leucovorin is readily converted to another reduced folate, 5,10-methylenetetrahydrofolate, which acts to stabilize the binding of fluorodeoxyuridylic acid to thymidylate synthase and thereby enhances the inhibition of this enzyme.

The pharmacokinetics after intravenous, intramuscular and oral administration of a 25 mg dose of leucovorin were studied in male volunteers. After intravenous administration, serum total reduced folates (as measured by *Lactobacillus casei* assay) reached a mean peak of 1259 ng/mL (range 897 to 1625). The mean time to peak was 10 minutes. This initial rise in total reduced folates was primarily due to the parent compound 5-formyl-THF (measured by *Streptococcus faecalis* assay) which rose to 1206 ng/mL at 10 minutes. A sharp drop in parent compound followed and coincided with the appearance of the active metabolite 5-methyl-THF which became the predominant circulating form of the drug.

The mean peak of 5-methyl-THF was 258 ng/mL and occurred at 1.3 hours. The terminal half-life for total reduced folates was 6.2 hours. The area under the concentration versus time curves (AUCs) for *l*-leucovorin, *d*-leucovorin and 5-methyltetrahydrofolate were  $28.4 \pm 3.5$ ,  $956 \pm 97$  and  $129 \pm 12$  (mg/min/L  $\pm$  S.E.). When a higher dose of *d,l*-leucovorin (200 mg/m<sup>2</sup>) was used, similar results were obtained. The *d*-isomer persisted in plasma at concentrations greatly exceeding those of the *l*-isomer.

After intramuscular injection, the mean peak of serum total reduced folates was 436 ng/mL (range 240 to 725) and occurred at 52 minutes. Similar to IV administration, the initial sharp rise was due to the parent compound. The mean peak of 5-formyl-THF was 360 ng/mL and occurred at 28 minutes. The level of the metabolite 5-methyl-THF increased subsequently over time until at 1.5 hours it represented 50% of the circulating total folates. The mean peak of 5-methyl-THF was 226 ng/mL at 2.8 hours. The terminal half-life of total reduced folates was 6.2 hours. There was no difference of statistical significance between IM and IV administration in the AUC for total reduced folates, 5-formyl-THF, or 5-methyl-THF.

After oral administration of leucovorin reconstituted with aromatic elixir, the mean peak concentration of serum total reduced folates was 393 ng/mL (range 160 to 550). The mean time to peak was 2.3 hours and the terminal half-life was 5.7 hours. The major component was the metabolite 5-methyltetrahydrofolate to which leucovorin is primarily converted in the intestinal mucosa. The mean peak of 5-methyl-THF was 367 ng/mL at 2.4 hours. The peak level of the parent compound was 51 ng/mL at 1.2 hours. The AUC of total reduced folates after oral administration of the 25 mg dose was 92% of the AUC after intravenous administration.

Following oral administration, leucovorin is rapidly absorbed and expands the serum pool of reduced folates. At a dose of 25 mg, almost 100% of the *l*-isomer but only 20% of the *d*-isomer is absorbed. Oral absorption of leucovorin is saturable at doses above 25 mg. The apparent bioavailability of leucovorin was 97% for 25 mg, 75% for 50 mg, and 37% for 100 mg.

In a randomized clinical study conducted by the Mayo Clinic and the North Central Cancer Treatment Group (Mayo/NCCTG) in patients with advanced metastatic colorectal cancer three treatment regimens were compared: Leucovorin (LV) 200 mg/m<sup>2</sup> and 5-fluorouracil (5-FU) 370 mg/m<sup>2</sup> versus LV 20 mg/m<sup>2</sup> and 5-FU 425 mg/m<sup>2</sup> versus 5-FU 500 mg/m<sup>2</sup>. All drugs were administered by slow intravenous infusion daily for 5 days repeated every 28 to 35 days. Response rates were 26% ( $p = 0.04$  versus 5-FU alone), 43% ( $p = 0.001$  versus 5-FU alone) and 10% for the high dose leucovorin, low dose leucovorin and 5-FU alone groups respectively. Respective median survival times were 12.2 months ( $p = 0.037$ ), 12 months ( $p = 0.050$ ), and 7.7 months. The low dose LV regimen gave a statistically significant improvement in weight gain of more than 5%, relief of symptoms, and improvement in performance status. The high dose LV regimen gave a statistically significant improvement in performance status and trended toward improvement in weight gain and in relief of symptoms but these were not statistically significant.

In a second Mayo/NCCTG randomized clinical study the 5-FU alone arm was replaced by a regimen of sequentially administered methotrexate (MTX), 5-FU, and LV. Response rates with LV 200 mg/m<sup>2</sup> and 5-FU 370 mg/m<sup>2</sup> versus LV 20 mg/m<sup>2</sup> and 5-FU 425 mg/m<sup>2</sup> versus sequential MTX and 5-FU and LV were respectively 31% ( $p < 0.01$ ), 42% ( $p < 0.01$ ), and 14%. Respective median survival times were 12.7 months ( $p < 0.04$ ), 12.7 months ( $p < 0.01$ ), and 8.4 months. No statistically significant difference in weight gain of more than 5% or in improvement in performance status was seen between the treatment arms.

The pharmacokinetics of 200 mg doses of Leucovorin administered intravenously and orally (reconstituted powder, not tablets) have been evaluated. The serum clearance corrected for bioavailability, terminal half-life, and apparent volume of distribution of total folate were not significantly different between the routes of administration. The oral bioavailability of the 200 mg dose was 31%. Eighty-three percent of the biologically active IV dose was recovered in the urine within 24 hours, 31% as 5-methyltetrahydrofolate. Twenty percent of the same oral dose was excreted in 24 hours, 16% as 5-methyltetrahydrofolate.<sup>1</sup>

## **INDICATIONS AND USAGE**

Leucovorin calcium rescue is indicated after high-dose methotrexate therapy in osteosarcoma. Leucovorin calcium is also indicated to diminish the toxicity and counteract the effects of impaired methotrexate elimination and of inadvertent overdosages of folic acid antagonists.

Leucovorin calcium is indicated in the treatment of megaloblastic anemias due to folic acid deficiency when oral therapy is not feasible.

Leucovorin is also indicated for use in combination with 5-fluorouracil to prolong survival in the palliative treatment of patients with advanced colorectal cancer.

Leucovorin should not be mixed in the same infusion as 5-fluorouracil because a precipitate may form.

## **CONTRAINDICATIONS**

Leucovorin is improper therapy for pernicious anemia and other megaloblastic anemias secondary to the lack of vitamin B<sub>12</sub>. A hematologic remission may occur while neurologic manifestations continue to progress.

## **WARNINGS**

In the treatment of accidental overdoses of folic acid antagonists, intravenous leucovorin should be administered as promptly as possible. As the time interval between antifolate administration [e.g., methotrexate (MTX)] and leucovorin rescue increases, leucovorin's effectiveness in counteracting toxicity decreases. In the treatment of accidental overdoses of intrathecally administered folic acid antagonists, do not administer leucovorin intrathecally. LEUCOVORIN MAY BE HARMFUL OR FATAL IF GIVEN INTRATHECALLY.

Monitoring of the serum MTX concentration is essential in determining the optimal dose and duration of treatment with leucovorin.

Delayed MTX excretion may be caused by a third space fluid accumulation (i.e., ascites, pleural effusion), renal insufficiency, or inadequate hydration. Under such circumstances, higher doses of leucovorin or prolonged administration may be indicated. Doses higher than those recommended for oral use must be given intravenously.

Because of the benzyl alcohol contained in certain diluents used for Leucovorin Calcium for Injection, when doses greater than 10 mg/m<sup>2</sup> are administered, Leucovorin Calcium for Injection should be reconstituted with Sterile Water for Injection, USP, and used immediately. (See **DOSAGE AND ADMINISTRATION**.)

Because of the calcium content of the leucovorin solution, no more than 160 mg of leucovorin should be injected intravenously per minute (16 mL of a 10 mg/mL, or 8 mL of a 20 mg/mL solution per minute).

Leucovorin enhances the toxicity of 5-fluorouracil. When these drugs are administered concurrently in the palliative therapy of advanced colorectal cancer, the dosage of 5-fluorouracil must be lower than usually administered. Although the toxicities observed in patients treated with the combination of leucovorin plus 5-fluorouracil are qualitatively similar to those observed in patients treated with 5-fluorouracil alone, gastrointestinal toxicities (particularly stomatitis and diarrhea) are observed more commonly and may be more severe and of prolonged duration in patients treated with the combination.

In the first Mayo/NCCTG controlled trial, toxicity, primarily gastrointestinal, resulted in 7% of patients requiring hospitalization when treated with 5-fluorouracil alone or 5-fluorouracil in combination with 200 mg/m<sup>2</sup> of leucovorin and 20% when treated with 5-fluorouracil in combination with 20 mg/m<sup>2</sup> of leucovorin. In the second Mayo/NCCTG trial, hospitalizations related to treatment toxicity also appeared to occur more often in patients treated with the low dose leucovorin/5-fluorouracil combination than in patients treated with the high dose combination — 11% versus 3%. Therapy with leucovorin/5-fluorouracil must not be initiated or continued in patients who have symptoms of gastrointestinal toxicity of any severity, until those symptoms have completely resolved. Patients with diarrhea must be monitored with particular care until the diarrhea has resolved, as rapid clinical deterioration leading to death can occur. In an additional study utilizing higher weekly doses of 5-FU and leucovorin, elderly and/or debilitated patients were found to be at greater risk for severe gastrointestinal toxicity.

Seizures and/or syncope have been reported rarely in cancer patients receiving leucovorin, usually in association with fluoropyrimidine administration, and most commonly in those with CNS metastases or other predisposing factors, however, a causal relationship has not been established.

The concomitant use of leucovorin with trimethoprim-sulfamethoxazole for the acute treatment of *Pneumocystis carinii* pneumonia in patients with HIV infection was associated with increased rates of treatment failure and morbidity in a placebo-controlled study.

## **PRECAUTIONS**

### **General**

Parenteral administration is preferable to oral dosing if there is a possibility that the patient may vomit and not absorb the leucovorin. Leucovorin has no effect on non-hematologic toxicities of MTX, such

as the nephrotoxicity resulting from drug and/or metabolite precipitation in the kidney.

**Since leucovorin enhances the toxicity of fluorouracil, leucovorin/5-fluorouracil combination therapy for advanced colorectal cancer should be administered under the supervision of a physician experienced in the use of antimetabolite cancer chemotherapy. Particular care should be taken in the treatment of elderly or debilitated colorectal cancer patients, as these patients may be at increased risk of severe toxicity.**

### Laboratory Tests

Patients being treated with the leucovorin/5-fluorouracil combination should have a CBC with differential and platelets prior to each treatment. During the first two courses a CBC with differential and platelets has to be repeated weekly and thereafter once each cycle at the time of anticipated WBC nadir. Electrolytes and liver function tests should be performed prior to each treatment for the first three cycles then prior to every other cycle. Dosage modifications of fluorouracil should be instituted as follows, based on the most severe toxicities:

Diarrhea and/or Stomatitis	WBC/mm <sup>3</sup> Nadir	Platelets/mm <sup>3</sup> Nadir	5-FU Dose
Moderate	1,000-1,900	25-75,000	decrease 20%
Severe	< 1,000	< 25,000	decrease 30%

If no toxicity occurs, the 5-fluorouracil dose may increase 10%. Treatment should be deferred until WBC's are 4,000/mm<sup>3</sup> and platelets 130,000/mm<sup>3</sup>. If blood counts do not reach these levels within two weeks, treatment should be discontinued. Patients should be followed up with physical examination prior to each treatment course and appropriate radiological examination as needed. Treatment should be discontinued when there is clear evidence of tumor progression.

### Drug Interactions

Folic acid in large amounts may counteract the antiepileptic effect of phenobarbital, phenytoin and primidone, and increase the frequency of seizures in susceptible pediatric patients.

Preliminary animal and human studies have shown that small quantities of systemically administered leucovorin enter the CSF primarily as 5-methyltetrahydrofolate and, in humans, remain 1 to 3 orders of magnitude lower than the usual methotrexate concentrations following intrathecal administration. However, high doses of leucovorin may reduce the efficacy of intrathecally administered methotrexate.

Leucovorin may enhance the toxicity of 5-fluorouracil. (See **WARNINGS**.)

### Pregnancy

Teratogenic Effects

#### *Pregnancy Category C*

Adequate animal reproduction studies have not been conducted with leucovorin. It is also not known whether leucovorin can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Leucovorin should be given to a pregnant woman only if clearly needed.

### Nursing Mothers

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when leucovorin is administered to a nursing mother.

### Pediatric Use

See **Drug Interactions**.

## Geriatric Use

Clinical studies of Leucovorin Calcium did not show differences in safety or effectiveness between subjects over 65 and younger subjects. Other clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older patients cannot be ruled out. This drug is known to be excreted by the kidney and the risk of toxic reactions to the drug may be greater in patients with impaired renal function. Because elder patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

## ADVERSE REACTIONS

Allergic sensitization, including anaphylactoid reactions and urticaria, has been reported following administration of both oral and parenteral leucovorin. No other adverse reactions have been attributed to the use of leucovorin *per se*.

Table 1 summarizes significant adverse events occurring in 316 patients treated with the leucovorin/5-fluorouracil combinations compared against 70 patients treated with 5-fluorouracil alone for advanced colorectal carcinoma. These data are taken from the Mayo/NCCTG large multicenter prospective trial evaluating the efficacy and safety of the combination regimen.

**TABLE 1 PERCENTAGE OF PATIENTS TREATED WITH LEUCOVORIN/FLUOROURACIL FOR ADVANCED COLORECTAL CARCINOMA REPORTING ADVERSE EXPERIENCES OR HOSPITALIZED FOR TOXICITY**

	(High LV)/5-FU (N=155)		(Low LV)/5-FU (N=161)		5-FU Alone (N=70)	
	Any (%)	Grade 3+ (%)	Any (%)	Grade 3+ (%)	Any (%)	Grade 3+ (%)
Leukopenia	69	14	83	23	93	48
Thrombocytopenia	8	2	8	1	18	3
Infection	8	1	3	1	7	2
Nausea	74	10	80	9	60	6
Vomiting	46	8	44	9	40	7
Diarrhea	66	18	67	14	43	11
Stomatitis	75	27	84	29	59	16
Constipation	3	0	4	0	1	-
Lethargy/Malaise/Fatigue	13	3	12	2	6	3
Alopecia	42	5	43	6	37	7
Dermatitis	21	2	25	1	13	-
Anorexia	14	1	22	4	14	-
Hospitalization for Toxicity	5%		15%		7%	

High LV = Leucovorin 200 mg/m<sup>2</sup>, Low LV = Leucovorin 20 mg/m<sup>2</sup>

Any = percentage of patients reporting toxicity of any severity

Grade 3+ = percentage of patients reporting toxicity of grade 3 or higher

**TABLE 2 GUIDELINES FOR LEUCOVORIN DOSAGE AND ADMINISTRATION DO NOT ADMINISTER LEUCOVORIN INTRATHECALLY**

Clinical Situation	Laboratory Findings	Leucovorin Dosage and Duration
Normal	Serum methotrexate level	15 mg PO, IM, or IV q 6 hours for

Methotrexate Elimination	approximately 10 micromolar at 24 hours after administration, 1 micromolar at 48 hours, and less than 0.2 micromolar at 72 hours.	60 hours (10 doses starting at 24 hours after start of methotrexate infusion).
Delayed Late Methotrexate Elimination	Serum methotrexate level remaining above 0.2 micromolar at 72 hours, and more than 0.05 micromolar at 96 hours after administration.	Continue 15 mg PO, IM, or IV q 6 hours, until methotrexate level is less than 0.05 micromolar.
Delayed Early Methotrexate Elimination and/or Evidence of Acute Renal Injury	Serum methotrexate level of 50 micromolar or more at 24 hours, or 5 micromolar or more at 48 hours after administration, OR; a 100% or greater increase in serum creatinine level at 24 hours after methotrexate administration (e.g., an increase from 0.5 mg/dL to a level of 1 mg/dL or more)	150 mg IV q 3 hours, until methotrexate level is less than 1 micromolar; then 15 mg IV q 3 hours until methotrexate level is less than 0.05 micromolar.

**To report SUSPECTED ADVERSE REACTIONS, contact Sagent Pharmaceuticals, Inc. at 1-866-625-1618 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

## OVERDOSAGE

Excessive amounts of leucovorin may nullify the chemotherapeutic effect of folic acid antagonists.

## DOSAGE AND ADMINISTRATION

### Advanced Colorectal Cancer

Either of the following two regimens is recommended:

1. Leucovorin is administered at 200 mg/m<sup>2</sup> by slow intravenous injection over a minimum of 3 minutes, followed by 5-fluorouracil at 370 mg/m<sup>2</sup> by intravenous injection.
2. Leucovorin is administered at 20 mg/m<sup>2</sup> by intravenous injection followed by 5-fluorouracil at 425 mg/m<sup>2</sup> by intravenous injection.

5-Fluorouracil and leucovorin should be administered separately to avoid the formation of a precipitate.

Treatment is repeated daily for five days. This five-day treatment course may be repeated at 4 week (28-day) intervals, for 2 courses and then repeated at 4 to 5 week (28 to 35 day) intervals provided that the patient has completely recovered from the toxic effects of the prior treatment course.

In subsequent treatment courses, the dosage of 5-fluorouracil should be adjusted based on patient tolerance of the prior treatment course. The daily dosage of 5-fluorouracil should be reduced by 20% for patients who experienced moderate hematologic or gastrointestinal toxicity in the prior treatment course, and by 30% for patients who experienced severe toxicity (see **PRECAUTIONS: Laboratory Tests**). For patients who experienced no toxicity in the prior treatment course, 5-fluorouracil dosages may be increased by 10%. Leucovorin dosages are not adjusted for toxicity.

Several other doses and schedules of leucovorin/5-fluorouracil therapy have also been evaluated in patients with advanced colorectal cancer; some of these alternative regimens may also have efficacy in the treatment of this disease. However, further clinical research will be required to confirm the safety and effectiveness of these alternative leucovorin/5-fluorouracil treatment regimens.

### Leucovorin Rescue After High-Dose Methotrexate Therapy

The recommendations for leucovorin rescue are based on a methotrexate dose of 12 to 15 grams/m<sup>2</sup> administered by intravenous infusion over 4 hours (see methotrexate package insert for full prescribing information).

Leucovorin rescue at a dose of 15 mg (approximately 10 mg/m<sup>2</sup>) every 6 hours for 10 doses starts 24 hours after the beginning of the methotrexate infusion. In the presence of gastrointestinal toxicity, nausea or vomiting, leucovorin should be administered parenterally. Do not administer leucovorin intrathecally.

Serum creatinine and methotrexate levels should be determined at least once daily. Leucovorin administration, hydration, and urinary alkalinization (pH of 7.0 or greater) should be continued until the methotrexate level is below  $5 \times 10^{-8}$  M (0.05 micromolar). The leucovorin dose should be adjusted or leucovorin rescue extended based on the guidelines provided in Table 2.

Patients who experience delayed early methotrexate elimination are likely to develop reversible renal failure. In addition to appropriate leucovorin therapy, these patients require continuing hydration and urinary alkalinization, and close monitoring of fluid and electrolyte status, until the serum methotrexate level has fallen to below 0.05 micromolar and the renal failure has resolved.

Some patients will have abnormalities in methotrexate elimination or renal function following methotrexate administration, which are significant but less severe than the abnormalities described in Table 2. These abnormalities may or may not be associated with significant clinical toxicity. If significant clinical toxicity is observed, leucovorin rescue should be extended for an additional 24 hours (total of 14 doses over 84 hours) in subsequent courses of therapy. The possibility that the patient is taking other medications which interact with methotrexate (e.g., medications which may interfere with methotrexate elimination or binding to serum albumin) should always be reconsidered when laboratory abnormalities or clinical toxicities are observed.

### **Impaired Methotrexate Elimination or Inadvertent Overdosage**

Leucovorin rescue should begin as soon as possible after an inadvertent overdosage and within 24 hours of methotrexate administration when there is delayed excretion (see **WARNINGS**). Leucovorin 10 mg/m<sup>2</sup> should be administered IV, IM, or PO every 6 hours until the serum methotrexate level is less than  $10^{-8}$  M. In the presence of gastrointestinal toxicity, nausea, or vomiting, leucovorin should be administered parenterally. Do not administer leucovorin intrathecally.

Serum creatinine and methotrexate levels should be determined at 24 hour intervals. If the 24 hour serum creatinine has increased 50% over baseline or if the 24 hour methotrexate level is greater than  $5 \times 10^{-6}$  M or the 48 hour level is greater than  $9 \times 10^{-7}$  M, the dose of leucovorin should be increased to 100 mg/m<sup>2</sup> IV every 3 hours until the methotrexate level is less than  $10^{-8}$  M.

Hydration (3 L/d) and urinary alkalinization with sodium bicarbonate solution should be employed concomitantly. The bicarbonate dose should be adjusted to maintain the urine pH at 7.0 or greater.

### **Megaloblastic Anemia Due to Folic Acid Deficiency**

Up to 1 mg daily. There is no evidence that doses greater than 1 mg/day have greater efficacy than those of 1 mg; additionally, loss of folate in urine becomes roughly logarithmic as the amount administered exceeds 1 mg.

Each 50, 100, and 200 mg vial of Leucovorin Calcium for Injection when reconstituted with 5, 10, and 20 mL, respectively, of sterile diluent yields a leucovorin concentration of 10 mg leucovorin per mL. Each 350 mg vial of Leucovorin Calcium for Injection when reconstituted with 17.5 mL of sterile diluent yields a leucovorin concentration of 20 mg leucovorin per mL. Leucovorin Calcium for Injection contains no preservative. Reconstitute the lyophilized vial products with Bacteriostatic Water for Injection, USP, which contains benzyl alcohol, or with Sterile Water for Injection, USP. When reconstituted with Bacteriostatic Water for Injection, USP, the resulting solution must be used within 7 days. If the product is reconstituted with Sterile Water for Injection, USP, use immediately and discard



unused portion.

Because of the benzyl alcohol contained in Bacteriostatic Water for Injection, USP, when doses greater than 10 mg/m<sup>2</sup> are administered Leucovorin Calcium for Injection should be reconstituted with Sterile Water for Injection, USP, and used immediately. (See **WARNINGS**.) Because of the calcium content of the leucovorin solution, no more than 160 mg of leucovorin should be injected intravenously per minute (16 mL of a 10 mg/mL, or 8 mL of a 20 mg/mL solution per minute).

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

Leucovorin should not be mixed in the same infusion as 5-fluorouracil, since this may lead to the formation of a precipitate.

## HOW SUPPLIED

Leucovorin Calcium for Injection is supplied as follows:

<b>NDC</b>	<b>Leucovorin Calcium for Injection</b>	<b>Package Factor</b>
25021-813-10	50 mg Single-Dose Vial	10 vials per carton
25021-814-30	100 mg Single-Dose Vial	1 vial per carton
25021-815-30	200 mg Single-Dose Vial	1 vial per carton
25021-816-30	350 mg Single-Dose Vial	1 vial per carton

## Storage Conditions

Store dry product and reconstituted solution at 20° to 25°C (68° to 77°F); excursions permitted between 15° and 30°C (59° and 86°F). [See USP Controlled Room Temperature.]

**Protect from light.** Retain in carton until time of use.

**Sterile, Nonpyrogenic, Preservative-free.**

**The container closure is not made with natural rubber latex.**

## REFERENCES

- McGuire B, Sia L, Haynes J, et al. "Absorption kinetics of orally administered Leucovorin Calcium," *NCI Monogr* 1987;5:47-56.

SAGENT™

Mfd. for SAGENT Pharmaceuticals

Schaumburg, IL 60195 (USA)

Made in India

©2012 Sagent Pharmaceuticals, Inc.

Revised: September 2012

PACKAGE LABEL – PRINCIPAL DISPLAY PANEL – Vial Label

NDC 25021-813-10

Rx only

Leucovorin Calcium for Injection

50 mg per vial

For IV or IM Use

Must Be Diluted

LYOPHILIZED  
Single-Dose Vial

NDC 25021-813-10 Rx only

**LEUCOVORIN**  
Calcium for Injection

**50 mg\* per vial**

**For IV or IM Use  
Must Be Diluted  
LYOPHILIZED  
Single-Dose Vial**

**SAGENT™**

**Sterile, Nonpyrogenic, Preservative-free.**  
\*Each vial contains: Leucovorin Calcium equivalent to 50 mg leucovorin as the base and 45 mg sodium chloride. Reconstitute with Bacteriostatic Water for Injection, USP (preserved with benzyl alcohol) and use within 7 days, or with Sterile Water for Injection, USP and use immediately and discard unused portion. When reconstituted with 5 mL sterile diluent, each mL of the resulting solution contains Leucovorin Calcium equivalent to 10 mg leucovorin.

**Usual Dosage:** See package insert for dosage information. Store dry product and reconstituted solution at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.]  
**Protect from light.** Retain in carton until time of use.

**Do not use after**

Lot:  
Exp.

(01)00325021813102

**SAGENT™**  
Mfd. for:  
SAGENT Pharmaceuticals  
Schaumburg, IL 60195 (USA)  
Made in India  
©2012 Sagent Pharmaceuticals, Inc.  
Code No.: GUJ/DRUG/1081

2029931

PACKAGE LABEL – PRINCIPAL DISPLAY PANEL – Vial Label

NDC 25021-814-30

Rx only

Leucovorin Calcium for Injection

100 mg per vial

For IV or IM Use

Must Be Diluted

LYOPHILIZED

Single-Dose Vial

NDC 25021-814-30 Rx only

**LEUCOVORIN**  
Calcium for Injection

**100 mg\* per vial**

**For IV or IM Use  
Must Be Diluted  
LYOPHILIZED  
Single-Dose Vial**

**SAGENT™**

**Sterile, Nonpyrogenic, Preservative-free.**  
\*Each vial contains: Leucovorin Calcium equivalent to 100 mg leucovorin as the base and 90 mg sodium chloride. Reconstitute with Bacteriostatic Water for Injection, USP (preserved with benzyl alcohol) and use within 7 days, or with Sterile Water for Injection, USP and use immediately and discard unused portion. When reconstituted with 10 mL sterile diluent, each mL of the resulting solution contains Leucovorin Calcium equivalent to 10 mg leucovorin.

**Usual Dosage:** See package insert for dosage information. Store dry product and reconstituted solution at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.]  
**Protect from light.** Retain in carton until time of use.

**Do not use after**

Lot:  
Exp.

(01)00325021814307

**SAGENT™**  
Mfd. for:  
SAGENT Pharmaceuticals  
Schaumburg, IL 60195 (USA)  
Made in India  
©2012 Sagent Pharmaceuticals, Inc.  
Code No.: GUJ/DRUG/1081

2029933

PACKAGE LABEL – PRINCIPAL DISPLAY PANEL – Vial Label

NDC 25021-815-30

Rx only

Leucovorin Calcium for Injection

200 mg per vial

For IV or IM Use

Must Be Diluted

LYOPHILIZED

Single-Dose Vial

NDC 25021-815-30 Rx only

**LEUCOVORIN**  
Calcium for Injection

**200 mg\* per vial**

For IV or IM Use  
Must Be Diluted  
LYOPHILIZED  
Single-Dose Vial

SAGENT

**Sterile, Nonpyrogenic, Preservative-free.**  
\*Each vial contains: Leucovorin Calcium equivalent to 200 mg leucovorin as the base and 180 mg sodium chloride.  
Reconstitute with Bacteriostatic Water for Injection, USP (preserved with benzyl alcohol) and use within 7 days, or with Sterile Water for Injection, USP and use immediately and discard unused portion. When reconstituted with 20 mL sterile diluent, each mL of the resulting solution contains Leucovorin Calcium equivalent to 10 mg leucovorin.  
**Usual Dosage:** See package insert for dosage information.  
Store dry product and reconstituted solution at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.]  
**Protect from light.** Retain in carton until time of use.  
**Do not use after** \_\_\_\_\_

(01)00325021815304

Lot: \_\_\_\_\_  
Exp: \_\_\_\_\_

SAGENT®  
Mfd. for:  
SAGENT Pharmaceuticals  
Schaumburg, IL 60195 (USA)  
Made in India  
©2012 Sagent Pharmaceuticals, Inc.  
Code No.: GUJ/DRUG/1081

2029935

PACKAGE LABEL – PRINCIPAL DISPLAY PANEL – Vial Label

NDC 25021-816-30

Rx only

Leucovorin Calcium for Injection

350 mg per vial

For IV or IM Use

Must Be Diluted

LYOPHILIZED

Single-Dose Vial

NDC 25021-816-30 Rx only

# LEUCOVORIN

Calcium for Injection

350 mg\* per vial

**For IV or IM Use  
Must Be Diluted  
LYOPHILIZED  
Single-Dose Vial**

**Sterile, Nonpyrogenic, Preservative-free.**  
\*Each vial contains: Leucovorin Calcium equivalent to 350 mg leucovorin and 140 mg sodium chloride.

Reconstitute with Bacteriostatic Water for Injection, USP (preserved with benzyl alcohol) and use within 7 days, or with Sterile Water for Injection, USP and use immediately and discard unused portion. When reconstituted with 17.5 mL sterile diluent, each mL of the resulting solution contains Leucovorin Calcium equivalent to 20 mg leucovorin.

**Usual Dosage:** See package insert for dosage information.  
Store dry product and reconstituted solution at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature.]  
**Protect from light.** Retain in carton until time of use.

**Do not use after**

(01)00325021816301

Lot: \_\_\_\_\_

Exp: \_\_\_\_\_

**SAGENT®**  
Mfd. for:  
SAGENT Pharmaceuticals  
Schaumburg, IL 60195 (USA)  
Made in India  
©2012 Sagent Pharmaceuticals, Inc.  
Code No.: GUJ/DRUG/1081

2029937

## LEUCOVORIN

leucovorin calcium injection, powder, for solution

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG LABEL	<b>Item Code (Source)</b>	NDC:25021-813
<b>Route of Administration</b>	INTRAVENOUS, INTRAMUSCULAR	<b>DEA Schedule</b>	

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
leucovorin calcium (leucovorin)	leucovorin	50 mg in 5 mL

### Inactive Ingredients

Ingredient Name	Strength
sodium chloride	
sodium hydroxide	
hydrochloric acid	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:25021-813-10	10 in 1 CARTON		
1		5 mL in 1 VIAL; Combination Product Type = C112160		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA200753	09/15/2012	

## LEUCOVORIN

leucovorin calcium injection, powder, for solution

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG LABEL	<b>Item Code (Source)</b>	NDC:25021-814
<b>Route of Administration</b>	INTRAVENOUS, INTRAMUSCULAR	<b>DEA Schedule</b>	

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
leucovorin calcium (leucovorin)	leucovorin	100 mg in 10 mL

### Inactive Ingredients

Ingredient Name	Strength
sodium chloride	
sodium hydroxide	
hydrochloric acid	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:25021-814-30	1 in 1 CARTON		
1		10 mL in 1 VIAL; Combination Product Type = C112160		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA200753	09/15/2012	

## LEUCOVORIN

leucovorin calcium injection, powder, for solution

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG LABEL	<b>Item Code (Source)</b>	NDC:25021-815
<b>Route of Administration</b>	INTRAVENOUS, INTRAMUSCULAR	<b>DEA Schedule</b>	

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
leucovorin calcium (leucovorin)	leucovorin	200 mg in 20 mL

### Inactive Ingredients

Ingredient Name	Strength
sodium chloride	
sodium hydroxide	
hydrochloric acid	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:25021-815-30	1 in 1 CARTON		
1		20 mL in 1 VIAL; Combination Product Type = C112160		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA200753	09/15/2012	

## LEUCOVORIN

leucovorin calcium injection, powder, for solution

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG LABEL	Item Code (Source)	NDC:25021-816
Route of Administration	INTRAVENOUS, INTRAMUSCULAR	DEA Schedule	

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
leucovorin calcium (leucovorin)	leucovorin	350 mg in 17.5 mL

### Inactive Ingredients

Ingredient Name	Strength
sodium chloride	
sodium hydroxide	
hydrochloric acid	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:25021-816-30	1 in 1 CARTON		
1		17.5 mL in 1 VIAL; Combination Product Type = C112160		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA200855	09/15/2012	

**Labeler** - Sagent Pharmaceuticals (796852890)

Revised: 9/2014

Sagent Pharmaceuticals