

BUDESONIDE- budesonide spray, metered Walgreens

Drug Facts

Active ingredient (in each spray)

Budesonide (glucocorticoid) 32 mcg

Purpose

Nasal allergy symptom reliever

Uses

Temporarily relieves these symptoms of hay fever or other upper respiratory allergies:

- nasal congestion
- runny nose
- itchy nose
- sneezing

Warnings

Do not use

- in children under 6 years of age
- if you have ever had an allergic reaction to any of the ingredients

Ask a doctor before use if you

- have had recent nose ulcers or nose surgery
- have had a nose injury that has not healed
- are using a steroid medicine for asthma, allergies or skin rash
- have an eye infection
- have or had glaucoma or cataracts

When using this product

- the growth rate of some children may be slower
- some symptoms may get better on the first day of treatment. It may take up to two weeks of daily use to feel the most symptom relief.
- do not share this bottle with anyone else as this may spread germs
- remember to tell your doctor about all the medicines you take, including this one

Stop use and ask a doctor if

- you have, or come into contact with someone who has, chickenpox, measles or tuberculosis
- you have or develop symptoms of an infection such as persistent fever
- you have any change in vision
- you have severe or frequent nosebleeds

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

Read insert (inside package) on how to:

- get a new bottle ready (primed) before first use
- prime bottle again if not used for two days
- use the spray
- clean the spray nozzle

ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER

- adults and children 12 years of age and older
- once daily, spray 2 times into each nostril while sniffing gently
 - once your allergy symptoms improve, reduce to 1 spray in each nostril per day

CHILDREN 6 TO UNDER 12 YEARS OF AGE

- the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year

- children 6 to under 12 years of age
- an adult should supervise use
 - once daily, spray 1 time into each nostril while sniffing gently
 - if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day

- children under 6 years of age
- do not use

- **do not use more than directed**
- if you forget a dose, do **not** double the

next dose

- do not spray into eyes or mouth
 - if allergy symptoms do not improve after two weeks, stop using and talk to a doctor
 - do not use for the common cold
 - shake well before each use
-

Other information

- **do not use if the safety seal labeled "sealed for your protection" is broken or missing.**
- keep package and insert. They contain important information.
- store upright at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Do not freeze. Protect from light.

Inactive ingredients

carboxymethyl cellulose sodium, dextrose anhydrous, edetate disodium dihydrate, hydrochloric acid (for pH adjustment), microcrystalline cellulose, polysorbate 80, potassium sorbate, purified water

Questions or comments?

call toll free **1-800-706-5575**

Principal Display Panel - Carton

CARTON LABEL - PRINCIPAL DISPLAY PANEL - 32 mcg per spray

Walgreens NDC 0363-0048-01

Budesonide Nasal Spray

Nasal Allergy Symptom Reliever

120 sprays

Relief of:

- Nasal Congestion
- Runny Nose
- Itchy Nose
- Sneezing



DO NOT USE IF CAPTIVE SEALS ARE BROKEN.

NG 073E-0

ORIGINAL PRESCRIPTION STRENGTH

NDC 0363-0048-01



24 HOUR Budesonide Nasal Spray

NASAL ALLERGY SYMPTOM RELIEVER 32 mcg PER SPRAY (GLUCOCORTICOID)

24 Hour

Non-Drowsy

- Relief of nasal congestion, runny nose, itchy nose & sneezing

120 METERED SPRAYS

0.285 FL OZ (8.43 mL)



Drug Facts

Active ingredient (in each spray)
Budesonide (glucocorticoid)
32 mcg..... Nasal allergy symptom reliever

Uses
Temporarily relieves these symptoms of hay fever or other upper respiratory allergies:
 ■ nasal congestion ■ runny nose
 ■ itchy nose ■ sneezing

Warnings
 Do not use
 ■ in children under 6 years of age
 ■ if you have ever had an allergic reaction to any of the ingredients

Ask a dealer before use if you
 ■ have had recent nose ulcers or nose surgery
 ■ have had a nose injury that has not healed
 ■ are using a steroid medicine for asthma, allergies or skin rash
 ■ have an eye infection
 ■ have or had glaucoma or cataracts

When using this product
 ■ the growth rate of some children may be slower
 ■ some symptoms may get better on the first day of treatment. It may take up to two weeks of daily use to feel the most symptom relief
 ■ do not share this bottle with anyone else as this may spread germs
 ■ remember to tell your doctor about all the medicines you take, including this one

Stop use and ask a dealer if
 ■ you have, or come into contact with someone who has chikungunya, measles or tuberculosis
 ■ you have or develop symptoms of an infection such as persistent fever

Drug Facts (continued)

■ you have any change in vision
 ■ you have severe or frequent nosebleeds
 If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions
 Read insert (inside package) on how to:
 ■ get a new bottle ready (primed) before first use
 ■ prime bottle again if not used for two days
 ■ use the spray
 ■ clean the spray nozzle

ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER
 adults and children 12 years of age and older
 ■ once daily, spray 2 times into each nostril while sniffing gently
 ■ once your allergy symptoms improve, reduce to 1 spray in each nostril per day

CHILDREN 6 TO UNDER 12 YEARS OF AGE
 ■ the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year.
 children 6 to under 12 years of age
 ■ an adult should supervise use
 ■ once daily, spray 1 time into each nostril while sniffing gently
 ■ if it is necessary, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day

Drug Facts (continued)

children under 6 years of age
 ■ do not use
 ■ do not use more than directed
 ■ if you forget a dose, do not double the next dose
 ■ do not spray into eyes or mouth
 ■ if allergy symptoms do not improve after two weeks, stop using and talk to a doctor
 ■ do not use for the common cold
 ■ shake well before each use

Other information
 ■ do not use if the safety seal is broken or missing.
 ■ keep package and insert. They contain important information.
 ■ store upright at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Do not freeze. Protect from light.

Inactive ingredients
 carboxymethyl cellulose sodium, dextrose anhydrous, edetate disodium dihydrate, hydrochloric acid (for pH adjustment), microcrystalline cellulose, polysorbate 80, potassium sorbate, purified water

Questions or comments?
 call toll free 1-800-706-5575

IMPORTANT: Read all product information before using. Keep package and insert. They contain important information.

Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.
 *This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Rhinocort® Allergy Spray.



17EM 03989 W000-000-0
 5 11917 18320 6
 ©2021 Walgreen Co. MADE IN CANADA
 walgreens.com
 100% SATISFACTION GUARANTEED
 DISTRIBUTED BY: WALGREEN CO.
 200 WILMOT RD., DEERFIELD, IL 60015

W3JORG0821-F

448553

Principal Display Panel - Bottle
BOTTLE LABEL - PRINCIPAL DISPLAY PANEL - 32 mcg per spray
Walgreens NDC 0363-0048-01
Budesonide Nasal Spray
Nasal Allergy Symptom Reliever

120 sprays



BUDESONIDE

budesonide spray, metered

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0048
Route of Administration	NASAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BUDESONIDE (UNII: Q3OKS62Q6X) (Budesonide - UNII:Q3OKS62Q6X)	BUDESONIDE	32 ug

Inactive Ingredients

Ingredient Name	Strength
EDETATE DISODIUM (UNII: 7FLD91C86K)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K679OBS311)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
DEXTRROSE, UNSPECIFIED FORM (UNII: IY9XDZ35W2)	
POTASSIUM SORBATE (UNII: 1VPU26JZZ4)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0048-01	1 in 1 BOTTLE, SPRAY	04/11/2016	
1		120 in 1 BOTTLE, SPRAY; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA078949	04/11/2016	

Labeler - Walgreens (008965063)

Registrant - Apotex Inc. (209429182)

Establishment

Name	Address	ID/FEI	Business Operations
Apotex Inc.		255092496	analysis(0363-0048) , manufacture(0363-0048)

Revised: 11/2023

Walgreens