

**DAYTIME MUCUS RELIEF SEVERE COLD AND NIGHTTIME COLD AND
FLU MAXIMUM STRENGTH- daytime acetaminophen, dextromethorphan hbr,
guaifenesin, phenylephrine hci, nighttime acetaminophen, diphenhydramine
hci, phenylephrine hci
TARGET Corporation**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

DRUG FACTS

Active ingredients for Nighttime (in each 20 mL)

Acetaminophen 650 mg

Diphenhydramine HCl 25 mg

Phenylephrine HCl 10 mg

Active ingredients for Daytime (in each 20 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Guaifenesin 400 mg

Phenylephrine HCl 10 mg

Purpose for Nighttime

Pain reliever/fever reducer

Antihistamine/cough suppressant

Nasal decongestant

Purpose for Daytime

Pain reliever / fever reducer

Cough suppressant

Expectorant

Nasal decongestant

Uses

Nighttime

- temporarily relieves these common cold and flu symptoms

- cough
- nasal congestion
- minor aches and pains
- sore throat
- headache
- runny nose
- sneezing
- temporarily reduces fever
- controls cough to help you get to sleep

Daytime

- temporarily relieves these common cold and flu symptoms
- cough
- nasal congestion
- minor aches and pains
- sore throat
- headache
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make cough more productive

Warnings

NIGHTTIME and DAYTIME

Liver warning: This product contain acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash.

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

Nighttime

- with any drug containing acetaminophen (prescription or nonprescription) . If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other drug containing diphenhydramine, even one used on the skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain

drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product

- for children under 12 years of age

Daytime

- with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI)(certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

Nighttime

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Daytime

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

Nighttime

- you are taking the blood thinning drug warfarin
- you are taking sedative or tranquilizers

Daytime

- taking the blood thinning drug warfarin

When using these products

Nighttime

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Daytime

- **do not use more than directed**

Stop use and ask a doctor if

Nighttime

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse, or lasts more than 7 days
- fever gets worse, or last more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be a signs of a serious condition

Daytime

- nervousness, dizziness or sleeplessness occur
- pain, nasal congestion, or cough gets worse, or lasts more than 7 days
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

Nighttime and DayTime

ask a health professional before use.

Keep out of reach of children.

Nighttime and DayTime

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Nighttime

- **do not take more than directed (see overdose warning)**

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device
- keep dosing cup with product
- mL = milliliter
- dose as follows or as directed by a doctor
- adults and children 12 years and older: 20 mL every 4 hours while symptoms last
- children under 12 years of age: do not use

Daytime

- **do not take more than directed (see Overdose warning)**
- do not take more than 6 doses in any 24-hours period
- measure only with dosing cup provided. Do not use any other dosing device.
- dose as follows or as directed by a doctor
- keep dosing cup with product
- mL = milliliter
- adults and children 12 years of age and older: 20 mL every 4 hours
- children under 12 years of age: do not use

Other information

Nighttime

- **each 20 mL contains:** sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Daytime

- **each 20 mL contains:** 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

Nighttime

citric acid, disodium EDTA, FD&C Blue #1, FD&C red #40, Flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Daytime

citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Principal Display Panel

NIGHTTIME

Compare to active ingredients in Maximum Strength Mucinex® Fast-Max® Night Time Cold & Flu**

maximum strength

nighttime

Cold & Flu

acetaminophen (pain reliever / fever reducer)

diphenhydramine HCl (antihistamine / cough suppressant)

phenylephrine HCl (nasal decongestant)

Relieves aches, fever, sore throat

controls cough

relieves nasal congestion

relieves runny nose and sneezing

AGES 12 + YEARS

DAYTIME

Compare to active ingredients in Maximum Strength Mucinex® Fast-Max® Day Time Severe Cold*

maximum strength

daytime

Severe Cold

acetaminophen (pain reliever / fever reducer)

dextromethorphan HBr (cough suppressant)

guaifenesin (expectorant)

phenylephrine HCl (nasal decongestant)

Relieves aches, fever & sore throat

Controls Cough

relieves nasal and chest congestion

thins and loosens mucus

AGES 12 + YEARS

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by Recitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® Day Time Severe Cold.

**This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® FAST-MAX® Night Time Cold & Flu.

Dist. by Target Corp.

Minneapolis, MN 55403

Questions? Call 1-800-910-6874

[illegible]

TARGET Maximum Strength Nighttime cold and flu Maximum Strength Daytime Severe Cold

daytime acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hci, nighttime acetaminophen, diphenhydramine hci, phenylephrine hci kit

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-463
---------------------	----------------	---------------------------	---------------

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-463-12	1 in 1 KIT; Type 0: Not a Combination Product	01/11/2018	

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE, PLASTIC	177 mL
Part 2	1 BOTTLE, PLASTIC	177 mL

Part 1 of 2

DAYTIME MUCUS RELIEF SEVERE COLD MAXIMUM STRENGTH

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hci liquid

Product Information

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 20 mL

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE CALCIUM DISODIUM (UNII: 25IH6R4SGF)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYL GALLATE (UNII: 8D4SNN7V92)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341		

Part 2 of 2

NIGHTTIME COLD AND FLU MAXIMUM STRENGTH

acetaminophen, diphenhydraminehcl, phenylephrine hcl liquid

Product Information

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 20 mL
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg in 20 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
PROPYL GALLATE (UNII: 8D4SNN7V92)	
XANTHAN GUM (UNII: TTV12P4NEE)	
EDETATE CALCIUM DISODIUM (UNII: 25IH6R4SGF)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

OTC monograph final	part341	01/11/2018	
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	01/11/2018	

Labeler - TARGET Corporation (006961700)

Revised: 11/2022

TARGET Corporation