

**ALLERGY MULTI SYMPTOM- acetaminophen, chlorpheniramine maleate, and phenylephrine hydrochloride tablet, coated**  
**AMERISOURCE BERGEN**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**GNP-1128B-2020-0820**

**Drug Facts**

| <b>Active ingredients<br/>(in each caplet)</b> | <b>Purpose</b>     |
|--|--------------------|
| Acetaminophen 325 mg                           | Pain reliever      |
| Chlorpheniramine maleate<br>2 mg               | Antihistamine      |
| Phenylephrine HCl 5 mg                         | Nasal decongestant |

**Uses**

- temporarily relieves these symptoms of hay fever or other upper respiratory allergies:
  - headache
  - sinus congestion and pressure
  - nasal congestion
  - runny nose and sneezing
  - minor aches and pains
- temporarily relieves these additional symptoms of hay fever:
  - itching of the nose or throat
  - itchy, watery eyes
- helps clear nasal passages
- helps decongest sinus openings and passages

**Warnings**

**Liver warning**

This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma

### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

### **When using this product**

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

### **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.**

## Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## Directions

- do not take more than directed (see overdose warning)

|   |  |
|---|--|
| adults and<br>children 12<br>years and over | <ul style="list-style-type: none"><li>take 2 caplets every 4 hours</li><li>swallow whole; do not crush, chew, or dissolve</li><li>do not take more than 10 caplets in 24 hours</li></ul> |
| children under<br>12 years                  | <ul style="list-style-type: none"><li>ask a doctor</li></ul>   |

## Other information

- store between 20°-25°C (68°-77°F) in a dry place
- retain carton for complete product information and warnings

## Inactive ingredients

acesulfame potassium, colloidal silicon dioxide, corn starch, croscarmellose sodium, crospovidone, flavor, iron oxide yellow, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, stearic acid, talc, titanium dioxide

## PRINCIPAL DISPLAY PANEL

GOOD NEIGHBOR PHARMACY®

NDC 46122-660-62

For Adults

Allergy Multi-Symptom

Acetaminophen, Phenylephrine HCl, Chlorpheniramine Maleate

Pain Reliever, Nasal Decongestant, Antihistamine

Relieves:

- Headache
- Sinus Pressure
- Nasal Congestion

- ## Actual Size



|   |                             |        |
|---|-----------------------------|--------|
| <b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)               | ACETAMINOPHEN               | 325 mg |
| <b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U) | CHLORPHENIRAMINE MALEATE    | 2 mg   |
| <b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV) | PHENYLEPHRINE HYDROCHLORIDE | 5 mg   |

### Inactive Ingredients

| Ingredient Name  | Strength |
|--|----------|
| <b>ACESULFAME POTASSIUM</b> (UNII: 23OV73Q5G9)             |          |
| <b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)                  |          |
| <b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)            |          |
| <b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)              |          |
| <b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)               |          |
| <b>CROSPVIDONE</b> (UNII: 2S7830E561)                      |          |
| <b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)      |          |
| <b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A) |          |
| <b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)   |          |
| <b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)            |          |
| <b>STARCH, CORN</b> (UNII: O8232NY3SJ)                     |          |
| <b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)                 |          |
| <b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)                     |          |
| <b>TALC</b> (UNII: 7SEV7J4R1U)                             |          |
| <b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)                 |          |

### Product Characteristics

|                 |                   |                     |          |
|-----------------|-------------------|---------------------|----------|
| <b>Color</b>    | white (Off-White) | <b>Score</b>        | no score |
| <b>Shape</b>    | OVAL              | <b>Size</b>         | 17mm     |
| <b>Flavor</b>   | MINT              | <b>Imprint Code</b> | AAA;1128 |
| <b>Contains</b> |                   |                     |          |

### Packaging

| # | Item Code        | Package Description                                     | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:46122-660-62 | 2 in 1 CARTON   | 11/13/2020           |                    |
| 1 |                  | 12 in 1 BLISTER PACK; Type 0: Not a Combination Product |                      |                    |

### Marketing Information

| Marketing Category  | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC monograph final | part341                                  | 11/13/2020           |                    |

**Labeler** - AMERISOURCE BERGEN (007914906)

