

**POTASSIUM CHLORIDE IN SODIUM CHLORIDE- sodium chloride and potassium chloride injection, solution**  
**ICU Medical Inc.**

-----  
**INTRAVENOUS SOLUTIONS**  
**with POTASSIUM CHLORIDE**

Potassium Chloride in 0.9% Sodium Chloride Injection, USP

**Flexible Plastic Container**

Rx only

**DESCRIPTION**

Intravenous solutions with potassium chloride (I.V. solutions with KCl) are sterile and nonpyrogenic solutions in water for injection. They are for administration by intravenous infusion only.

See Table for summary of content and characteristics of these solutions.

The solutions contain no bacteriostat, antimicrobial agent or added buffer and each is intended only for use as a single-dose injection. When smaller doses are required the unused portion should be discarded.

These solutions are parenteral fluid and/or electrolyte replenishers.

Potassium Chloride, USP is chemically designated KCl, a white granular powder freely soluble in water.

Sodium Chloride, USP is chemically designated NaCl, a white crystalline powder freely soluble in water.

Water for Injection, USP is chemically designated H<sub>2</sub>O.

The flexible plastic container is fabricated from a specially formulated polyvinylchloride. Water can permeate from inside the container into the overwrap but not in amounts sufficient to affect the solution significantly. Solutions inside the plastic container also can leach out certain of its chemical components in very small amounts before the expiration period is attained. However, the safety of the plastic has been confirmed by tests in animals according to USP biological standards for plastic containers.

**CLINICAL PHARMACOLOGY**

When administered intravenously, these solutions provide a source of water and potassium chloride with 0.9% sodium chloride.

Intravenous solutions containing potassium chloride are particularly intended to provide needed potassium cation (K<sup>+</sup>). Potassium is the chief cation of body cells (160 mEq/liter of intracellular water). It is found in low concentration in plasma and extracellular fluids (3.5 to 5.0 mEq/liter in a healthy adult). Potassium plays an important role in electrolyte balance. Normally about 80 to 90% of the potassium intake is excreted in the urine; the remainder in the stools and to a small extent, in the perspiration. The kidney does not conserve potassium well so that during fasting or in patients on a potassium-free diet, potassium loss from the body continues resulting in potassium depletion. A deficiency of either potassium or chloride will lead to a deficit of the other.

Sodium chloride in water dissociates to provide sodium (Na<sup>+</sup>) and chloride (Cl<sup>-</sup>) ions. Sodium (Na<sup>+</sup>) is the principal cation of the extracellular fluid and plays a large part in the therapy of fluid and electrolyte disturbances. Chloride (Cl<sup>-</sup>) has an integral role in buffering action when oxygen and carbon dioxide exchange occurs in the red blood cells. The distribution and excretion of sodium (Na<sup>+</sup>) and chloride (Cl<sup>-</sup>) are largely under the control of the kidney which maintains a balance between intake and output.

Water is an essential constituent of all body tissues and accounts for approximately

70% of total body weight. Average normal adult daily requirement ranges from two to three liters (1.0 to 1.5 liters each for insensible water loss by perspiration and urine production).

Water balance is maintained by various regulatory mechanisms. Water distribution depends primarily on the concentration of electrolytes in the body compartments and sodium ( $\text{Na}^+$ ) plays a major role in maintaining physiologic equilibrium.

## INDICATIONS AND USAGE

These solutions are indicated in patients requiring parenteral administration of potassium chloride and sodium chloride.

## CONTRAINDICATIONS

Solutions containing potassium chloride are contraindicated in diseases where high potassium levels may be encountered.

## WARNINGS

Solutions which contain potassium ions should be used with great care, if at all, in patients with hyperkalemia, severe renal failure and in conditions in which potassium retention is present.

To avoid potassium intoxication, do not infuse these solutions rapidly. In patients with severe renal insufficiency or adrenal insufficiency, administration of potassium chloride may cause potassium intoxication.

Solutions containing sodium ions should be used with great care, if at all, in patients with congestive heart failure, severe renal insufficiency and in clinical states in which there exists edema with sodium retention.

In patients with diminished renal function, administration of solutions containing sodium or potassium ions may result in sodium or potassium retention.

The intravenous administration of these solutions can cause fluid and/or solute overloading resulting in dilution of serum electrolyte concentrations, overhydration, congested states or pulmonary edema.

The risk of dilutional states is inversely proportional to the electrolyte concentration of administered parenteral solutions. The risk of solute overload causing congested states with peripheral and pulmonary edema is directly proportional to the electrolyte concentrations of such solutions.

## PRECAUTIONS

Clinical evaluation and periodic laboratory determinations are necessary to monitor changes in fluid balance, electrolyte concentrations and acid-base balance during prolonged parenteral therapy or whenever the condition of the patient warrants such evaluation.

Caution must be exercised in the administration of parenteral fluids, especially those containing sodium ions, to patients receiving corticosteroids or corticotropin.

**Potassium replacement therapy should be guided primarily by serial electrocardiograms. Plasma potassium levels are not necessarily indicative of tissue potassium levels.**

High plasma concentrations of potassium may cause death through cardiac depression, arrhythmias or arrest.

Potassium-containing solutions should be used with caution in the presence of cardiac disease, particularly in digitalized patients or in the presence of renal disease.

Care should be exercised to insure that the needle (or catheter) is well within the lumen

of the vein and that extravasation does not occur.

Do not administer unless solution is clear and container is undamaged. Discard unused portion.

Pregnancy Category C: Animal reproduction studies have not been conducted with potassium chloride or sodium chloride. It is also not known whether potassium chloride or sodium chloride can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Potassium chloride or sodium chloride should be given to a pregnant woman only if clearly needed.

### **Pediatric Use**

The safety and effectiveness in the pediatric population are based on the similarity of the clinical conditions of the pediatric and adult populations. In neonates or very small infants the volume of fluid may affect fluid and electrolyte balance.

### **ADVERSE REACTIONS**

Reactions which may occur because of the solutions or technique of administration include febrile response, infection at the site of injection, venous thrombosis or phlebitis extending from the site of injection, extravasation and hypervolemia.

If an adverse reaction does occur, discontinue the infusion, evaluate the patient, institute appropriate therapeutic countermeasures and save the remainder of the fluid for examination if deemed necessary.

Nausea, vomiting, abdominal pain and diarrhea have been reported with potassium therapy. The signs and symptoms of potassium intoxication include paresthesias of the extremities, flaccid paralysis, listlessness, mental confusion, weakness and heaviness of the legs, hypotension, cardiac arrhythmias, heart block, electrocardiographic abnormalities such as disappearance of P waves, spreading and slurring of the QRS complex with development of a biphasic curve and cardiac arrest.

Potassium-containing solutions are intrinsically irritating to tissues. Therefore, extreme care should be taken to avoid perivascular infiltration. Local tissue necrosis and subsequent sloughing may result if extravasation occurs. Chemical phlebitis and venospasm have also been reported.

Should perivascular infiltration occur, I.V. administration at that site should be discontinued at once. Local infiltration of the affected area with procaine hydrochloride, 1%, to which hyaluronidase may be added, will often reduce venospasm and dilute the potassium remaining in the tissues locally. Local application of heat may also be helpful.

### **OVERDOSAGE**

In the event of potassium overdosage, discontinue the infusion immediately and institute intensive corrective therapy to reduce serum potassium levels. (See **WARNINGS** and **PRECAUTIONS**.)

### **DOSAGE AND ADMINISTRATION**

These solutions should be administered only by intravenous infusion and as directed by the physician. The dose and rate of injection are dependent upon the age, weight and clinical condition of the patient. If the serum potassium level is greater than 2.5 mEq/liter, potassium should be given at a rate not to exceed 10 mEq/hour in a concentration less than 30 mEq/liter. Somewhat faster rates and greater concentrations (usually up to 40 mEq/liter) of potassium may be indicated in patients with more severe potassium deficiency. The total 24-hour dose should not generally exceed 200 mEq of potassium.

### **Drug Interactions**

Additives may be incompatible. Consult with pharmacist, if available. When introducing additives, use aseptic technique, mix thoroughly and do not store.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit. (See **PRECAUTIONS**.)

## INSTRUCTIONS FOR USE

### To Open

Tear outer wrap at notch and remove solution container. If supplemental medication is desired, follow directions below before preparing for administration. Some opacity of the plastic due to moisture absorption during the sterilization process may be observed. This is normal and does not affect the solution quality or safety. The opacity will diminish gradually.

### To Add Medication

1. Prepare additive port.
2. Using aseptic technique and an additive delivery needle of appropriate length, puncture resealable additive port at target area, inner diaphragm and inject. Withdraw needle after injecting medication.
3. The additive port may be protected by covering with an additive cap.
4. Mix container contents thoroughly.

### Preparation for Administration

#### (Use aseptic technique)

1. Close flow control clamp of administration set.
2. Remove cover from outlet port at bottom of container.
3. Insert piercing pin of administration set into port with a twisting motion until the set is firmly seated. **NOTE:** See full directions on administration set carton.
4. Suspend container from hanger.
5. Squeeze and release drip chamber to establish proper fluid level in chamber.
6. Open flow control clamp and clear air from set. Close clamp.
7. Attach set to venipuncture device. If device is not indwelling, prime and make venipuncture.
8. Regulate rate of administration with flow control clamp.

**WARNING: Do not use flexible container in series connections.**

## HOW SUPPLIED

Intravenous solutions with potassium chloride (I.V. solution with KCl) are supplied in single-dose flexible plastic containers. See Table:

Potassium Chloride in										
0.9% Sodium Chloride										
Inj., USP			COMPOSITION				Approx. Ionic Concentrations			
			(g/L)		Calculated	pH	(mEq/L)			
NDC No.	mEq Potassium	Size (mL)	Sodium Chloride	Potassium Chloride	Osmolarity (mOsmol/L)	(range)	Sodium (Na <sup>+</sup> )	Potassium (K <sup>+</sup> )	Chloride (Cl <sup>-</sup> )	Approximate kcal/L
0409-7115-09	20 mEq	1000	9	1.49	348	4.8 (3.5 to 6.5)	154	20	174	0
0990-7115-09	20 mEq	1000	9	1.49	348	4.8 (3.5 to 6.5)	154	20	174	0
0409-7116-	40 mEq	1000	9	2.98	388	4.8 (3.5 to 6.5)	154	40	194	0

09										
0990-7116-09	40 mEq	1000	9	2.98	388	4.8 (3.5 to 6.5)	154	40	194	0

ICU Medical is transitioning NDC codes from the "0409" to a "0990" labeler code. Both NDC codes are expected to be in the market for a period of time.

Store at 20 to 25°C (68 to 77°F). [See USP Controlled Room Temperature.] Protect from freezing.

Revised: March, 2020

IFU0000168

ICU Medical, Inc., Lake Forest, Illinois, 60045, USA

## **PRINCIPAL DISPLAY PANEL - 20 mEq Bag Label**

20 mEq POTASSIUM

1000 mL

NDC 0990-7115-09

20

mEq

POTASSIUM

CHLORIDE

in 0.9%

Sodium Chloride

Injection, USP

EACH 100 mL CONTAINS POTASSIUM CHLORIDE 149 mg; SODIUM CHLORIDE 900 mg IN WATER FOR INJECTION. ELECTROLYTES PER 1000 mL: POTASSIUM 20 mEq; SODIUM 154 mEq; CHLORIDE; 174 mEq. 348 mOsmol/LITER (CALC.) pH 4.8 (3.5 to 6.5)

ADDITIVES MAY BE INCOMPATIBLE. CONSULT WITH PHARMACIST, IF AVAILABLE. WHEN INTRODUCING ADDITIVES, USE ASEPTIC TECHNIQUE, MIX THOROUGHLY AND DO NOT STORE.

SINGLE-DOSE CONTAINER. FOR I.V. USE. USUAL DOSAGE: SEE INSERT. STERILE, NONPYROGENIC. USE ONLY IF SOLUTION IS CLEAR AND CONTAINER IS UNDAMAGED. MUST NOT BE USED IN SERIES CONNECTIONS.

Rx ONLY

3

v

CONTAINS DEHP

ICU Medical, Inc.,  
Lake Forest, Illinois, 60045, USA

icumedical  
IMP0000049

# 20 mEq POTASSIUM

1000 mL



NDC 0990-7115-09

**20  
mEq**

**POTASSIUM  
CHLORIDE**

in 0.9%

**Sodium Chloride**

**Injection, USP**

EACH 100 mL CONTAINS POTASSIUM CHLORIDE 149 mg; SODIUM CHLORIDE 900 mg IN WATER FOR INJECTION. ELECTROLYTES PER 1000 mL: POTASSIUM 20 mEq; SODIUM 154 mEq; CHLORIDE; 174 mEq.

348 mOsmol/LITER (CALC.) pH 4.8 (3.5 to 6.5)

ADDITIVES MAY BE INCOMPATIBLE.  
CONSULT WITH PHARMACIST, IF AVAILABLE.  
WHEN INTRODUCING ADDITIVES, USE  
ASEPTIC TECHNIQUE,  
MIX THOROUGHLY AND DO NOT STORE.

SINGLE-DOSE CONTAINER. FOR I.V. USE.

USUAL DOSAGE: SEE INSERT.

STERILE, NONPYROGENIC. USE ONLY IF  
SOLUTION IS CLEAR AND CONTAINER IS  
UNDAMAGED. MUST NOT BE USED IN SERIES  
CONNECTIONS.

Rx ONLY



CONTAINS DEHP

ICU Medical, Inc.,  
Lake Forest, Illinois, 60045, USA

**icumedical**  
IMP0000049

## PRINCIPAL DISPLAY PANEL - 40 mEq Bag Label

40 mEq POTASSIUM

1000 mL

NDC 0990-7116-09

40

mEq

POTASSIUM

CHLORIDE

in 0.9%

Sodium Chloride

Injection, USP

EACH 100 mL CONTAINS POTASSIUM CHLORIDE 298 mg; SODIUM CHLORIDE 900 mg IN WATER FOR INJECTION. ELECTROLYTES PER 1000 mL: POTASSIUM 40 mEq; CHLORIDE 194 mEq; SODIUM 154 mEq. 388 mOsmol/LITER (CALC.)

pH 4.8 (3.5 to 6.5)

ADDITIVES MAY BE INCOMPATIBLE. CONSULT WITH PHARMACIST, IF AVAILABLE. WHEN INTRODUCING ADDITIVES, USE ASEPTIC TECHNIQUE, MIX THOROUGHLY AND DO NOT STORE.

SINGLE-DOSE CONTAINER. FOR I.V. USE. USUAL DOSAGE: SEE INSERT. STERILE, NONPYROGENIC. USE ONLY IF SOLUTION IS CLEAR AND CONTAINER IS UNDAMAGED. MUST NOT BE USED IN SERIES CONNECTIONS.

Rx ONLY

3

v

CONTAINS DEHP

IMP0000050

ICU Medical, Inc.,  
Lake Forest, Illinois, 60045, USA

icumedical

# 40 mEq POTASSIUM

1000 mL



NDC 0990-7116-09

— 1

**40  
mEq**

**POTASSIUM  
CHLORIDE**

— 2

in 0.9%

— 3

**Sodium Chloride  
Injection, USP**

— 4

EACH 100 mL CONTAINS POTASSIUM CHLORIDE  
298 mg; SODIUM CHLORIDE 900 mg IN  
WATER FOR INJECTION. ELECTROLYTES PER  
1000 mL: POTASSIUM 40 mEq; CHLORIDE  
194 mEq; SODIUM 154 mEq.

— 5

388 mOsmol/LITER (CALC.) pH 4.8 (3.5 to 6.5)

— 6

ADDITIVES MAY BE INCOMPATIBLE. CONSULT  
WITH PHARMACIST, IF AVAILABLE. WHEN  
INTRODUCING ADDITIVES,  
USE ASEPTIC TECHNIQUE,  
MIX THOROUGHLY AND DO NOT STORE.

— 7

SINGLE-DOSE CONTAINER. FOR I.V. USE.  
USUAL DOSAGE: SEE INSERT.

— 8

STERILE, NONPYROGENIC. USE ONLY IF  
SOLUTION IS CLEAR AND CONTAINER IS  
UNDAMAGED. MUST NOT BE USED IN SERIES  
CONNECTIONS.

— 9

Rx ONLY



— 10

IMPO000050

CONTAINS DEHP

ICU Medical, Inc.,  
Lake Forest, Illinois, 60045, USA



**icu**medical

— 11

— 12

— 13

— 14

— 15

## PRINCIPAL DISPLAY PANEL - 1000 mL Bag Pouch

### TO OPEN TEAR AT NOTCH

DO NOT REMOVE FROM OVERWRAP UNTIL READY FOR USE. AFTER REMOVING  
THE OVERWRAP, CHECK FOR MINUTE LEAKS BY SQUEEZING CONTAINER FIRMLY.  
IF LEAKS ARE FOUND, DISCARD SOLUTION AS STERILITY MAY BE IMPAIRED.  
RECOMMENDED STORAGE: ROOM TEMPERATURE (25°C). AVOID EXCESSIVE  
HEAT. PROTECT FROM FREEZING. SEE INSERT.

98-4321-R14-3/98

2

HDPE



## TO OPEN TEAR AT NOTCH



DO NOT REMOVE FROM OVERWRAP UNTIL READY FOR USE. AFTER REMOVING THE OVERWRAP, CHECK FOR MINUTE LEAKS BY SQUEEZING CONTAINER FIRMLY. IF LEAKS ARE FOUND, DISCARD SOLUTION AS STERILITY MAY BE IMPAIRED. RECOMMENDED STORAGE: ROOM TEMPERATURE (25°C). AVOID EXCESSIVE HEAT. PROTECT FROM FREEZING. SEE INSERT.

98-4321-R14-3/98

### POTASSIUM CHLORIDE IN SODIUM CHLORIDE

sodium chloride and potassium chloride injection, solution

#### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0990-7115
Route of Administration	INTRAVENOUS		

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37)	SODIUM CHLORIDE	9 g in 1000 mL
POTASSIUM CHLORIDE (UNII: 660YQ98I10) (POTASSIUM CATION - UNII:295O53K152)	POTASSIUM CHLORIDE	1.49 g in 1000 mL

#### Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0K00R)	

#### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0990-7115-09	12 in 1 CASE	11/01/2019	
1		1 in 1 POUCH		
1		1000 mL in 1 BAG; Type 0: Not a Combination Product		

#### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA019686	11/01/2019	

### POTASSIUM CHLORIDE IN SODIUM CHLORIDE

sodium chloride and potassium chloride injection, solution

#### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0990-7116
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37)		SODIUM CHLORIDE	9 g in 1000 mL	
POTASSIUM CHLORIDE (UNII: 660YQ98I10) (POTASSIUM CATION - UNII:295O53K152)		POTASSIUM CHLORIDE	2.98 g in 1000 mL	
Inactive Ingredients				
Ingredient Name		Strength		
WATER (UNII: 059QF0KO0R)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0990-7116-09	12 in 1 CASE	12/01/2019	
1		1 in 1 POUCH		
1		1000 mL in 1 BAG; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA		NDA019686	12/01/2019	

**Labeler -** ICU Medical Inc. (118380146)

## Establishment

Name	Address	ID/FEI	Business Operations
Hospira, Inc.		093132819	ANALYSIS(0990-7115, 0990-7116) , MANUFACTURE(0990-7115, 0990-7116) , PACK(0990-7115, 0990-7116) , LABEL(0990-7115, 0990-7116)

Revised: 10/2021

ICU Medical Inc.