

**ASPIRIN REGULAR STRENGTH- aspirin tablet, film coated**  
**FOODHOLD U.S.A., LLC**

-----  
**Care One 44-157**

***Active ingredient (in each tablet)***

Aspirin 325 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever/fever reducer

***Uses***

temporarily relieves:

- minor pain of arthritis
- headache
- muscle pain
- menstrual pain
- toothache
- pain and fever of colds

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- asthma (wheezing)
- hives
- facial swelling
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- take more or for a longer time than directed
- have 3 or more alcoholic drinks every day while using this product

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**

- you are taking a diuretic
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - vomit blood
  - have bloody or black stools
  - feel faint
  - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- redness or swelling is present
- new symptoms occur
- fever gets worse or lasts more than 3 days

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

***Directions***

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

**Other information**

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

**Inactive ingredients**

corn starch, hypromellose, polyethylene glycol, propylene glycol

**Questions or comments?**

**1-800-426-9391**

**Principal Display Panel**

CAREONE®

**NDC 41520-157-87**

*Compare to the  
active ingredient in  
Genuine Bayer® Aspirin<sup>†</sup>*

**Regular Strength  
ASPIRIN**

**325 mg**

Pain Reliever  
Fever Reducer (**NSAID**)

Safe Pain Relief

*Actual  
Size*

**300 TABLETS**

**OUR PHARMACISTS RECOMMEND RX**

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

<sup>†</sup>This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Genuine Bayer® Aspirin.  
50844 REV0122B15717


**DISTRIBUTED BY FOODHOLD U.S.A., LLC  
LANDOVER, MD 20785 • 1-877-846-9949**

**©2022 S&S Brands, LLC • Quality guaranteed or your money back.**

CAREone®  
NDC 41520-157-87  
Compare to the  
active ingredient in  
Genuine Bayer® Aspirin®

Regular Strength  
**ASPIRIN**

325 mg  
Pain Reliever  
Fever Reducer (NSAID)  
Safe Pain Relief

  
Actual  
Size

OUR PHARMACISTS  
RECOMMEND

300 TABLETS


TAMPER EVIDENCE: DO NOT USE IF IMPRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

**Drug Facts**  
**Active ingredient (in each tablet)** **Purpose**  
Aspirin 325 mg (NSAID)\* ..... Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses**  
temporarily relieves:  
■ minor pain of arthritis  
■ headache  
■ muscle pain  
■ menstrual pain  
■ toothache  
■ pain and fever of colds

**Warnings**  
Reye's syndrome: Children and teenagers who have or  
are recovering from chicken pox or flu-like symptoms  
should not use this product. When using this product, if  
changes in behavior with nausea and vomiting occur,  
consult a doctor because these symptoms could be an  
early sign of Reye's syndrome, a rare but serious illness.  
**Allergy alert:** Aspirin may cause a severe allergic reaction,  
which may include:  
■ asthma (wheezing) ■ hives  
■ facial swelling ■ shock  
**Stomach bleeding warning:** This product contains an NSAID,  
which may cause severe stomach bleeding. The chance is higher  
if you ■ are age 60 or older  
■ have had stomach ulcers or bleeding problems  
■ take a blood thinning (anticoagulant) or steroid drug  
■ take other drugs containing prescription or nonprescription  
NSAIDs (aspirin, ibuprofen, naproxen, or others)  
■ take more or for a longer time than directed  
■ have 3 or more alcoholic drinks every day while using this  
product  
**Do not use** ■ if you are allergic to aspirin or any other pain  
reliever/fever reducer  
■ if you have ever had an allergic reaction to this product or any  
of its ingredients

\*This product is not manufactured or distributed by Bayer AG, owner of the  
registered trademark Genuine Bayer® Aspirin.  
50844 REV0122815717  
DISTRIBUTED BY FOODMOLD U.S.A., LLC  
LANDOVER, MD 20785 • 1-877-846-9949  
©2022 SAS Brands, LLC • Quality guaranteed or your money back.

  
3 41520 31361 5

PEEL HERE FOR MORE DRUG FACTS

**STOP PEELING**

**Drug Facts (continued)**  
Ask a doctor before use if ■ you are taking a diuretic  
■ stomach bleeding warning applies to you  
■ you have a history of stomach problems, such as  
heartburn ■ you have asthma  
■ you have high blood pressure, heart disease, liver  
disease, or kidney disease  
**Ask a doctor or pharmacist before use if you are taking**  
a prescription drug for ■ gout ■ diabetes ■ arthritis  
**Stop use and ask a doctor if** ■ an allergic reaction  
occurs. Seek medical help right away.  
■ you experience any of the following signs of stomach  
bleeding: ■ vomit blood  
■ have bloody or black stools ■ feel faint  
■ have stomach pain that does not get better  
■ ringing in the ears or a loss of hearing occurs  
■ pain gets worse or lasts more than 10 days  
■ redness or swelling is present  
■ new symptoms occur  
■ fever gets worse or lasts more than 3 days  
These could be signs of a serious condition.  
**If pregnant or breast-feeding, ask a health professional**  
before use. It is especially important not to use aspirin at  
20 weeks or later in pregnancy unless definitely directed  
to do so by a doctor because it may cause problems in the  
unborn child or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get  
medical help or contact a Poison Control Center right  
away.

**Directions**  
■ do not take more than directed  
■ drink a full glass of water with each dose  
■ adults and children 12 years and over: take 1 or 2  
tablets every 4 hours or 3 tablets every 6 hours not to  
exceed 12 tablets in 24 hours unless directed by a  
doctor  
■ children under 12 years: do not use unless directed by  
a doctor  
**Other information** ■ store at 25°C (77°F);  
excursions permitted between 15°-30°C (59°-86°F)  
■ use by expiration date on package  
**Inactive ingredients** corn starch, hypromellose,  
polyethylene glycol, propylene glycol  
**(Questions or comments?) 1-800-426-9391**

CareOne 44-157

ASPIRIN REGULAR STRENGTH			
aspirin tablet, film coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41520-157
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)		ASPIRIN	325 mg
Inactive Ingredients			
Ingredient Name			Strength
STARCH, CORN (UNII: O8232NY35J)			
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)			
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			

Product Characteristics				
Color		white	Score	no score
Shape		ROUND	Size	10mm
Flavor			Imprint Code	44;157;ASPIRIN
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41520-157-78	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/10/1996	
2	NDC:41520-157-87	300 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/10/1996	
3	NDC:41520-157-90	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/10/1996	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M013	02/10/1996	

**Labeler** - FOODHOLD U.S.A., LLC (809183973)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(41520-157)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(41520-157)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(41520-157)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(41520-157)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(41520-157)