

**UPSET STOMACH RELIEF TO GO- bismuth subsalicylate powder**  
**Breakthrough Products Inc.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Upset Stomach Relief to Go™**

**Drug Facts**

**Active ingredient (per powder)**

Bismuth subsalicylate 525mg

**Purpose**

Upset stomach reliever/antidiarrheal

**Uses**

Relieves:

- traveler's diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
  - heartburn
  - indigestion
  - nausea
  - gas
  - belching
  - fullness

**Warnings**

**Reye's syndrome**

Children and teenagers who have or are recovering from chickenpox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert**

Contains salicylate.

**Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

**Do not use if you have**

- an ulcer
- a bleeding problem
- bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool

**Ask a doctor or pharmacist before use if you are** taking any drug for:

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

**When using this product** a temporary, but harmless, darkening of the stool and/or tongue may occur.

**Stop use and ask a doctor if**

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts for more than 2 days

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions****adults and children 12 years and over:**

- see instructions in red box for opening packet
- place 1 powder on the tongue and swallow with or without water every ½ to 1 hour as needed
- do not exceed 8 doses in 24 hours
- use until diarrhea stops but not more than 2 days
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**children under 12 years: ask a doctor****Other information**

each dose contains salicylate 199mg

**Inactive ingredients**

citric acid, flavor, glucose, silica, sucralose

**Questions?**

1-888-998-7436 (Mon-Fri 9AM-5PM MDT)

<b>Do not use if pouch is torn or open</b>
------------------------------------------------

*distributed by*

**URGENT Rx®**

**Breakthrough Products, Inc.**

Denver, CO 80202

**PRINCIPAL DISPLAY PANEL - 1 Powder Packet**

*New!*

**NO  
WATER  
REQUIRED**  
POUR DIRECTLY IN MOUTH

**URGENT Rx<sup>®</sup>  
FAST  
POWDERS<sup>™</sup>**

▣ SEE OPENING INSTRUCTIONS ON BACK PANEL ▣

**UPSET  
STOMACH**

**URGENT Rx<sup>®</sup>**

**RELIEF  
TO-GO<sup>™</sup>**

BISMUTH SUBSALICYLATE/  
UPSET STOMACH RELIEVER & ANTIDIARRHEAL

**cherry**

distributed by UrgentRx  
1 POWDER PACK

New!

POUR NO WATER

Do not use if pouch is torn or open

Upset Stomach Relief to Go™

distributed by URGENT Rx®

Breakthrough Products, Inc. Denver, CO 80202

Fold top and store in purse or wallet

US Pat. No 8,376,140

CCUS074-A



TO OPEN

- 1) FOLD ON LINE TO EXPOSE NOTCH
- 2) WHILE FOLDED, TEAR AT NOTCH

(or use scissors)

Drug Facts

Active Ingredient (per powder)

Bismuth subsalicylate 525mg ..... Upset stomach reliever/antidiarrheal

Uses Relieves: • traveler's diarrhea • diarrhea • upset stomach due

to overindulgence in food and drink, including: • heartburn

• indigestion • nausea • gas • belching • fullness

Warnings Children and teenagers who have or are

recovering from chickenpox or flu-like symptoms should not use this

product. When using this product, if changes in behavior with nausea

and vomiting occur, consult a doctor because these symptoms could

be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are • allergic

to salicylates (including aspirin) • taking other salicylate products Do

not use if you have • an ulcer • a bleeding problem • bloody or black stool Ask a

doctor before use if you have • fever • mucus in the stool Ask a doctor or

pharmacist before use if you are taking any drug for: • anticoagulation (thinning the

blood) • diabetes • gout • arthritis When using this product a temporary, but

harmless, darkening of the stool and/or tongue may occur. Stop use and ask a doctor

if • symptoms get worse or last more than 2 days • ringing in the ears or loss of hearing

occurs • diarrhea lasts for more than 2 days If pregnant or breast-feeding, ask a

health professional before use. Keep out of reach of children. In case of overdose, get

medical help or contact a Poison Control Center right away. Directions adults and

children 12 years and over: • see instructions in red box for opening packet • place 1

powder on the tongue and swallow with or without water every ½ to 1 hour as needed •

do not exceed 8 doses in 24 hours • use until diarrhea stops but not more than 2 days •

drink plenty of clear fluids to help prevent dehydration caused by diarrhea children

under 12 years: ask a doctor Other information each dose contains salicylate

199mg inactive ingredients citric acid, flavor, glucose, silica, sucralose

Questions? 1-888-998-7436 (Mon-Fri 9AM-5PM MDT)

WATER  
REQUIRED  
DIRECTLY IN MOUTH

URGENT Rx<sup>®</sup>  
**FAST**  
POWDERS™

◀ SEE OPENING INSTRUCTIONS ON BACK PANEL ▶

**UPSET  
STOMACH**



**RELIEF  
TO-GO™**

BISMUTH SUBSALICYLATE/  
UPSET STOMACH RELIEVER & ANTIDIARRHEAL

cherry

distributed by UrgentRx

1 POWDER PACK

**UPSET STOMACH RELIEF TO GO**

bismuth subsalicylate powder

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG LABEL	<b>Item Code (Source)</b>	NDC:51596-004
<b>Route of Administration</b>	ORAL	<b>DEA Schedule</b>	

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
Bismuth subsalicylate (SALICYLIC ACID and BISMUTH CATION)	Bismuth subsalicylate	525 mg

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
anhydrous citric acid	
dextrose	
silicon dioxide	
sucralose	

**Product Characteristics**

<b>Color</b>	WHITE	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CHERRY	<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51596-004-05	5 in 1 BOX		
1	NDC:51596-004-01	1 in 1 POUCH; Combination Product Type = C112160		
2	NDC:51596-004-12	12 in 1 BOX		
2	NDC:51596-004-01	1 in 1 POUCH; Combination Product Type = C112160		
3	NDC:51596-004-24	24 in 1 BOX		
3	NDC:51596-004-01	1 in 1 POUCH; Combination Product Type = C112160		
4	NDC:51596-004-10	10 in 1 BOX		
4		1 in 1 PACKET; Combination Product Type = C112160		

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC MONOGRAPH FINAL	part335	09/01/2011	

**Labeler** - Breakthrough Products Inc. (962008251)