

# VANCOMYCIN HYDROCHLORIDE- vancomycin hydrochloride capsule

## PAI Holdings, LLC

### HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use VANCOMYCIN HYDROCHLORIDE CAPSULES safely and effectively. See full prescribing information for VANCOMYCIN HYDROCHLORIDE CAPSULES. VANCOMYCIN HYDROCHLORIDE capsules, for oral use Initial U.S. Approval: 1986 To reduce the development of drug-resistant bacteria and maintain the effectiveness of vancomycin hydrochloride capsules and other antibacterial drugs, vancomycin hydrochloride capsules should be used only to treat infections that are proven or strongly suspected to be caused by susceptible bacteria.

### INDICATIONS AND USAGE

Treatment of: ( 1 )

- *C. difficile* -associated diarrhea
- Enterocolitis caused by *Staphylococcus aureus* (including methicillin- resistant strains)

Important Limitations: ( 1 ) ( 5.1 )

- Orally administered vancomycin hydrochloride capsules is not effective for other types of infections.

### DOSAGE AND ADMINISTRATION

- *C. difficile*- associated diarrhea:
  - Adult Patients ( $\geq 18$  years of age): 125 mg orally 4 times daily for 10 days. ( 2.1 )
  - Pediatric Patients (<18 years of age): 40 mg/kg in 3 or 4 divided doses for 7 to 10 days. The total daily dosage should not exceed 2 g. ( 2.2 )
- Staphylococcal enterocolitis:
  - Adult Patients ( $\geq 18$  years of age): 500 mg to 2 g orally in 3 or 4 divided doses for 7 to 10 days. ( 2.1 )
  - Pediatric Patients (<18 years of age): 40 mg/kg in 3 or 4 divided doses for 7 to 10 days. The total daily dosage should not exceed 2 g. ( 2.2 )

### DOSAGE FORMS AND STRENGTHS

- 125 mg capsules ( 3 )
- 250 mg capsules ( 3 )

### CONTRAINDICATIONS

Hypersensitivity to vancomycin ( 4 ) (4)

### WARNINGS AND PRECAUTIONS

- **Vancomycin hydrochloride capsules must be given orally for treatment of staphylococcal enterocolitis and *C. difficile* -associated diarrhea. Orally administered vancomycin hydrochloride capsules are not effective for other types of infections.** ( 5.1 )
- Clinically significant serum concentrations have been reported in some patients who have taken multiple oral doses of vancomycin hydrochloride capsules for active *C. difficile* -associated diarrhea. Monitoring of serum concentrations may be appropriate in some instances. ( 5.2 )
- Nephrotoxicity has occurred following oral vancomycin hydrochloride capsules therapy and can occur either during or after completion of therapy. The risk is increased in geriatric patients ( 5.3 ) Monitor renal function.
- Ototoxicity has occurred in patients receiving vancomycin hydrochloride capsules. ( 5.4 ) Assessment of auditory function may be appropriate in some instances.
- Prescribing vancomycin hydrochloride capsules in the absence of a proven or strongly suspected bacterial infection is unlikely to provide benefit to the patient and increases the risk of the development of drug resistant bacteria. ( 5.6 )
- The 125 mg strength capsules contains FD&C Yellow No. 5 (tartrazine) which may cause allergic-type reactions (including bronchial asthma) in certain susceptible persons. ( 5.7 )

### ADVERSE REACTIONS

The most common adverse reactions ( $\geq 10\%$ ) were nausea (17%), abdominal pain (15 %), and hypokalemia (13%). ( 6.1 )

To report SUSPECTED ADVERSE REACTIONS, contact Pharmaceutical Associates, Inc. at 1-800-845-8210 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch) .

### DRUG INTERACTIONS

No drug interaction studies have been conducted. ( 7 ) (7)

### USE IN SPECIFIC POPULATIONS

- **Pediatrics:** Safety and effectiveness in patients <18 years of age have not been established. ( 8.4 )
- **Geriatrics:** In patients >65 years of age, including those with normal renal function prior to treatment, renal function should be monitored during and following treatment with vancomycin hydrochloride capsules to detect potential vancomycin induced nephrotoxicity. ( 5.3 ) ( 6.1 ) ( 8.5 ) ( 14.1 )

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 5/2019

## **2 DOSAGE AND ADMINISTRATION**

2.1 Adults

2.2 Pediatric Patients

## **3 DOSAGE FORMS AND STRENGTHS**

## **4 CONTRAINDICATIONS**

## **5 WARNINGS AND PRECAUTIONS**

5.1 Oral Use Only

5.2 Potential for Systemic Absorption

5.3 Nephrotoxicity

5.4 Ototoxicity

5.5 Superinfection

5.6 Development of Drug-Resistant Bacteria

5.7 Risk of Allergic Reactions due to Tartrazine

## **6 ADVERSE REACTIONS**

6.1 Clinical Trial Experience

6.2 Postmarketing Experience

## **7 DRUG INTERACTIONS**

## **8 USE IN SPECIFIC POPULATIONS**

8.1 Pregnancy

8.3 Nursing Mothers

8.4 Pediatric Use

8.5 Geriatric Use

## **10 OVERDOSAGE**

## **11 DESCRIPTION**

## **12 CLINICAL PHARMACOLOGY**

12.1 Mechanism of Action

12.3 Pharmacokinetics

12.4 Microbiology

## **13 NONCLINICAL TOXICOLOGY**

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

## **14 CLINICAL STUDIES**

14.1 Diarrhea Associated With *Clostridium Difficile*

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

## **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

Vancomycin Hydrochloride Capsules are indicated for the treatment of *C. difficile* - associated diarrhea. Vancomycin Hydrochloride Capsules is also used for the treatment of enterocolitis caused by *Staphylococcus aureus* (including methicillin-resistant strains). Parenteral administration of vancomycin is not effective for the above infections; therefore, Vancomycin Hydrochloride Capsules must be given orally for these infections.

**Orally administered Vancomycin Hydrochloride Capsules is not effective for other types of infections.**

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Vancomycin Hydrochloride Capsules and other antibacterial drugs, Vancomycin Hydrochloride Capsules should be used only to treat infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

### **2 DOSAGE AND ADMINISTRATION**

#### **2.1 Adults**

Vancomycin Hydrochloride Capsules are used in treating *C. difficile* -associated diarrhea

and staphylococcal enterocolitis.

- *C. difficile*- associated diarrhea: The recommended dose is 125 mg administered orally 4 times daily for 10 days.
- Staphylococcal enterocolitis: Total daily dosage is 500 mg to 2 g administered orally in 3 or 4 divided doses for 7 to 10 days.

## 2.2 Pediatric Patients

The usual daily dosage is 40 mg/kg in 3 or 4 divided doses for 7 to 10 days. The total daily dosage should not exceed 2 g.

## 3 DOSAGE FORMS AND STRENGTHS

Vancomycin Hydrochloride Capsules USP 125 mg\* CAPSULES have a blue cap and yellow body imprinted with "OP64" on both cap and body in white ink.

Vancomycin Hydrochloride Capsules USP 250 mg\* CAPSULES have a blue cap and dark-yellow body imprinted with "OP65" on both cap and body in white ink.

\*Equivalent to vancomycin.

## 4 CONTRAINDICATIONS

Vancomycin Hydrochloride Capsules are contraindicated in patients with known hypersensitivity to vancomycin.

## 5 WARNINGS AND PRECAUTIONS

### 5.1 Oral Use Only

**This preparation for the treatment of colitis is for oral use only and is not systemically absorbed. Vancomycin Hydrochloride Capsules must be given orally for treatment of staphylococcal enterocolitis and Clostridium difficile-associated diarrhea. Orally administered Vancomycin Hydrochloride Capsules are not effective for other types of infections.**

**Parenteral administration of vancomycin is not effective for treatment of staphylococcal enterocolitis and C. difficile-associated diarrhea. If parenteral vancomycin therapy is desired, use an intravenous preparation of vancomycin and consult the package insert accompanying that preparation .**

### 5.2 Potential for Systemic Absorption

Clinically significant serum concentrations have been reported in some patients who have taken multiple oral doses of Vancomycin Hydrochloride Capsules for active *C. difficile*-associated diarrhea. Some patients with inflammatory disorders of the intestinal mucosa also may have significant systemic absorption of vancomycin. These patients may be at risk for the development of adverse reactions associated with higher doses of Vancomycin Hydrochloride Capsules; therefore, monitoring of serum concentrations of vancomycin may be appropriate in some instances, e.g., in patients with renal insufficiency and/or colitis or in those receiving concomitant therapy with an aminoglycoside antibiotic.

### 5.3 Nephrotoxicity

Nephrotoxicity (e.g., reports of renal failure, renal impairment, blood creatinine increased) has occurred following oral Vancomycin Hydrochloride Capsules therapy in randomized controlled clinical studies, and can occur either during or after completion of therapy.

The risk of nephrotoxicity is increased in patients >65 years of age [see ADVERSE REACTIONS, Clinical Trial Experience ( 6.1 ) and USE IN SPECIFIC POPULATIONS, Geriatric Use ( 8.5 )] .

In patients >65 years of age, including those with normal renal function prior to treatment, renal function should be monitored during and following treatment with Vancomycin Hydrochloride Capsules to detect potential vancomycin induced nephrotoxicity.

## 5.4 Ototoxicity

Ototoxicity has occurred in patients receiving vancomycin. It may be transient or permanent. It has been reported mostly in patients who have been given excessive intravenous doses, who have an underlying hearing loss, or who are receiving concomitant therapy with another ototoxic agent, such as an aminoglycoside. Serial tests of auditory function may be helpful in order to minimize the risk of ototoxicity [see *ADVERSE REACTIONS, Postmarketing Experience* ( 6.2 )].

## 5.5 Superinfection

Use of Vancomycin Hydrochloride Capsules may result in the overgrowth of nonsusceptible bacteria. If superinfection occurs during therapy, appropriate measures should be taken.

## 5.6 Development of Drug-Resistant Bacteria

Prescribing Vancomycin Hydrochloride Capsules in the absence of a proven or strongly suspected bacterial infection is unlikely to provide benefit to the patient and increases the risk of the development of drug resistant bacteria.

## 5.7 Risk of Allergic Reactions due to Tartrazine

The 125 mg strength capsules contains FD&C Yellow No. 5 (tartrazine) which may cause allergic-type reactions (including bronchial asthma) in certain susceptible persons. Although the overall incidence of FD&C Yellow No. 5 (tartrazine) sensitivity in the general population is low, it is frequently seen in patients who also have aspirin hypersensitivity.

## 6 ADVERSE REACTIONS

### 6.1 Clinical Trial Experience

Because clinical studies are conducted under widely varying conditions, adverse reaction rates observed in the clinical studies of a drug cannot be directly compared to rates in the clinical studies of another drug and may not reflect the rates observed in practice.

The data described below reflect exposure to vancomycin hydrochloride in 260 adult subjects in two Phase 3 clinical trials for the treatment of diarrhea associated with *C. difficile*. In both trials, subjects received vancomycin hydrochloride capsules 125 mg orally four times daily. The mean duration of treatment was 9.4 days. The median age of patients was 67, ranging between 19 and 96 years of age. Patients were predominantly Caucasian (93%) and 52% were male.

Adverse reactions occurring in  $\geq 5\%$  of vancomycin hydrochloride-treated subjects are shown in Table 1. The most common adverse reactions associated with vancomycin hydrochloride ( $\geq 10\%$ ) were nausea, abdominal pain, and hypokalemia.

**Table 1: Common ( $\geq 5\%$ ) Adverse Reactions for Vancomycin Hydrochloride Reported in Clinical Trials for Treatment of Diarrhea Associated with *C. difficile***

System/Organ Class	Adverse Reaction	VancomycinHydrochloride% (N=260)
<b>Gastrointestinal disorders</b>	Nausea	17
	Abdominal pain	15
	Vomiting	9
	Diarrhea	9
	Flatulence	8
<b>General disorders and administration site conditions</b>	Pyrexia	9
	Edema peripheral	6
	Fatigue	5
<b>Infections and infestations</b>	Urinary tract infection	8
<b>Metabolism and nutrition disorders</b>	Hypokalemia	13
<b>Musculoskeletal and connective tissue disorders</b>	Back pain	6
<b>Nervous system disorders</b>	Headache	7

Nephrotoxicity (e.g., reports of renal failure, renal impairment, blood creatinine increased) occurred in 5% of subjects treated with vancomycin hydrochloride.

Nephrotoxicity following vancomycin hydrochloride typically first occurred within one week after completion of treatment (median day of onset was Day 16). Nephrotoxicity following vancomycin hydrochloride occurred in 6% of subjects >65 years of age and 3% of subjects ≤65 years of age [see **WARNINGS AND PRECAUTIONS, Nephrotoxicity ( 5.3 )**].

The incidences of hypokalemia, urinary tract infection, peripheral edema, insomnia, constipation, anemia, depression, vomiting, and hypotension were higher among subjects >65 years of age than in subjects ≤65 years of age [see **USE IN SPECIFIC POPULATIONS, Geriatric Use ( 8.5 )**].

Discontinuation of study drug due to adverse events occurred in 7% of subjects treated with vancomycin hydrochloride. The most common adverse events leading to discontinuation of vancomycin hydrochloride were *C. difficile* colitis (<1%), nausea (<1%), and vomiting (<1%).

## 6.2 Postmarketing Experience

The following adverse reactions have been identified during post-approval use of vancomycin hydrochloride. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

**Ototoxicity:** Cases of hearing loss associated with intravenously administered vancomycin have been reported. Most of these patients had kidney dysfunction or a preexisting hearing loss or were receiving concomitant treatment with an ototoxic drug [see **WARNINGS AND PRECAUTIONS, Ototoxicity ( 5.4 )**]. Vertigo, dizziness, and tinnitus have been reported.

**Hematopoietic:** Reversible neutropenia, usually starting 1 week or more after onset of intravenous therapy with vancomycin or after a total dose of more than 25 g, has been reported for several dozen patients. Neutropenia appears to be promptly reversible when vancomycin is discontinued. Thrombocytopenia has been reported.

**Miscellaneous:** Patients have been reported to have had anaphylaxis, drug fever, chills, nausea, eosinophilia, rashes (including exfoliative dermatitis), Stevens-Johnson syndrome, toxic epidermal necrolysis, and rare cases of vasculitis in association with the administration of vancomycin.

A condition has been reported that is similar to the IV-induced syndrome with symptoms consistent with anaphylactoid reactions, including hypotension, wheezing, dyspnea, urticaria, pruritus, flushing of the upper body ("Red Man Syndrome"), pain and muscle spasm of the chest and back. These reactions usually resolve within 20 minutes but may persist for several hours.

## 7 DRUG INTERACTIONS

No drug interaction studies have been conducted.

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

Pregnancy Category B – The highest doses of vancomycin tested were not teratogenic in rats given up to 200 mg/kg/day IV (1180 mg/m<sup>2</sup> or 1 times the recommended maximum human dose based on body surface area) or in rabbits given up to 120 mg/kg/day IV (1320 mg/m<sup>2</sup> or 1.1 times the recommended maximum human dose based body surface area). No effects on fetal weight or development were seen in rats at the highest dose tested or in rabbits given 80 mg/kg/day (880 mg/m<sup>2</sup> or 0.74 times the recommended maximum human dose based on body surface area).

In a controlled clinical study, the potential ototoxic and nephrotoxic effects of vancomycin on infants were evaluated when the drug was administered intravenously to pregnant women for serious staphylococcal infections complicating intravenous drug abuse. Vancomycin was found in cord blood. No sensorineural hearing loss or nephrotoxicity attributable to vancomycin was noted. One infant whose mother received vancomycin in the third trimester experienced conductive hearing loss that was not attributed to the administration of vancomycin. Because the number of subjects treated in this study was limited and vancomycin was administered only in the second and third

trimesters, it is not known whether vancomycin causes fetal harm. Because animal reproduction studies are not always predictive of human response, vancomycin hydrochloride capsules should be given to a pregnant woman only if clearly needed.

### 8.3 Nursing Mothers

Vancomycin is excreted in human milk based on information obtained with the intravenous administration of vancomycin. However, systemic absorption of vancomycin is very low following oral administration of vancomycin hydrochloride capsules [see *CLINICAL PHARMACOLOGY, Pharmacokinetics ( 12.3 )*]. It is not known whether vancomycin is excreted in human milk, as no studies of vancomycin concentration in human milk after oral administration have been done. Caution should be exercised when vancomycin hydrochloride capsules is administered to a nursing woman. Because of the potential for adverse events, a decision should be made whether to discontinue nursing or discontinue the drug, taking into account the importance of the drug to the mother.

### 8.4 Pediatric Use

Safety and effectiveness in pediatric patients have not been established.

### 8.5 Geriatric Use

In clinical trials, 54% of vancomycin hydrochloride-treated subjects were >65 years of age. Of these, 40% were between the ages of >65 and 75, and 60% were >75 years of age.

Clinical studies with vancomycin hydrochloride in diarrhea associated with *Clostridium difficile* have demonstrated that geriatric subjects are at increased risk of developing nephrotoxicity following treatment with oral vancomycin hydrochloride, which may occur during or after completion of therapy. In patients >65 years of age, including those with normal renal function prior to treatment, renal function should be monitored during and following treatment with vancomycin hydrochloride to detect potential vancomycin induced nephrotoxicity [see *WARNINGS AND PRECAUTIONS, Nephrotoxicity ( 5.3 )*; *ADVERSE REACTIONS, Clinical Trial Experience ( 6.1 )* and *CLINICAL STUDIES, Diarrhea Associated with Clostridium difficile ( 14.1 )*].

Patients >65 years of age may take longer to respond to therapy compared to patients ≤65 years of age [see *CLINICAL STUDIES, Diarrhea Associated with Clostridium difficile ( 14.1 )*]. Clinicians should be aware of the importance of appropriate duration of vancomycin hydrochloride treatment in patients >65 years of age and not discontinue or switch to alternative treatment prematurely.

## 10 OVERDOSAGE

Supportive care is advised, with maintenance of glomerular filtration. Vancomycin is poorly removed by dialysis. Hemofiltration and hemoperfusion with polysulfone resin have been reported to result in increased vancomycin clearance.

To obtain up-to-date information about the treatment of overdose, a good resource is your certified Regional Poison Control Center. Telephone numbers of certified poison control centers are listed in the *Physicians' Desk Reference (PDR)*. In managing overdosage, consider the possibility of multiple drug overdoses, interaction among drugs, and unusual drug kinetics.

## 11 DESCRIPTION

Vancomycin Hydrochloride Capsules, USP for oral administration contain chromatographically purified vancomycin hydrochloride, a tricyclic glycopeptide antibiotic derived from *Amycolatopsis orientalis* (formerly *Nocardia orientalis*), which has the chemical formula  $C_{66}H_{75}Cl_2N_9O_{24} \cdot HCl$ . The molecular weight of vancomycin hydrochloride is 1485.73; 500 mg of the base is equivalent to 0.34 mmol.

The 125 mg capsules contain vancomycin hydrochloride equivalent to 125 mg (0.08 mmol) vancomycin. These capsules also contain polyethylene glycol 6000, FD&C Yellow No. 5, titanium dioxide, FD&C Blue No. 1, sodium lauryl sulfate and gelatin.

The 250 mg capsules contain vancomycin hydrochloride equivalent to 250 mg (0.17 mmol) vancomycin. These capsules also contain polyethylene glycol 6000, yellow iron



## Gram-positive bacteria

*Staphylococcus aureus* (including methicillin-resistant isolates ) associated with enterocolitis

## Anaerobic Gram-positive bacteria

*Clostridium difficile* isolates associated with *C. difficile* associated diarrhea.

## 13 NONCLINICAL TOXICOLOGY

### 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

No long-term carcinogenesis studies in animals have been conducted.

At concentrations up to 1000 µg/mL, vancomycin had no mutagenic effect *in vitro* in the mouse lymphoma forward mutation assay or the primary rat hepatocyte unscheduled DNA synthesis assay. The concentrations tested *in vitro* were above the peak plasma vancomycin concentrations of 20 to 40 µg/mL usually achieved in humans after slow infusion of the maximum recommended dose of 1 g. Vancomycin had no mutagenic effect *in vivo* in the Chinese hamster sister chromatid exchange assay (400 mg/kg IP) or the mouse micronucleus assay (800 mg/kg IP).

No definitive fertility studies have been conducted.

## 14 CLINICAL STUDIES

### 14.1 Diarrhea Associated With *Clostridium Difficile*

In two trials, Vancomycin Hydrochloride Capsules 125 mg orally four times daily for 10 days was evaluated in 266 adult subjects with *C. difficile* -associated diarrhea (CDAD). Enrolled subjects were 18 years of age or older and received no more than 48 hours of treatment with oral vancomycin hydrochloride or oral/intravenous metronidazole in the 5 days preceding enrollment. CDAD was defined as  $\geq 3$  loose or watery bowel movements within the 24 hours preceding enrollment, and the presence of either *C. difficile* toxin A or B, or pseudomembranes on endoscopy within the 72 hours preceding enrollment. Subjects with fulminant *C. difficile* disease, sepsis with hypotension, ileus, peritoneal signs or severe hepatic disease were excluded.

Efficacy analyses were performed on the Full Analysis Set (FAS), which included randomized subjects who received at least one dose of vancomycin hydrochloride and had any post-dosing investigator evaluation data (N=259; 134 in Trial 1 and 125 in Trial 2).

The demographic profile and baseline CDAD characteristics of enrolled subjects were similar in the two trials. Vancomycin Hydrochloride -treated subjects had a median age of 67 years, were mainly white (93%), and male (52%). CDAD was classified as severe (defined as 10 or more unformed bowel movements per day or  $WBC \geq 15000/mm^3$ ) in 25% of subjects, and 47% were previously treated for CDAD.

Efficacy was assessed by using clinical success, defined as diarrhea resolution and the absence of severe abdominal discomfort due to CDAD, on Day 10. An additional efficacy endpoint was the time to resolution of diarrhea, defined as the beginning of diarrhea resolution that was sustained through the end of the prescribed active treatment period.

The results for clinical success for vancomycin hydrochloride -treated subjects in both trials are shown in Table 2.

**Table 2: Clinical Success Rates (Full Analysis Set)**

	<b>Clinical Success Rate</b>	<b>95% Confidence Interval</b>
	<b>Vancomycin Hydrochloride% (N)</b>	
<b>Trial 1</b>	81.3 (134)	(74.4, 88.3)
<b>Trial 2</b>	80.8 (125)	(73.5, 88.1)

The median time to resolution of diarrhea was 5 days and 4 days in Trial 1 and Trial 2, respectively. For subjects older than 65 years of age, the median time to resolution was 6 days and 4 days in Trial 1 and Trial 2, respectively. In subjects with diarrhea resolution at end-of-treatment with vancomycin hydrochloride capsules, recurrence of CDAD

during the following four weeks occurred in 25 of 107 (23%) and 18 of 102 (18%) in Trial 1 and Trial 2, respectively.

Restriction Endonuclease Analysis (REA) was used to identify *C. difficile* baseline isolates in the BI group. In Trial 1, the Vancomycin Hydrochloride-treated subjects were classified at baseline as follows 31 (23%) with BI strain, 69 (52%) with non-BI strain, and 34 (25%) with unknown strain. Clinical success rates were 87% for BI strain, 81% for non-BI strain, and 76% for unknown strain. In subjects with diarrhea resolution at end-of treatment with vancomycin hydrochloride, recurrence of CDAD during the following four weeks occurred in 7 of 26 subjects with BI strain, 12 of 56 subjects with non-BI strain, and 6 of 25 subjects with unknown strain.

## 16 HOW SUPPLIED/STORAGE AND HANDLING

Vancomycin Hydrochloride Capsules, USP are available in:

The 125 mg\* CAPSULES have a blue cap and yellow body imprinted with "OP64" on both cap and body in white ink.

The 250 mg\* CAPSULES have a blue cap and dark-yellow body imprinted with "OP65" on both cap and body in white ink.

Strength	Pack	NDC Number
Vancomycin 125 mg	A carton contains 2 blister packs. Each blister pack contains 10 capsules for a total of 20 capsules per carton.	NDC No. 0121-0867-20
	Each bottle contains 50 capsules.	NDC No. 0121-0867-50
Vancomycin 250 mg	A carton contains 2 blister packs. Each blister pack contains 10 capsules for a total of 20 capsules per carton.	NDC No. 0121-0890-20
	Each bottle contains 50 capsules.	NDC No. 0121-0890-50

Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F).  
[See USP Controlled Room Temperature]

\*Equivalent to vancomycin.

## 17 PATIENT COUNSELING INFORMATION

Patients should be counseled that antibacterial drugs including Vancomycin Hydrochloride should only be used to treat bacterial infections. They do not treat viral infections (e.g., the common cold). When Vancomycin Hydrochloride is prescribed to treat a bacterial infection, patients should be told that although it is common to feel better early in the course of therapy, the medication should be taken exactly as directed. Skipping doses or not completing the full course of therapy may (1) decrease the effectiveness of the immediate treatment and (2) increase the likelihood that bacteria will develop resistance and will not be treatable by Vancomycin Hydrochloride or other antibacterial drugs in the future.

Manufactured by:

Orient Pharma Co., Ltd.

8 Kehu 1st Road, Huwei Chen, Yunlin, Taiwan 63247

Product of Taiwan Distributed by:

Pharmaceutical Associates, Inc.

1700 Perimeter Road,

Greenville, SC 29605

Rx only

To request additional information or if you have questions concerning vancomycin hydrochloride capsules please phone Pharmaceutical Associates, Inc. at 1-800-845-

**PACKAGE/LABEL DISPLAY PANEL**

**Vancomycin Hydrochloride Capsules USP, Equiv. to 125 mg vancomycin  
NDC 0121-0867-50, Rx Only 50 Capsules**

<p><b>Each Capsule Contains:</b> Vancomycin hydrochloride USP equivalent to 125 mg vancomycin</p> <p><b>Usual Total Daily Adult Dose:</b> 500 mg to 2 g administered in divided doses every 6 to 8 hours. Not for treatment of systemic infections.</p> <p>See package insert for other dosage information.</p> <p><b>Storage:</b> Store at Controlled Room Temperature 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F). [See USP Controlled Room Temperature]</p>	<p>NDC 0121-0867-50</p> <p><b>Vancomycin Hydrochloride Capsules, USP</b></p> <p><b>125 mg*</b></p> <p>* Equivalent to 125 mg vancomycin</p> <p><b>Rx only</b> 50 Capsules</p> 	<p><b>KEEP THIS AND ALL MEDICATION OUT OF THE REACH OF CHILDREN.</b></p> <p>Contains color additives including FD&amp;C Yellow No. 5 (tartrazine)</p> <p><b>Distributed by:</b> Pharmaceutical Associates, Inc. 1700 Perimeter Road Greenville, SC 29605</p> <p><b>Manufactured by:</b> Orient Pharma Co., Ltd. 8 Keahu 1st Road Huwei Chen, Yunlin 63247, Taiwan</p> <p>4000569 Rev.002</p>	
---	---	--	---

**Vancomycin Hydrochloride Capsules USP, Equiv. to 250 mg vancomycin  
NDC 0121-0890-50, Rx Only 50 Capsules**

<p><b>Each Capsule Contains:</b> Vancomycin hydrochloride USP equivalent to 250 mg vancomycin</p> <p><b>Usual Total Daily Adult Dose:</b> 500 mg to 2 g administered in divided doses every 6 to 8 hours. Not for treatment of systemic infections.</p> <p>See package insert for other dosage information.</p> <p><b>Storage:</b> Store at Controlled Room Temperature 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F). [See USP Controlled Room Temperature]</p> <p><b>KEEP THIS AND ALL MEDICATION OUT OF THE REACH OF CHILDREN.</b></p>	<p>NDC 0121-0890-50</p> <p><b>Vancomycin Hydrochloride Capsules, USP</b></p> <p><b>250 mg*</b></p> <p>* Equivalent to 250 mg vancomycin</p> <p><b>Rx only</b> 50 Capsules</p> 	<p style="text-align: right;">4000570 Rev.002</p> <p><b>Manufactured by:</b> Orient Pharma Co., Ltd. 8 Keahu 1st Road Huwei Chen, Yunlin 63247, Taiwan</p> <p><b>Distributed by:</b> Pharmaceutical Associates, Inc. 1700 Perimeter Road Greenville, SC 29605</p>	
--	---	---	---

<b>VANCOMYCIN HYDROCHLORIDE</b>			
vancomycin hydrochloride capsule			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0121-0867
<b>Route of Administration</b>	ORAL		
<b>Active Ingredient/Active Moiety</b>			
	<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
	VANCOMYCIN HYDROCHLORIDE (UNII: 71W0621TJD) (VANCOMYCIN - UNII: 6Q205EH1VU)	VANCOMYCIN	125 mg

Inactive Ingredients	
Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C YELLOW NO. 5 (UNII: I753WB2F1M)	
GELATIN (UNII: 2G86QN327L)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
POLYETHYLENE GLYCOL 6000 (UNII: 30IQX730WE)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SHELLAC (UNII: 46N107B71O)	
GLYCERYL MONOOLEATE (UNII: C4YAD5F5G6)	

  

Product Characteristics			
Color	blue (Blue Cap) , yellow (Yellow Body)	Score	no score
Shape	CAPSULE (Capsule)	Size	18mm
Flavor		Imprint Code	OP;64
Contains			

  

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0121-0867-50	50 in 1 BOTTLE; Type 0: Not a Combination Product	03/10/2020	

  

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA210729	07/25/2019	

VANCOMYCIN HYDROCHLORIDE			
vancomycin hydrochloride capsule			

  

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0121-0890
Route of Administration	ORAL		

  

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
VANCOMYCIN HYDROCHLORIDE (UNII: 71W0621TJD) (VANCOMYCIN - UNII:6Q205EH1VU)	VANCOMYCIN	250 mg

  

Inactive Ingredients	
Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
POLYETHYLENE GLYCOL 6000 (UNII: 30IQX730WE)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SHELLAC (UNII: 46N107B71O)	
GLYCERYL MONOOLEATE (UNII: C4YAD5F5G6)	

  

Product Characteristics			
Color	blue (Blue Cap) , yellow (Dark-Yellow Body)	Score	no score

<b>Shape</b>	CAPSULE (Capsule)	<b>Size</b>	22mm
<b>Flavor</b>		<b>Imprint Code</b>	OP;65
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0121-0890-50	50 in 1 BOTTLE; Type 0: Not a Combination Product	03/10/2020	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA210729	07/25/2019	

**Labeler - PAI Holdings, LLC (044940096)**

Revised: 1/2023

PAI Holdings, LLC