VITAFOL-OB PLUS DHA PRENATAL SUPPLEMENT PLUS DHA- vitamin a, ascorbic acid, vitamin d, .alpha.-tocopherol, thiamine mononitrate, riboflavin, niacin, pyridoxine hydrochloride, folic acid, cyanocobalamin, calcium, iron, magnesium, zinc, copper, and doconexent Exeltis USA, Inc

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Vitafol [®]-OB+DHA

0642-0076-30

Prenatal Supplement with DHA

Rx

COMPOSITION

Each Vitafol [®]-OB caplet contains:

Vitamin A (as beta carotene)	810 mcg RAE
Vitamin C (as ascorbic acid)	70 mg
Vitamin D (as cholecalciferol)	10 mcg
Vitamin E (as dl-alpha tocopheryl acetate)	13.5 mg
Thiamin (Vitamin B1 as thiamine mononitrate)	1.6 mg
Riboflavin (Vitamin B2)	1.8 mg
Niacin (as niacinamide)	18 mg NE
Vitamin B6 (as pyridoxine hydrochloride)	2.5 mg
Folate (as folic acid)	1700 mcg
	DFE
Vitamin B12 (as cyanocobalamin)	12 mcg
Calcium (as calcium carbonate)	100 mg
lron (as ferrous fumarate)	65 mg
Magnesium (as magnesium oxide)	25 mg
Zinc (as zinc oxide)	25 mg
Copper (as copper oxide)	2 mg
Each DHA softgel capsule contains:	
Docosahexaenoic acid (DHA) (from natural algal oil)	250 mg

Other ingredients: gelatin (bovine), gylcerin, microcrystalline cellulose, croscarmellose sodium, silicon dioxide, maltodextrin, copovidone, stearic acid, water, hydroxypropyl methylcellulose, dicalcium phosphate, acacia gum, titanium dioxide, polydextrose, starch, magnesium stearate, triacetin, modified food starch, mannitol, vitamin E alcohol,

polyethylene glycol, talc, FD&C Blue #1, FD&C Blue #2. Contains: Soy.

INDICATIONS AND USAGE

Vitafol [®]-OB+DHA is indicated to provide vitamin, mineral and omega-3 fatty acid supplementation prior to conception, throughout pregnancy, and during the postnatal period for the lactating and non-lactating mother.*

Vitafol [®]-OB+DHA does not contain fish, fish oils, fish proteins or fish byproducts.

CONTRAINDICATIONS

Vitafol [®]-OB+DHA is contraindicated in patients with hypersensitivity to any of its components or color additives.

Folic acid is contraindicated in patients with untreated and uncomplicated pernicious anemia, and in those with anaphylactic sensitivity to folic acid.

Iron therapy is contraindicated in patients with hemochromatosis and patients with iron storage disease or the potential for iron storage disease due to chronic hemolytic anemia (e.g., inherited anomalies of hemoglobin structure or synthesis and/or red cell enzyme deficiencies, etc.), pyridoxine responsive anemia, or cirrhosis of the liver.

Cyanocobalamin is contraindicated in patients with sensitivity to cobalt or to cyanocobalamin (vitamin B-12).

Warning

Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. Keep this product out of reach of children. In case of accidental overdose, call a doctor or a Poison Control Center immediately.

WARNINGS/PRECAUTIONS

Vitamin D supplementation should be used with caution in those with hypercalcemia or conditions that may lead to hypercalcemia such as hyperparathryroidism and those who form calcium-containing kidney stones. High doses of vitamin D can lead to elevated levels of calcium that reside in the blood and soft tissues. Bone pain, high blood pressure, formation of kidney stones, renal failure, and increased risk of heart disease can occur.

Prolonged use of iron salts may produce iron storage disease.

Folic acid, especially in doses above 0.1 mg daily, may obscure pernicious anemia, in that hematologic remission may occur while neurological manifestations remain progressive.

The use of folic acid doses above 1 mg daily may precipitate or exacerbate the neurological damage of vitamin B12 deficiency. Consumption of more than 3 grams of omega-3 fatty acids per day from all sources may lead to excessive bleeding.

Supplemental intake of omega-3 fatty acids, such as DHA exceeding 2 grams per day is not recommended.

Avoid overdosage. Keep out of the reach of children.

Drug Interactions

High doses of folic acid may result in decreased serum levels of the anticonvulsant drugs.

Vitamin D supplementation should not be given with large amounts of calcium in those with hypercalcemia or conditions that may lead to hypercalcemia such as hyperparathyroidism and those who form calcium-containing kidney stones.

Zinc can inhibit the absorption of certain antibiotics; take at least 2 hours apart to minimize interactions.

Consult appropriate references for additional specific vitamin-drug interactions.

Information for Patients

Patients should be counseled to disclose all medical conditions, including use of all medications, vitamins and supplements, pregnancy, and breast-feeding.

Pediatric Use

Not for pediatric use.

ADVERSE REACTIONS

Adverse reactions have been reported with specific vitamins and minerals, but generally at doses substantially higher than those in Vitafol [®]-OB+DHA. Allergic reactions have been reported with some forms of gum acacia to include respiratory problems and skin lesions.

Contact your doctor for medical advice about serious adverse events. To report a serious adverse event or obtain product information, contact 1-877-324-9349.

DOSAGE AND ADMINISTRATION

Before, during and after pregnancy, one caplet and one soft-gel capsule daily, or as directed by a physician.

HOW SUPPLIED

Vitafol [®]-OB+DHA is available as a light blue caplet debossed EV0079 and one ambercolored DHA softgel capsule. Available in Box of Unit-Dose pack of 30 (5 child resistant blister cards of 6 caplets and 6 softgel capsules each) (0642-0076-30) and as professional samples (0642-0076-03).

Store at room temperature, approximately 15°-30°C (59°-86°F), avoid excessive heat

and moisture.

*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

Rx

Distributed by:

Exeltis USA, Inc.

Florham Park, NJ 07932

1-877-324-9349

www.exeltisusa.com

©2021 Exeltis USA, Inc.

U.S. Patent No. 6,814,983; 7,390,509

Vitafol[®] is a trademark of Exeltis USA, Inc.

Rev. May 2021

0763001-02

PRINCIPAL DISPLAY PANEL - Kit Carton

0642-0076-30 Vitafol [®]-OB+DHA Prenatal Supplement with DHA SUGAR, LACTOSE, GLUTEN AND IODINE FREE New Smaller DHA Softgel DOES NOT CONTAIN FISH OIL R _X Unit Dose Pack

30 Caplets and 30 Softgel Capsules

U.S. Patented



VITAFOL-OB PLUS DHA PRENATAL SUPPLEMENT PLUS DHA

vitamin a, ascorbic acid, vitamin d, .alpha.-tocopherol, thiamine mononitrate, riboflavin, niacin, pyridoxine hydrochloride, folic acid, cyanocobalamin, calcium, iron, magnesium, zinc, copper, and doconexent kit

Product Information									
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0642-0076						
Deckering									
Packaging									
# Item Code	Package Description	Marketing Start Date	Marketing End Date						

	Quantity of Parts						
	Total Product Qua						
Part 2 1 BOX 30	1						
	30						

Part 1 of 2

VITAFOL-OB

vitamin a, ascorbic acid, vitamin d, .alpha.-tocopherol, thiamine mononitrate, riboflavin, niacin, pyridoxine hydrochloride, folic acid, cyanocobalamin, calcium, iron, magnesium, zinc, and copper tablet, coated

Product Information

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
VITAMIN A (UNII: 81G40H8B0T) (VITAMIN A - UNII:81G40H8B0T)	VITAMIN A	810 ug
ASCORBIC ACID (UNII: PQ6CK8PD0R) (ASCORBIC ACID - UNII:PQ6CK8PD0R)	ASCORBIC ACID	70 mg
VITAMIN D (UNII: 9VU1KI44GP) (CHOLECALCIFEROL - UNII:1C6V77QF41)	VITAMIN D	10 ug
.ALPHATOCOPHEROL (UNII: H4N855PNZ1) (.ALPHATOCOPHEROL - UNII: H4N855PNZ1)	.ALPHATOCOPHEROL	13.5 mg
THIAMINE MONONITRATE (UNII: 8K0I04919X) (THIAMINE ION - UNII:4ABT0J945J)	THIAMINE	1.6 mg
RIBOFLAVIN (UNII: TLM29760FR) (RIBOFLAVIN - UNII:TLM29760FR)	RIBOFLAVIN	1.8 mg
NIACIN (UNII: 2679MF687A) (NIACIN - UNII:2679MF687A)	NIACIN	18 mg
PYRIDOXINE HYDROCHLORIDE (UNII: 68Y4CF58BV) (PYRIDOXINE - UNII:KV2JZ1BI6Z)	PYRIDOXINE HYDROCHLORIDE	2.5 mg
FOLIC ACID (UNII: 935E97BOY8) (FOLIC ACID - UNII:935E97BOY8)	FOLIC ACID	1700 ug
CYANOCOBALAMIN (UNII: P6YC3EG204) (CYANOCOBALAMIN - UNII:P6YC3EG204)	CYANOCOBALAMIN	12 ug
CALCIUM (UNII: SY7Q814VUP) (CALCIUM - UNII:SY7Q814VUP)	CALCIUM	100 mg
IRON (UNII: E1UOL152H7) (IRON - UNII:E1UOL152H7)	IRON	65 mg
MAGNESIUM (UNII: I38ZP9992A) (MAGNESIUM - UNII:I38ZP9992A)	MAGNESIUM	25 mg
ZINC (UNII: J41CSQ7QDS) (ZINC - UNII: J41CSQ7QDS)	ZINC	25 mg
COPPER (UNII: 789U1901C5) (COPPER - UNII:789U1901C5)	COPPER	2 mg

Inactive Ingredients						
Ingredient Name	Strength					
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)						
STEARIC ACID (UNII: 4ELV7Z65AP)						
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)						

GELATIN, UNS	SPECIFIE	D (UNII: 20	6860N327L)					
				E (UNII: L11K75P92J)	1			
SILICON DIO				_ (,				
		-		CINIC ANHYDRIDE) (UN	NII: 461P5CIN6T)		
SUCROSE (UN					, (···· ··· ··· ··· ··· · · · · · · · · ·		
STARCH, COP			51)					
SODIUM ASC			-					
TOCOPHERO								
MAGNESIUM								
TITANIUM DIG								
ACACIA (UNII:			- _ J. /					
			ECIFIED (UN	II: 3WQ0SDW1A)				
MEDIUM-CHA								
SORBIC ACID								
TRICALCIUM	-	• •	K4C08XP666)	1				
HYPROMELLO								
POLYDEXTRO				,				
TRIACETIN (U			-,					
			1600000 W	AMW) (UNII: RFW2E	T671	P)		
FD&C BLUE N				(on 11 12 2	1071	.,		
FD&C BLUE N								
D&C YELLOW	•		• •					
ALUMINUM O	XIDE (UN	III: LMI2606	933)					
Product C	haract		Je	Score			no score	
			/AL				10 score 19mm	
Shape		0	/AL	Size		-		
Flavor				Imprint Code		t	EV0079	
Contains								
Packaging							NAi	
# Item Code		Packa	Package Description		Iv	larketing Start Date	Marketing End Date	
1	30 in 1 Packag		1: Convenie	nce Kit of Co-				
Market		Format	ion					
Marketi Marketi	-			er or Monograp	h	Marketing Start	Marl	ceting End
Catego	ry	, pp i co	Cita		••			Date
unapproved dr other	ug					02/16/2007		
Part 2 o	F2							

DHA

Product Inform a Route of Administr	ation						
	ation						
Route of Administr							
	ration	ORAL					
Active Ingredien	nt/Active	Moiety					
J		dient Name			Basis of St	rength	Strength
DOCONEXENT (UNII: Z	Z AD9OKH9JC) (DOCONEXENT - UNII:ZA	D90KH9JC)		DOCONEXENT		250 mg
Inactive Ingredie	ents						
		Ingredient Name				Str	ength
GLYCERIN (UNII: PDC6							
		86QN327L)					
WATER (UNII: 059QF0k	(OUR)						
Product Charact	teristics						
Color	yellow (A	Amber)	Score			no score	
Shape	OVAL	Size				15mm	
Flavor			Imprint Co	ode			
Contains							
Packaging							
# Item Code	Packa	ge Description	Ma	arketi Da	ng Start Ite		ting End ate
1 30 in 1 Packag		1: Convenience Kit of Co-					
Marketing In	format	ion					
Marketing Category	Applicat	tion Number or Mono Citation		Marketing Start Date			eting End Date
DIETARY SUPPLEMENT			C)2/16/20	007		
Marketing In	format	ion					
Marketing Category	Applicat	tion Number or Mono Citation	graph	Mark	eting Start Date		eting End Date
unapproved drug other			C)2/16/2	007		

Registrant - Exeltis USA, Inc (071170534)

Revised: 12/2022

Exeltis USA, Inc