


HYDROCORTISONE ACETATE AND PRAMOXINE HYDROCHLORIDE- Acella Pharmaceuticals, LLC


Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Hydrocortisone Acetate 2.5%-Pramoxine HCl 1% Cream Rx Only

DESCRIPTION: Hydrocortisone Acetate 2.5%-Pramoxine HCl 1 % Cream is a topical preparation containing Hydrocortisone Acetate 2.5% w/w and Pramoxine HCl 1 % w/w in a hydrophilic base containing purified water, cetostearyl alcohol, mineral oil, white petrolatum, propylparaben, triethanolamine laurel sulfate, citric acid, sodium citrate, isopropyl palmitate, stearic acid, lanolin alcohol, methyl paraben, potassium sorbate, sorbic acid, PEG-40 stearate and ceresin wax. Topical corticosteroids are anti-inflammatory and anti-pruritic agents. The structural formula, the chemical name, molecular formula and molecular weight for active ingredients are presented below.



hydrocortisone acetate
Pregn-4-ene-3,20-dione, 21-(acetyloxy)-11,
17-dihydroxy-, (11-beta)-C₂₂H₃₃O₆; mol. wt. 404.50



pramoxine hydrochloride
4-(3-(n-butylphenoxy)propyl)morpholine hydroxide
C₁₇H₂₇NO₃·HCl; mol. wt. 329.87

CLINICAL PHARMACOLOGY: Topical corticosteroids share anti-inflammatory, anti-pruritic and vasoconstrictive actions.

The mechanism of anti-inflammatory activity of topical corticosteroids is unclear. Various laboratory methods, including vasoconstrictor assays, are used to compare and predict potencies and/or clinical efficacies of the topical corticosteroids. There is some evidence to suggest that a recognizable correlation exists between vasoconstrictor potency and therapeutic efficacy in man.

Pramoxine hydrochloride is a topical anesthetic agent which provides temporary relief from itching and pain. It acts by stabilizing the neuronal membrane of nerve endings with which it comes into contact.

Pharmacokinetics: The extent of percutaneous absorption of topical corticosteroids is determined by many factors including the vehicle, the integrity of the epidermal barrier and the use of occlusive dressings.

Topical corticosteroids can be absorbed from normal intact skin. Inflammation and/or other disease processes in the skin increase percutaneous absorption. Occlusive dressings substantially increase the percutaneous absorption of topical corticosteroids. Thus, occlusive dressings may be a valuable therapeutic adjunct for treatment of resistant dermatoses. (See DOSAGE AND ADMINISTRATION.)

Once absorbed through the skin, topical corticosteroids are handled through pharmacokinetic pathways similar to systemically administered corticosteroids. Corticosteroids are bound to plasma proteins in varying degrees. Corticosteroids are metabolized primarily in the liver and are then excreted by the kidneys. Some of the topical corticosteroids and their metabolites are also excreted into the bile.

INDICATIONS AND USAGE: Topical corticosteroids are indicated for the relief of the inflammatory and pruritic manifestations of corticosteroid-responsive dermatoses.

CONTRAINDICATIONS: Topical corticosteroids are contraindicated in those patients with a history of hypersensitivity to any of the components of the preparation.

PRECAUTIONS: General: Systemic absorption of topical corticosteroids has produced reversible hypothalamic-pituitary-adrenal (HPA) axis suppression, manifestations of Cushing's syndrome, hyperglycemia and glucosuria in some patients. Conditions which augment systemic absorption include the application of the more potent steroids, use over large surface areas, prolonged use and the addition of occlusive dressings. Therefore, patients receiving a large dose of a potent topical steroid applied to a large surface area and under an occlusive dressing should be evaluated periodically for evidence of HPA axis suppression by using the urinary free cortisol and ACTH stimulation tests. If HPA axis suppression is noted, an attempt should be made to withdraw the drug, to reduce the frequency of application or to substitute a less potent steroid. Recovery of HPA axis function is generally prompt and complete upon discontinuation of the drug. Infrequently, signs and symptoms of steroid withdrawal may occur, requiring supplemental systemic corticosteroids. Children may absorb proportionally larger amounts of topical corticosteroids and thus be more susceptible to systemic toxicity. (See Precautions-Pediatric Use.)

If irritation develops, topical corticosteroids should be discontinued and appropriate therapy instituted. In the presence of dermatological infections, the use of an appropriate antifungal or antibacterial agent should be instituted. If a favorable response does not occur promptly, the corticosteroid should be discontinued until the infection has been adequately controlled.

Information for the patient: Patients using topical corticosteroids should receive the following information and instructions:

1. This medication is to be used as directed by the physician. It is for external use only. Avoid contact with eyes.
2. Patients should be advised not to use this medication for any disorder other than for which it was prescribed.
3. The treated skin area should not be bandaged or otherwise covered or wrapped as to be occlusive unless directed by the physician.
4. Patients should report any signs of local adverse reactions especially under occlusive dressings.
5. Parents of pediatric patients should be advised not to use tight-fitting diapers or plastic pants on a child being treated in the diaper area, as these garments may constitute occlusive dressings.

Laboratory Tests: Urinary free cortisol test and ACTH stimulation test may be helpful in evaluating the HPA axis suppression.

Carcinogenesis, Mutagenesis and Impairment of Fertility: Long-term animal studies have not been performed to evaluate the carcinogenic potential or the effect on fertility of topical corticosteroids. Studies to determine mutagenicity with prednisolone and hydrocortisone have revealed negative results.

Pregnancy: Teratogenic Effects: Pregnancy Category C: Corticosteroids are generally teratogenic in laboratory animals when administered systemically at relatively low dosage levels. The more potent corticosteroids have been shown to be teratogenic after dermal application in laboratory animals. There

are no adequate and well-controlled studies in pregnant women on teratogenic effects from topically applied corticosteroids. Therefore, topical corticosteroids should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. Drugs of this class should not be used extensively on pregnant patients, in large amounts or for prolonged periods of time.

Nursing Mothers: It is not known whether topical administration of corticosteroids could result in sufficient systemic absorption to produce detectable amounts in breast milk. Systemically administered corticosteroids are secreted into breast milk in quantities NOT likely to have a deleterious effect on the infant. Nevertheless, caution should be exercised when topical corticosteroids are administered to a nursing woman.

Pediatric Use: Pediatric patients may demonstrate greater susceptibility to topical corticosteroids induced HPA axis suppression and Cushing's syndrome than mature patients because of larger skin surface area to body weight ratio. Hypothalamic-pituitary-adrenal (HPA) axis suppression, Cushing's syndrome and intracranial hypertension have been reported in children receiving topical corticosteroids. Manifestations of adrenal suppression in children include linear growth retardation, delayed weight gain, low plasma cortisol levels and absence of response to ACTH stimulation. Manifestations of intracranial hypertension include bulging fontanelles, headaches and bilateral papilledema. Administration of topical corticosteroids to children should be limited to the least amount compatible with an effective therapeutic regimen. Chronic corticosteroid therapy may interfere with the growth and development of children.

ADVERSE REACTIONS: The following local adverse reactions are reported infrequently with topical corticosteroids, but may occur more frequently with the use of occlusive dressings. These reactions are listed in an approximate decreasing order of occurrence: burning, itching, irritation, dryness, folliculitis, hypertrichosis, acneiform eruptions, hypopigmentation, perioral dermatitis, allergic contact dermatitis, maceration of the skin, secondary infection, skin atrophy, striae, miliaria.

OVERDOSAGE: Topically applied corticosteroids can be absorbed in sufficient amounts to produce systemic effects (See PRECAUTIONS.)

DOSAGE AND ADMINISTRATION: Topical corticosteroids are generally applied to the affected area as a thin film three to four times daily depending on the severity of the condition. Occlusive dressings may be used for the management of psoriasis or recalcitrant conditions. If an infection develops, the use of occlusive dressings should be discontinued and appropriate antimicrobial therapy instituted.

HOW SUPPLIED:

Hydrocortisone Acetate 2.5%-Pramoxine HCl 1 % Cream
1 ounce tube (NDC 42192-107-01) 30 x 4 gram tubes (NDC 42192-108-04)

Hydrocortisone Acetate 2.5%-Pramoxine HCl 1 % Cream Kit with Pramoxine Wipes
1 ounce tube (NDC 42192-118-07) 30 x 4 gram tubes (NDC 42192-119-30)

Storage Conditions: Store at 20° - 25°C (68° - 77°F); excursions permitted to 15° - 30°C (59° - 86°F) [see USP Controlled Room Temperature].

All prescription substitutions and / or recommendations using this product shall be made subject to state and federal statutes as applicable. **Please NOTE: This is not an Orange Book product and has not been subjected to FDA therapeutic equivalency or other equivalency testing.** No representation is made as to generic status or bioequivalency. Each person recommending a prescription substitution using this product shall make such recommendation based on his / her professional knowledge and opinion, upon evaluating the active ingredients, inactive ingredients, excipients and chemical information provided herein."

Manufactured For:

Acella Pharmaceuticals, LLC • Alpharetta, GA 30022

PRINCIPAL DISPLAY PANEL - Kit

NDC 42192-118-07

**Hydrocortisone Acetate 2.5% -
Pramoxine HCl 1% Cream Kit**

Packaged together for convenience

**Hydrocortisone Acetate 2.5% - Pramoxine HCl 1% Cream
and
Pramoxine HCl 1% Wipes**

Rx Only


Contents: Hydrocortisone Acetate 2.5% - Pramoxine HCl 1% Cream

Net Wt. 1 oz. tube and applicator

6 Pramoxine HCl 1% Pre-moistened Anorectal Wipes

CONTENTS NOT FOR INDIVIDUAL SALE

**Acella
PHARMACEUTICALS, LLC**



hydrocortisone acetate and pramoxine hydrochloride kit

Product Information

Product Type	HUMAN PRESCRIPTION DRUG LABEL	Item Code (Source)	NDC:42192-118
---------------------	-------------------------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:42192-118-07	1 in 1 CARTON		

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 TUBE, WITH APPLICATOR	28 g
Part 2	1 PACKAGE, COMBINATION	1 g

Part 1 of 2

HYDROCORTISONE ACETATE AND PRAMOXINE HYDROCHLORIDE

hydrocortisone acetate and pramoxine hydrochloride cream

Product Information

Route of Administration	TOPICAL	DEA Schedule	
--------------------------------	---------	---------------------	--

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HYDROCORTISONE ACETATE (HYDROCORTISONE)	HYDROCORTISONE ACETATE	2.5 g in 100 g
PRAMOXINE HYDROCHLORIDE (PRAMOXINE)	PRAMOXINE HYDROCHLORIDE	1 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
WATER	
CETOSTEARYL ALCOHOL	
MINERAL OIL	
PETROLATUM	
PROPYLPARABEN	
TRIETHANOLAMINE LAURYL SULFATE	
CITRIC ACID MONOHYDRATE	
SODIUM CITRATE	
ISOPROPYL PALMITATE	
STEARIC ACID	
LANOLIN ALCOHOLS	
METHYLPARABEN	

POTASSIUM SORBATE	
SORBIC ACID	
PEG-40 STEARATE	
CERESIN	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		28 g in 1 TUBE, WITH APPLICATOR		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
UNAPPROVED DRUG OTHER		02/01/2010	

Part 2 of 2

INERT

inert swab

Product Information

Route of Administration	TOPICAL	DEA Schedule	
-------------------------	---------	--------------	--

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PRAMO XINE HYDRO CHLORIDE (PRAMO XINE)	PRAMO XINE HYDROCHLORIDE	1 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN	
WATER	
CITRIC ACID MONOHYDRATE	
SODIUM CITRATE	
POTASSIUM SORBATE	
PHENOXYETHANOL	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		1 g in 1 PACKAGE, COMBINATION		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
UNAPPROVED DRUG OTHER		02/01/2010	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
UNAPPROVED DRUG OTHER		02/01/2010	

Labeler - Acella Pharmaceuticals, LLC (825380939)

Registrant - Acella Pharmaceuticals, LLC (825380939)

Revised: 9/2014

Acella Pharmaceuticals, LLC