

**GOOD SENSE TUSSIN CF- dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride solution**

**L. Perrigo Company**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Perrigo Tussin CF Drug Facts**

**Active ingredients (in each 10 mL)**

Dextromethorphan HBr, USP 20 mg

Guaifenesin, USP 200 mg

Phenylephrine HCl, USP 10 mg

**Purposes**

Cough suppressant

Expectorant

Nasal decongestant

**Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
- nasal congestion
- cough due to minor throat and bronchial irritation

**Warnings**

**Do not use**

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

**When using this product**

do not use more than directed

**Stop use and ask a doctor if**

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	10 mL every 4 hours
children under 12 years	do not use

**Other information**

- each 10 mL contains: sodium 5 mg
- store at 20-25°C (68-77°F).  
Do not refrigerate.

**Inactive ingredients**

anhydrous citric acid, edetate disodium, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

**Questions or comments?**

1-800-719-9260

**Principal Display Panel**

Non-Drowsy

Tussin CF

Cough Suppressant (Dextromethorphan HBr)

Expectorant (Guaifenesin)

Nasal Decongestant (Phenylephrine HCl)

Multi-Symptom Cold

Relieves:

Nasal Congestion

Cough

Mucus

Peak Cold

Adult

For Ages 12 & Over

Compare to active ingredients of Robitussin® Multi-Symptom Cold

4 FL OZ (118 mL)

NDC 0113-0703-26

GOODSENSE®

Non-Drowsy

**Tussin**  
**CF** Cough Suppressant  
(Dextromethorphan HBr)  
Expectorant (Guaifenesin)  
Nasal Decongestant  
(Phenylephrine HCl)

**Multi-Symptom Cold**

Relieves:

- Nasal Congestion
- Cough • Mucus



Peak Cold

**Adult**  
For Ages 12 & Over

Compare to active ingredients of  
Robitussin® Multi-Symptom  
Cold



4 FL OZ (118 mL)

NDC 0113-0703-26

GOODSENSE®

Non-Drowsy

**Tussin**  
**CF** Cough Suppressant  
(Dextromethorphan HBr)  
Expectorant (Guaifenesin)  
Nasal Decongestant  
(Phenylephrine HCl)

**Multi-Symptom Cold**

Relieves:

- Nasal Congestion
- Cough • Mucus




Peak Cold

**Adult**  
For Ages 12 & Over

Compare to active ingredients of  
Robitussin® Multi-Symptom  
Cold



4 FL OZ (118 mL)



**CF Multi-Symptom Cold**  
**Tussin**

**GOODSENSE**

## Drug Facts

### Active ingredients (in each 10 mL)

Dextromethorphan HBr,  
USP 20 mg.....Cough suppressant  
Guaifenesin, USP 200 mg.....Expectorant  
Phenylephrine HCl,  
USP 10 mg.....Nasal decongestant

### Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
  - nasal congestion
  - cough due to minor throat and bronchial irritation

### Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

When using this product do not use more than directed

### Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition. ►

## Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	10 mL every 4 hours
children under 12 years	do not use

### Other information

- each 10 mL contains: sodium 5 mg
  - store at 20-25°C (68-77°F).
- Do not refrigerate.

**Inactive ingredients** anhydrous citric acid, edetate disodium, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

### Questions or comments?

1-800-719-9260

Gluten Free

Dosage Cup Included

**DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING**

Distributed By **Perrigo**® Allegan, MI 49010

**PARENTS:**

Learn about teen medicine abuse

[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)



LOT NO.

EXP.

: 51626 C2 C7

GOOD SENSE TUSSIN CF

dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride solution

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0113-0703
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 10 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 10 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 10 mL

Inactive Ingredients	
Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics				
Color		RED (clear)	Score	
Shape			Size	
Flavor		FRUIT	Imprint Code	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0113-0703-26	1 in 1 CARTON	07/29/2014	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0113-0703-34	1 in 1 CARTON	07/29/2014	
2		237 mL in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final		part341	07/29/2014	

**Labeler** - L. Perrigo Company (006013346)