

**BISACODYL- bisacodyl suppository**  
**Geritrex LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Bisacodyl Suppositories**

**Drug Facts**

Active ingredient (in each suppository) Purpose

Bisacodyl USP 10 mg ..... Stimulant laxative

**Uses**

relieves occasional constipation.

this product generally produces a bowel movement in 15 minutes to 1 hour.

**Directions**

tear off one suppository from the strip.

tear and peel the wrapper in downward motion.

continue tearing downward to almost the full length of the suppository (see illustration at the right)

gently remove the suppository from the wrapper.

lie on your side and push suppository. with pointed end first, high into the rectum so it will not slip out

retain it for 15 to 20 minutes.

Do not exceed recommended dose

if suppository seems soft, place in refrigerator for a short time before use

Adults and children 12 years and over 1 suppository once daily.

Children under 12 years Ask a doctor.

Adults and children 12 years and over 1 suppository once daily.

Children under 12 years Ask a doctor.

**WARNINGS**

For rectal use only.

Do not use when abdominal pain, nausea, or vomiting are present.

Ask a doctor before use if you have a sudden change in bowel habits that lasts over a period of 2 weeks.

When using this product you may experience. rectal burning. mild cramps.abdominal discomfort. faintness.

stop use and ask a doctor if you have rectal bleeding. you fail to have a bowel movement after using a laxative. These may be signs of a serious condition.

you need to use a laxative for more than one week.

if pregnant or breast-feeding, ask a health professional before use.

**INACTIVE INGREDIENT**


hydrogenated vegetable oil

## STORAGE

store at temperatures below 30°C (86°F)

## OTC - KEEP OUT OF REACH OF CHILDREN

If swallowed, get medical help or contact a poison control center right away.



<b>BISACODYL</b>			
bisacodyl suppository			
<b>Product Information</b>			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:54162-006
Route of Administration	RECTAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
BISACODYL (UNII: 10 X0 709 Y6 I) (BISACODYL - UNII:10 X0 709 Y6 I)	BISACODYL	0.01 g in 2 g

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
HYDROGENATED SOYBEAN OIL (UNII: A2M9 1M9 18 C)	

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:54162-006-01	2 g in 1 BOX, UNIT-DOSE; Type 0: Not a Combination Product		

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC monograph not final	part334	07/31/2015	

**Labeler** - Geritrex LLC (112796248)

Revised: 1/2016

Geritrex LLC